



# Selected Plants of Navajo Rangelands

## with Navajo names

---

**Diné bikéyah Chi'I noosélgíí Baḡhą́ą́nosin** | Take care of our Navajo Rangelands

Developed by Navajo Tri-State Federally-Recognized Tribes Extension  
Program in collaboration with New Mexico State University

For spoken audio and additional content, visit [NavajoRange.nmsu.edu](https://NavajoRange.nmsu.edu)



**Cover image:**

Top (left to right): Indian ricegrass (Amy Smith Muese), cliffrose (Gerald Moore), black grama (Patrick Alexander), western wheatgrass (Patrick Alexander)

Bottom: Navajo Nation flag



For spoken audio and additional content, or to download this booklet to a device or computer, visit **NavajoRange.nmsu.edu**

You can help document Navajo rangelands for future generations. Send photos of rangeland sites to: Gerald Moore (germoore@nmsu.edu), Navajo Tri-State Federally-Recognized Tribes Extension Program (FRTEP), Grey Farrell (gfarrell@cals.arizona.edu), University of Arizona Extension, Tuba City, AZ, or Treva Henio (tjhenio@gmail.com), BIA-Natural Resources, Crownpoint, NM, labeled with date, location, and your comments.

This book can help agricultural professionals, ranchers, and others on the Navajo Nation to identify typical range plants and maximize productivity and sustainability of rangeland ecology. Once downloaded to a device, it can be used even in areas without mobile or Wi-Fi coverage. Plants are arranged by plant type. Use the index to find plants sorted by common name, scientific name, habitat, growing season, flower color, or livestock special considerations. Plant scientific names were drawn primarily from the USDA PLANTS database.

Produced by the Department of Innovative Media Research and Extension, New Mexico State University. Send comments or corrections to mmedia@nmsu.edu. © 2017, 2018

New Mexico State University Board of Regents. This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, through the Western Sustainable Agriculture Research and Education program Professional Development Grant Program under project number EW15-023. USDA is an equal opportunity employer and service provider. NMSU is an equal opportunity employer and educator.



United States Department of Agriculture  
National Institute of Food and Agriculture



**BE BOLD.** Shape the Future.  
**College of Agricultural, Consumer  
and Environmental Sciences**  
Cooperative Extension Service



# Table of Contents

<b>Does the Range Concern You?</b>	<b>6</b>
<b>Rangeland Management Tips</b>	<b>7</b>
<b>Plant Types</b>	<b>9</b>

## Grasses

Bluegrass, Kentucky	10
Bluestem, Little False	12
Bottlebrush Squirreltail	13
<b>Bromes</b>	
Brome, Field	14
Brome, Japanese	16
Brome, Mountain	18
Brome, Red	20
Brome, Ripgut	22
Brome, Smooth	24
Cheatgrass	26
Cordgrass, Alkali	28
<b>Dropseeds</b>	
Dropseed, Giant	30
Dropseed, Pine	32
Dropseed, Sand	34
Dropseed, Spike	36
Sacaton, Alkali	38
<b>Fescues</b>	
Fescue, Arizona	40
Fescue, Six Weeks	42
Foxtail Barley	44
Galleta, James'	46
Goatgrass, Jointed	48
<b>Gramas</b>	
Grama, Black	50
Grama, Blue	52
Grama, Sideoats	54
Johnsongrass	56
Junegrass, Prairie	58
<b>Muhlys</b>	
Muhly, Mountain	60
Muhly, Ring	62

## Brush

Algerita	110
Antelope Bitterbrush	112

<b>Muhly</b> <i>continued</i>	
Muhly, Sandhill	64
Muhly, Spike	66
Mutton Grass	68
Needle-and- Threadgrass	69
<b>Needlegrasses</b>	
Needlegrass, Desert	71
Needlegrass, Scribner	73
New Mexico Feathergrass	75
Redtop	77
Ricegrass, Indian	79
Saltgrass	81
Sandbur, Field	83
<b>Three-Awns</b>	
Three-Awns (Annual)	85
Three-Awns (Perennial)	87
Tufted Horse Grass	89
<b>Wheatgrasses</b>	
Wheatgrass, Crested	91
Wheatgrass, Slender	93
Wheatgrass, Western	95
<b>Wildrye</b>	
Wildrye, Canadian	97
Wildrye, Russian	99
<b>Grasslike</b>	
Arrowgrass	101
<b>Rushes</b>	
Bulrush	102
Rushes	104
<b>Sedges</b>	
Nutsedge, Yellow	106
Sedges	108

Biscuitroot	114
Blackbrush	116

## Table of Contents

*continued*

### Brush *continued*

Buckwheat	118	Sagebrush, Bigelow	146
Camelthorn	121	<b>Sagebrushes</b>	
Chokecherry	123	Sagebrush, Black	148
Cliff Fendlerbush	125	Sagebrush, Fringed	150
Cliffrose	127	Sagebrush, Sand	152
Four-Winged Saltbush	128	Shadscale	154
Greasewood	130	Skunkbush	156
Mormon Tea	132	Snakeweed	158
Oregon Grape	134	Snowberry	160
Pingue	136	Wax Currant	162
Rabbitbrush	138	Wild Rose	164
Rabbitbrush, Green	140	Winterfat	166
<b>Rabbitbrushes</b>		Wolfberry	168
Roundleaf Buffaloberry	142	Field Bindweed	170
Sagebrush, Big	144	Puncturevine	171

### Vines

Antelopehorn	173	Beeplant, Rocky Mountain	175
--------------	-----	--------------------------	-----

### Forbs

Birdbeak	177	Globemallow	219
Bitterweed	179	Groundsel, Threadleaf	221
Blanket Flower	181	Halogeton	223
Blue-Eyed Grass	183	Horsemint	225
Bluebells	185	Indian Paintbrush	227
Bundle Flower	187	Iris	229
California Burclover	189	Jacob's Ladder	231
Cocklebur	191	Jimmyweed	233
Common Purslane	193	Jimsonweed	235
Copperweed	195	<b>Knapweeds</b>	
Cow Parsnip	196	Knapweed	237
Curlycup Gumweed	198	Knapweed, Diffuse	239
Daisy, Rock	200	Knapweed, Russian	241
Dandelion	202	Kochia	243
Death Camas	204	Lambsquarters	245
Desert Sand Verbena	206	Larkspur	247
Devilweed	207	Leafy Spurge	249
Evening Primrose	209	<b>Locoweeds / Milkvetches</b>	
Filaree	211	Loco, Ellis	251
Four O'Clock	213	Loco, Rattleweed	253
Geranium	215	Loco, Woolly	255
Gilia	217	Red Stem Peavine	257

## Table of Contents

*continued*

### Forbs *continued*

Lupine	259
Milkweed	261
Mullein	263
Mustard	265
Navajo Tea	267
Navajo Tobacco	269
Navajo Wild Carrot	271
Nightshade	273
Penstemon	275
Pepperweed	277
Poison Ivy	279
Rubberweed, Bitter	281
Sego Lily	283
Senecio	285
Sneezeweed	287

### Cacti

Cholla	313
Prickly Pear	315

### Trees

Alder	319
Colorado Piñon	321
Coyote Willow	323
Fremont Cottonwood	325
Gambel Oak	327
Juniper	328
Mountain Mahogany	330
Ponderosa Pine	332
Quaking Aspen	334

### Other

<b>Horsetails</b>	
Horsetail	352
Horsetail, Smooth	354

Spreading Wallflower	289
St. Johnswort	291
Sulphur Cinquefoil	293
Sunflower	295
Thistle, Musk	297
Thistle, Canada	299
Thistle, Scotch	301
Tumbleweed	303
<b>Thistles</b>	
Water Hemlock	305
Western Yarrow	307
Whitetop	308
Wooly Indian Wheat	310
Wormsweed	312

Yucca	317
-------	-----

Red Plum	336
Russian Olive	338
Saltcedar	340
Siberian Elm	342
Subalpine Fir	344
Tree of Heaven	346
Utah Serviceberry	348
Water Birch	350

Lichen	356
--------	-----

<b>Acknowledgments</b>	<b>358</b>
------------------------	------------

<b>Credits</b>	<b>359</b>
----------------	------------

<b>Glossary</b>	<b>361</b>
-----------------	------------

<b>Index</b>	<b>365</b>
--------------	------------



# Does the Range Concern You?

Yes, each one of us has a real interest in our rangeland. It is the main source of our meat supply. It is important in the production of wool and leather products. It is a valuable source of water, clean air, and wildlife. It offers many types of recreation. It is just as important for the non-user of rangeland to promote its conservation and protection for the future as it is for the person raising livestock and living on the range.

---

## Know Your Range!

You can keep the range healthy and productive by knowing the plants and animals that live there and how to manage them properly. Plants and animals live together as a **community**. When livestock or game animals graze an area, changes take place in the plant community. To be a good range manager, you must recognize the changes that result from grazing. You must be able to tell why the changes took place and whether the changes are good or bad. To know about the rangelands and how to manage and care for them, you must have some knowledge of soils, water, climate, and plants.

Plants will tell you what kind of range you have and how good a manager you are. Each plant will tell you a story of what is happening. The presence or absence of certain plants tells the condition of the range, how the range is being used, and what needs to be done to improve and protect the range. This plant identification tool provides a selection of the most common of hundreds of range plants that grow on the Navajo Reservation.

This tool is intended to be of assistance to Navajos in knowing both Navajo and English names for the plants they encounter and also as an aid to those working with Navajos. In discussing range or livestock, it is often necessary to refer to a specific plant by name. This tool should provide a means of knowing some of these names in Navajo so the Navajos involved will know exactly which plant is being referred to. This is strictly a preliminary list. Many of the plants will be known by other names, either Navajo or English, in other areas.

## For More Information

Information in this handbook is based on the Selected Plants of Navajo Rangelands database: [navajorange.nmsu.edu](http://navajorange.nmsu.edu). This version is intended as a downloadable version of the database for offline use. Please visit the site for more information.

“Does the Range Concern You?” was adapted with permission from the 1981 *Navajo Nation range management handbook*, by Frank Parrill, then Range Conservationist, Navajo Tribe, Window Rock, AZ, and Allan Blacksheep, Jr., then Agricultural Extension Agent, University of Arizona, Ft. Defiance, AZ. Their handbook also formed the basis of the plant list on which this tool is based. Without their original work, our task would have been much more difficult.

# Rangeland Management Tips

from range professionals in the region

---

## Gerald Moore

Navajo Tri-State Federally-Recognized Tribes Extension  
Coordinating Extension Agent

1. A rancher's knowledge of livestock is incomplete without a knowledge of the types of range plants they graze.
2. Remember how important the range is to livestock. Don't concentrate on livestock first; concentrate on the land first.
3. The land is your bank. If you contribute to it, it will provide for you in the future and provide a future for the next generations.
4. In times when there's less grass, it's important to control where your herd goes. Reduction is always a management option.
5. The success and productivity of a livestock operation depends largely on the productivity and management of your natural resources.

## Renee Benally

Western Navajo Agency, Bureau of Indian Affairs (BIA)  
Natural Resource Specialist

Animals (Nal golshi), plants (Ch'il), and soil (Leezh) all play a part in rangeland grazing. To achieve ecological and economic objectives, ranchers and land managers must first understand how plants grow and reproduce, and then how grazing animals affect plants through defoliation, hoof action, and other factors.

Also important are grazing animals' specific behaviors and nutritional needs at critical points of the production cycle.

With this understanding in hand, ranch and land managers can consider:

1. Timing (When should grazing occur?)
2. Frequency (How often should defoliation occur?)
3. Intensity (How much forage should be removed, or how much residual plant material should remain after grazing?)
4. Stocking rate and type (How many and what type of livestock should be grazed?)

Plants respond differently to grazing depending on its timing, intensity, and frequency as well as the physiological and morphological characteristics of the plants and how these affect plant growth before and after grazing events.

## Nick Ashcroft

New Mexico State University  
Extension Range Specialist

1. Watch the plants and not the animals. You're not ranchers first, you're grass farmers first.
2. Range management means allowing the plants some rest, not over utilizing.
3. When identifying plants, the first question is:
  - What is it?
  - Is it poisonous to my animals?
  - Is it noxious/invasive?


To prevent noxious/invasive plants from taking over, take care of perennial plants by not over utilizing them.

4. Grass plants need to replace 30% of their roots every year. If you have a drought year, or if you over utilize, that root system isn't replaced, and it can't continue.
5. Remember the basics of what a plant needs: sunlight, air, soil/nutrients, water. If you take off too much leaf area by grazing, the plant quits photosynthesizing and quits growing. It's also important to leave enough plant there to put the water into the soil instead of letting it run off.
6. Every site is different, with different precipitation and different challenges. On the Navajo Nation one unique challenge is that there are no fences. That means animals are moving where the feed is, and you have to keep an even closer eye on the plants to see what is happening to them.



# Plant Types

## Grasses




Grasses have a main stem with a sheath and leaves wrapping around the stem below. Their leaves have parallel veins. Growth pattern is alternate (leaves staggered on the stem) or basal.

## Brush




Also called shrubs, plants in this category are woody and are usually multi stemmed and less than 16 feet tall (with some exceptions). They have shallower roots than trees.

## Vines




Twining or climbing plants with relatively long stems, vines grow on something, attached with tendrils. They may have prominent flowers and can be woody or herbaceous.

## Forbs



Often called "weeds," forbs usually have non woody stems (easily bent) and prominent flowers. They may be annual, biennial, or perennial. They can have single or multiple stems or grow in a basal pattern, meaning all growth comes from the base.

## Cacti



Stems of cacti are thick and succulent, sometimes forming pads. Leaves are reduced to spines and barbs, grouped in clusters on the stem. Different species have many differently colored flowers.

## Trees



Trees are woody and have bark. They usually have a single trunk (with some exceptions) and taproots. Different species have different shapes and kinds of leaves, such as needles, evergreen leaves, or deciduous leaves.



Patrick Alexander

# Bluegrass, Kentucky

*Poa pratensis*

## Tsétahtł'oh



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Cool season

### Special Considerations:

- ✓ Good livestock forage
- ! Potentially invasive

**Kentucky bluegrass** is a perennial, sod-forming grass native to Europe. The species is highly palatable to horses, cattle, and sheep. The plant becomes dormant during the heat of summer but regains or maintains its green color in fall.

This plant may become weedy or invasive in some regions or habitats and may displace desirable vegetation if not properly managed.

## Grasses

### Bluegrass, Kentucky

*Poa pratensis*

Tsétaht'oh



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander





Patrick Alexander

# Bluestem, Little False

*Schizachyrium scoparium* var. *scoparium*

Tł'ohdeitł'ichi'í



## Alternative Names:

Beard grass

## Plant Type:

Grass

## Habitat:

Grasslands

Ponderosa pine/mixed conifer

## Growing Season:

Warm season

## Special Considerations:

None

**Little bluestem** is a medium height grass with coarse stems and basal leaves. Plants are green, but often purplish at base of stem. The entire plant has a reddish cast after frost.

Little bluestem can be a dominant understory species in open-canopy pine stands. Seed is generally wind-dispersed only short distances from the parent plant.



Patrick Alexander



Patrick Alexander



Max Licher

# Bottlebrush Squirreltail

*Elymus elymoides*

**Azéé'ilwo'iindtsaaí**



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Brushlands

Grasslands

Wastelands

## Growing Season:

Cool season

## Special Considerations:

- ✓ Provides good forage

**Bottlebrush squirreltail** is a, native perennial bunchgrass. The plant grows 8 to 25 inches tall, and occurs primarily on dry, gravelly, or saline soils. It is common on hillsides and alkaline flats.

Considered good forage before seed development and after seed shatter, it receives its name because the seedhead resembles a bottlebrush.



Patrick Alexander



Bildoj [CC BY-SA 3.0], from  
Wikimedia Commons

# Brome, Field

*Bromus arvensis*



## Alternative Names:

*Cenchrus incertus*

## Plant Type:

Grass

## Habitat:

Brushlands

Grasslands

River valleys

## Growing Season:

Cool season

## Special Considerations:

! Noxious

**Field brome** is a winter annual that produces dense, low, leafy growth in the fall and starts growing early in the spring. It does not have creeping stolons or rhizomes, but tillers profusely. It produces seedheads in late spring or early summer. Seed stalks are 2 to 3 feet tall. It has an extensive fibrous root system and relatively short top growth.

It grows vigorously under high fertility conditions and often smothers other grasses or weeds. It is an excellent seed producer and can maintain itself as a reseeding annual.



# **Brome, Field** *Bromus arvensis*



Robert H. Mohlenbrock, Courtesy of USDA NRCS  
Wetland Science Institute, from Wikimedia Commons

Seedheads



Bildoj [GFDL or CC BY-SA 3.0], from  
Wikimedia Commons

Seeds



Daderot [CC0 or CC0], from Wikimedia Commons

Early growth habit



Russ Kleinman, Bill Norris, &  
Richard Felger

# Brome, Japanese

*Bromus japonicus*



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Grasslands

Wastelands

## Growing Season:

Cool season

## Special Considerations:

! Noxious

**Japanese brome** is an introduced, annual grass from 8 to 48 inches tall. Its sheaths are hairy; the blades are glabrous to hairy. The inflorescence is an open panicle bearing 6 to 13 caryopses. Roots may be shallow, or surprisingly deep for an annual. Japanese brome has florets with somewhat flattened lemmas with awns mostly  $\frac{1}{4}$  to  $\frac{1}{3}$  inch long. It is found in sandy areas with abundant moisture. The stems and leaves are covered in dense, white pubescence.



**Brome, Japanese**

*Bromus japonicus*



*Russ Kleinman, Bill Norris, & Richard Felger*



*Patrick Alexander*



*Russ Kleinman, Bill Norris, & Richard Felger*



*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*



Russ Kleinman &amp; Karen Blisard

# Brome, Mountain

*Bromus carinatus*

Tł'ohntsohndtł'zí



## Alternative Names:

California brome

## Plant Type:

Grass

## Habitat:

Grasslands

## Growing Season:

Cool season

## Special Considerations:

- ✓ Good spring forage

**Mountain brome** is a native, annual or perennial bunchgrass that lives three to five years and grows to be two to four feet tall. The roots of mountain brome are fibrous, grow very quickly, and become deep and widespread. Young plants are erect, but older stems are decumbent: they grow along the ground with only the apical tips remaining erect.. Stems are robust with hairy sheaths. Leaf blades are  $\frac{1}{16}$  to  $\frac{1}{2}$  inch wide and 6 to 12 inches long. They can be pubescent or glabrous. The inflorescence is a stiff, open panicle, 4 to 8 inches long, and it droops at maturity. The spikelets are flattened, measuring  $\frac{3}{4}$  to  $1\frac{1}{2}$  inches long and  $\frac{1}{5}$  to  $\frac{1}{4}$  inch wide. They have five to seven flowers. Lemmas are over  $\frac{3}{8}$  inch long, flattened, keeled and usually pubescent. The awns are  $\frac{2}{16}$  to  $\frac{5}{16}$  inches long. Seeds mature in May and June at low elevations and by late August at high elevations.

Mountain brome is common in the mountains and foothills of the Intermountain West. It is often found in relatively moist habitats in mountain big sagebrush, mountain shrub, aspen, and spruce–fir communities and up to sub alpine mountain meadows. It can be found at elevations ranging from 5,000 to 10,500 feet.

Mountain brome is an excellent plant for the revegetation of livestock and big game ranges in foothill and mountain locations. It is highly palatable in the spring, providing good forage for wildlife and livestock. The leaves provide excellent grazing for elk, cattle, and horses and are also eaten by sheep and deer. The seeds are readily eaten by small mammals and birds.

**Brome, Mountain**

*Bromus carinatus*

**Tł'ohdsohndtł'zí**



Russ Kleinman



Russ Kleinman & Karen Blisard



Russ Kleinman



Russ Kleinman





Ken Gishi

# Brome, Red

*Bromus rubens*

## Bi'zé yilwo' lichi'í



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

! Noxious

! Can be harmful to livestock

**Red brome** is native to southern Europe and is an invasive weed in desert shrub and annual grassland communities. It commonly occurs in small patches on shallow soils, growing best where there is little competition from other annuals.

A tufted, annual bunchgrass, it grows from 8 to 20 inches tall.

Often found in areas with relatively high levels of sulfur dioxide pollution, it is of little value to livestock and big game. The long awns of red brome are harmful to livestock when seeds are ripe. Under some circumstances, red brome may provide a source of forage.

**Brome, Red**

*Bromus rubens*

**Bi'zé yilwo' lichí'í**



Patrick Alexander



Patrick Alexander



Ken Gishi





Russ Kleinman

# Brome, Ripgut

*Bromus diandrus*

## Tł'oh da a gighí



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

- ! Can injure livestock and pets

**Ripgut brome** is easily recognized by its stiff awns measuring  $1\frac{1}{4}$  to  $2\frac{3}{8}$  inches, carried in an open and drooping panicle. It grows around buildings and in other disturbed areas. It burns easily when dry. When flowering, its seedheads have tiny, rough teeth that can injure livestock.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*

**Brome, Ripgut**

*Bromus diandrus*


**Tł'oh da a gighí**



Russ Kleinman



Joseph M. DiTomaso, University of California - Davis



Barry Rice, sarracenia.com, Bugwood.org

Prominent awns



Copyright © 2007 The Regents of the University of California. All rights reserved. 5387158

Joseph M. DiTomaso, University of California - Davis

Panicle



Patrick Alexander

# Brome, Smooth

*Bromus inermis*

**bį́įh tł'óh**



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Grasslands

## Growing Season:

Cool season

## Special Considerations:

✓ Invasive

! Provides good forage

**Smooth brome** is a leafy, sod-forming, perennial grass that spreads by rhizomes. It is native in some areas of the United States, and provides good erosion control in areas that receive over 20 inches of rain per year.

The grass is highly palatable, is high in protein content, and relatively low in crude fiber content, making it a good forage species for livestock. The stems vary in height from 2 to 4 feet. The plant produces numerous basal and stem leaves that vary in length from 4 to 10 inches. Frequently the leaves are marked by a transverse wrinkle resembling a “W” a short distance below the tip. The flowerhead develops a characteristic rich purplish-brown color when mature. The seed is produced in semi-compact 5-inch-long panicles with ascending branches. The flat, compressed seed is usually awnless, about ⅓ inch long, and smooth.



## Grasses

### Brome, smooth

*Bromus inermis*

bijh t'óh



Patrick Alexander




Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Steve Perkins

# Cheatgrass

*Bromus tectorum*

## Shíyínáldzidí



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Cool season

### Special Considerations:

! Invasive

**Cheatgrass** is an invasive cool-season annual grass from the Mediterranean. It has solitary stems 4 to 24 inches tall, with soft drooping seedheads. Cheatgrass can provide good forage for a short time, but matures quickly and becomes unpalatable. At maturity, the plant turns reddish and cures to a light tan. After curing, late spring cheatgrass becomes highly flammable and has changed the fire frequency of western rangelands.

In Europe, cheatgrass historically grew in the decaying straw of thatched roofs. “Tectum” is Latin for roof, hence the name *Bromus tectorum*, “brome of the roofs.” Cheatgrass has developed into a severe weed in several agricultural systems throughout North America, particularly western pastureland, rangeland, and winter wheat fields. It is now estimated to infest more than 101 million acres in western states.

**Cheatgrass**  
*Bromus tectorum*  
**Shíyínáldzidí**



Patrick Alexander



Patrick Alexander

Drooping seedhead of mature cheatgrass, showing sharp-tipped seeds and barbed awns



Matt Lavin

# Cordgrass, Alkali

*Spartina gracilis*

## Tł'ohdá'ákáłitsoh



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Good for erosion control

**Alkali cordgrass** is a robust grass with erect stems  $\frac{1}{2}$  to 3 feet tall. Reproduction is by seed and by rhizomes. It starts growth in late spring and begins flowering in July. Seedheads are clusters about 2 to 10 inches long. Leaf blades are less than  $\frac{1}{4}$  inch wide and less than or up to 12 inches long, flat or with edges rolled upward and in on drying, and gradually tapered to more or less finely pointed tips. Alkali cordgrass is found at elevations from 4,000 to 6,500 feet where annual precipitation is from 12 to 30 inches, but it is not particularly sensitive to precipitation zones since it grows along watercourses, in wetlands, or in wet meadows. Alkali cordgrass is an excellent grass for erosion control along rivers and streams. Its foliage is coarse but readily grazed by cattle and horses. Historically, alkali cordgrass was a major constituent of native meadow hay but does not withstand close harvest and has been replaced with tall wheatgrass in many meadow areas. It regrows slowly, but is fire tolerant.

*\*Description based on Utah State University's Range Plants of Utah description.*



**Cordgrass, Alkali**

*Spartina gracilis*

**Tł'ohdá'ákáilitsoh**



Robert H. Mohlenbrock, hosted by the USDA-NRCS, Western wetland flora: Field office guide to plant species (1992), West Region, Sacramento



Matt Lavin



Matt Lavin



Matt Lavin



Patrick Alexander

# Dropseed, Giant

*Sporobolus giganteus*

Tł'ohyilzólítsoh



## Alternative Names:

Giant sacaton

## Plant Type:

Grass

## Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

River valleys

## Growing Season:

Warm season

## Special Considerations:

None

**Giant dropseed** is a perennial with a hairy ligule. It grows as high as 6 ½ feet tall and can easily be taller than the botanist observing it. The panicles are closed and dense.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*


**Dropseed, Giant**

*Sporobolus giganteus*

**Tł'ohyilzólítsoh**



*Patrick Alexander*



*Russ Kleinman & Richard Felger*



Max Licher

# Dropseed, Pine

*Blepharoneuron tricholepis*

## Tł'ohyilzólí



### Alternative Names:

Hairy dropseed

### Plant Type:

Grass

### Habitat:

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

### Growing Season:

Warm season

### Special Considerations:

- ✓ Provides livestock forage
- ✓ Indicator of healthy rangeland

**Pine dropseed** is a slender, erect, densely tufted bunchgrass growing 10 to 13 inches tall, with basal leaves and fibrous roots. It reproduces from seeds and tillers. It starts growth in late June or early July, completes growth in September, and generally comprises only a small portion of the vegetation.

The panicle-type inflorescence is 2 to 9 inches long, and can be open or contracted. Spikelets are small, containing one floret with a distinctive greenish-gray color, and with silky hairs.

Pine dropseed is usually found in open parks and meadows in the subalpine zone and in the open timber of the ponderosa pine and piñon-juniper types.

The palatability and quality of young pine dropseed plants is very good for all classes of livestock. The quality rapidly declines with maturity. The stems are neglected or only slightly grazed after maturing. It is generally one of the highest quality grasses in timbered areas.

A healthy pine dropseed community in areas where it is native is an indicator of rangeland in good condition.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



**Dropseed, Pine**  
*Blepharoneuron tricholepis*  
**Tł'ohyilzólí**



Patrick Alexander



Patrick Alexander



Patrick Alexander





Max Licher

# Dropseed, Sand

*Sporobolus cryptandrus*

Tł'ohzhóó'



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

None

**Sand dropseed** is a perennial bunchgrass growing from 2 to 3 feet high. Sand dropseed is extremely drought tolerant and is adapted to sites receiving 7 to 16 inches of annual precipitation. Its fine root system allows sand dropseed to extract water from depths of up to 12 inches. During periods of summer drought, sand dropseed leaves roll up to reduce surface area and evapotranspiration.

This species spreads naturally from seed once established and increases on depleted rangelands and wastelands. Sand dropseed plants are able to withstand heavy use due to their protected root crown and late maturity and because they are less preferred than other species. Plants can be killed by overgrazing as a result of continued close cropping; however, when grazed properly, sand dropseed increases on sites with poor conditions.

**Dropseed, Sand**  
*Sporobolus cryptandrus*  
**Tł'ohzhóó'**



*Matt Lavin*



*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*



Patrick Alexander

# Dropseed, Spike

*Sporobolus contractus*

## Tł'ohyilzólídeeníní



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Provides wildlife and livestock forage

**Spike dropseed** is a native, perennial bunchgrass with erect stems that grow three feet tall. Spike dropseed begins growing in the spring when moisture levels increase, and it and matures to produce seed in October. It can reseed itself following overgrazing or drought; however, proper management is required to ensure the plant is not overgrazed or over utilized by livestock or wildlife.

Spike dropseed has fair nutritional value, and many species of livestock will graze it. Various wildlife species, particularly upland birds, also feed on it, and it provides good cover for birds and small mammals.



**Dropseed, Spike**  
*Sporobolus contractus*  
**Tł'ohyilzólídeeníní**



Patrick Alexander



Patrick Alexander

Sheaths of *Sporobolus contractus* leaves are hairy around the margins



Patrick Alexander



Patrick Alexander



Patrick Alexander

# Sacaton, Alkali

*Sporobolus airoides*

## Tłohdá'ákáłitsoh



### Alternative Names:

Salt grass

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Good forage
- ✓ Stabilizes soil

**Alkali sacaton** is a perennial bunchgrass. It ranges in height from 20 to 60 inches, with flat leaves  $\frac{1}{16}$  to  $\frac{1}{4}$  inches wide that taper from the base of the leaf. The inflorescence is an open panicle 8 to 20 inches long with a pyramidal shape. The small seeds rest singly on branches in the loose, open seedhead.

Alkali sacaton blooms from April to May in the Southwest. It grows in both saline and nonsaline coarse, medium, and fine textured soils. This grass is tolerant of salinity and a broad range in pH.

It is good forage for horses and cattle as well as deer, small mammals (it is relished by jackrabbits), and birds. It is frequently utilized for seeding and stabilizing disturbed areas. Due to its salt tolerance, it is recommended for seeding sites such as oil well pits and saline waste from power plants.

After establishment, alkali sacaton is tolerant of drought and water inundation. However, it is intolerant of shade and is usually found growing in open areas.



## Grasses

### Sacaton, Alkali *Sporobolus airoides* Tłohdá'ákáłitsoh



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander

# Fescue, Arizona

*Festuca arizonica*

## Tsinyaat'ohyilzólí

**Alternative Names:**

None

**Plant Type:**

Grass

**Habitat:**

Grasslands

**Growing Season:**

Cool season

**Special Considerations:**

None

**Arizona fescue** is a native, long-lived perennial bunchgrass. The plant has no rhizomes. Culms (stems) are 1 to 3 feet tall. They are clustered, pale blue-green, and rough to the touch. Leaf blades are mostly basal, pale blue-green, 6 to 10 inches long, shaped like a string, and rough to the touch. The inflorescence is a panicle about 3 to 8 inches long. It has a deep, dense fibrous root system.

**Fescue, Arizona**

*Festuca arizonica*

**Tsinyaat'ohyilzólí**



Patrick Alexander



Patrick Alexander



Patrick Alexander





Russ Kleinman

# Fescue, Six Weeks

*Vulpia octoflora*

Tł'ohyilzólídiijoolí



## Alternative Names:

Sixweeks fescue

## Plant Type:

Grass

## Habitat:

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

## Growing Season:

Cool Season

## Special Considerations:

None

**Six weeks fescue** has solitary stems or stems in small tufts generally 3 to 12 inches tall. Under unusually wet conditions it may grow to 24 inches tall. It is erect or occasionally bent upward at the lower nodes. It is shallow rooted and easily pulled from the soil. Its seedhead is a panicle 1 to 4 inches long with short branches compressed against the rachis (stem). Spikelets contain 5 to 15 florets. Glumes are unequal with the first being awl-shaped, sharp, and  $\frac{1}{4}$  inch long. The second glume is narrow,  $\frac{1}{8}$  to  $\frac{1}{4}$  inch long, and sharp or with an awn tip.

Leaf blades are narrow and boat-shaped. Sheaths are smooth or with backward hairs.

It is adapted to a wide range of soils and soil textures but is most common on sterile, rocky ground and disturbed areas.

Associated species include cheatgrass, red brome, creosotebush, bursage, big sagebrush, and storksbill.

Six weeks fescue is generally so small that it provides little forage for large animals. It may be used by desert tortoises and small desert animals since it is common in their habitats. It provides little erosion control.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



## Fescue, Six Weeks

*Vulpia octoflora*

Tł'ohyilzólíidijoolí



Russ Kleinman, Richard Felger,  
Kevin Keith, & Sarah Johnson



Russ Kleinman, Richard Felger, Kevin Keith, & Sarah Johnson



Russ Kleinman, Richard Felger, Kevin Keith, & Sarah Johnson



Russ Kleinman, Bill Norris, &  
Denise Friedrich

# Foxtail Barley

*Hordeum jubatum*

## Azéé'iilwo'itsoh



### Alternative Names:

Bobtail barley

Squirreltail barley

### Plant Type:

Grass

### Habitat:

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

River valleys

Wastelands

### Growing Season:

Cool season

### Special Considerations:

! May injure grazing animals

**Foxtail barley** is a native grass that looks quite a bit like longleaf squirreltail. In foxtail barley, however, spikelets are borne in sets of three. The middle one is sessile, and the two outer spikelets are on pedicels. Foxtail barley propagates by seed and is extremely tolerant of saline soils. It inhabits both dry and wet areas and can be drought tolerant.



## Grasses

### Foxtail Barley

*Hordeum jubatum*

Azéé'ilwo'itsoh



Russ Kleinman



Russ Kleinman



Russ Kleinman, Bill Norris, & Denise Friedrich



Russ Kleinman



Patrick Alexander

# Galleta, James'

*Hilaria jamesii*

## Tł'ohdich'ízhí



### Alternative Names:

Galleta

Formerly *Pleuraphis jamesii*

### Plant Type:

Grass

### Habitat:

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

Wastelands

### Growing Season:

Warm season

### Special Considerations:

None

**Galleta** is a perennial grass with strong rhizomes, but commonly is found growing in bunches. Stems are erect, 11 to 19 inches tall. It is a desirable forage plant for cattle, horses, and sheep during active plant growth in late spring and early summer. Palatability is moderate during active growth. Galleta is relatively unpalatable during dormant periods. Galleta greens up in early summer if rains are adequate. Growth is dependent on precipitation and is tolerant of close grazing and trampling.

Sheep typically feed upon central portions of galleta tufts, leaving the coarser growth around the outer edges. It is used to some extent by deer and antelope. Galleta is commonly found between 3,500 and 7,500 feet elevation in zones of 5 to 16 inches of annual precipitation.

Galleta has been used for basketry, coiled trays, and as floor and hair brushes. It has also been used as a dietary/pediatric aid to encourage babies' appetites. It is also used by horses, deer, and antelope because it withstands trampling and close grazing.



**Galleta, James'**

*Hilaria jamesii*

**Tł'ohdich'ízhí**



Matt Lavin



Matt Lavin



Matt Lavin



Patrick Alexander



Patrick Alexander

# Goatgrass, Jointed

*Aegilops cylindrica*

## Cl'oh ałhe ni'lí



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

Piñon-juniper woodland

### Growing Season:

Cool season

### Special Considerations:

! Noxious

**Jointed goatgrass** is an introduced, annual grass that has spread into wildlands from winter wheat croplands. It is listed as a noxious weed in Arizona and New Mexico. Plants resemble winter wheat, but seedlings have fine hairs along leaf margins, sheaths, and auricles. The first leaf to emerge on seedling jointed goatgrass is reddish to brownish-green, whereas wheat tends to emerge as a whitish-green leaf.

Stems are hollow and may be upright, slightly lying down, or abruptly bent near the base. Stems branch at the base; the stems that branch off the main stem are called tillers. Jointed goatgrass is a primary concern within or near winter wheat fallow production areas where the species can contribute to reduced yield, loss of profitability, and degraded land values. Preventing the introduction of jointed goatgrass should be the key objective when managing this weedy species. The best way to prevent introduction is to always use certified weed-free seed from a reliable seed source when planting winter wheat in any wildland situations. Once jointed goatgrass is established, it spreads solely via seed. Control strategies should focus on removing seed production and reducing the opportunity for germination of seed already present in the soil. In most cases, several years of treatment are necessary to control jointed goatgrass.

**Goatgrass, Jointed**

*Aegilops cylindrica*

**Cl'oh ałhe ni'lı**



USDA APHIS PPQ - Oxford, North Carolina, USDA APH



Matt Lavin



Joseph M. DiTomaso, UC Davis



Joseph M. DiTomaso, UC Davis



Patrick Alexander





Matt Lavin

# Gramma, Black

*Bouteloua eriopoda*

Tł'ohnást'ąsłizhiní



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Wastelands

Grasslands

## Growing Season:

Warm season

## Special Considerations:

- ✓ Good livestock forage
- Easily killed by overgrazing

**Black grama** is a native, perennial grass 10 to 20 inches tall. It has crooked, woolly stems with narrow leaves, less than  $\frac{1}{8}$  inch wide, inrolled and wavy. Seedheads are similar to blue grama, except narrower. It is one of the region's most valuable and nutritious grasses.

It grows mostly on gravelly or sandy soils that are dry and seldom grows on clay loams or adobe (clay) flats. Black grama is characteristically a lower altitude grass (3,500 to 5,500 feet elevation), but occasionally grows at elevations of 7,000 feet.

Overgrazing easily kills this grass. To improve black grama ranges, defer grazing two successive years during the growing season, and do not graze more than 50% of current growth by weight during dormancy. Once the improvement is achieved, *maintain* ranges by deferring grazing for one growing season every third year.



## Grasses

### Grama, Black *Bouteloua eriopoda* Tł'ohnást'ąsłizhinií



Patrick Alexander



Patrick Alexander



Patrick Alexander

Black grama range in winter



Patrick Alexander

Black grama range in summer



Patrick Alexander

# Gramma, Blue

*Bouteloua gracilis*

Tł'ohnást'ąsi



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

- ✓ Good livestock forage
- ✓ Cures well on stem

**Blue grama** is a major warm-season grass, 10 to 20 inches tall with narrow basal leaves of 3 to 6 inches. Blue grama grows in definite bunches and reproduces by tillering and by seed. Mature seedheads are curved, resembling a human eyebrow.

Once the grass is established, it is very palatable to livestock all year long. Since growing points are at or near the ground, the grass withstands fairly close grazing. Blue grama cures well on stem, making it a good grass for grazing during the dormant season.

For best yields, defer grazing during the growing season every two to three years.



**Grama, Blue**  
*Bouteloua gracilis*  
**Tł'ohnást'ąsi**



Patrick Alexander


Blue grama rangeland in the dry season



Renee Benally



Amy Smith Muiise



Renee Benally



Patrick Alexander

# Grama, Sideoats

*Bouteloua curtipendula*

Tł'ohłichi'í



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

- ✓ Good livestock forage

**Sideoats grama** is a medium-size perennial bunchgrass, 15 to 30 inches tall or occasionally taller. This is the largest and most coarse of the grama grasses. It has a blue-green color, sometimes with a purplish cast (especially in the spring), and cures to a reddish-brown or straw color. Leaves are coarser than other species of gramas, straight, comparatively stiff, and mostly basal. Ten to thirty small, non-comb-like spikes are borne mostly along one side of each central seed stalk. These spikes drop when mature, leaving a long zigzag stalk. It produces high quality, nutritious forage that is relished by all classes of livestock throughout the summer and fall, and it remains moderately palatable into winter. This makes it one of the most important range grass species.

Sideoats is moderately drought tolerant, but less so than blue grama. It is moderately tolerant of semi-shaded conditions and can be found in open woodlands. It is fairly tolerant of fire (when in a dormant state) and of spring flooding. It probably has the widest range of adaptation of any of the warm-season perennial grasses.




## Grama, Sideoats

*Bouteloua curtipendula*


Tł'ohłichi'í



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



University of Wisconsin-Madison,  
Wisconsin State Herbarium

# Johnsongrass

*Sorghum x alnum*

## Akál



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Warm season

### Special Considerations:

- ! Noxious
- ! May suppress other plants
- ! Sometimes toxic to livestock, especially cattle

**Johnsongrass** is a tall perennial with long rhizomes. It grows in moist places and can form large colonies. The flowering segments are called rames. In the rames, a pedicelled spikelet can give rise to more pedicelled spikelets.

Johnsongrass was introduced to the United States as a forage crop, and generally provides good feed. At certain stages of development, however, it can be toxic to livestock. Toxins may also be present when the grass grows in adverse conditions such as drought, frost, or extreme heat. Toxins may also be present when the grass is wet. Toxins tend to affect cattle more than horses. Stalks grow to a height of eight feet. The leaves have a distinctive white midrib. The flowers occur in a purplish cluster. The plants spread through rhizomes and contain allelopathic chemicals, toxins that may affect the health and growth of other plants.

# Johnsongrass

*Sorghum x alnum*

Akál



Russ Kleinman

*Sorghum halapense*



Russ Kleinman

*Sorghum halapense*, detail of glume



D. Walters and C. Southwick, Table Grape Weed Disseminule ID, USDA APHIS PPQ, Bugwood.org

*Sorghum halapense*, strongly rhizomatous



Bruce Ackley, The Ohio State University, Bugwood.org



Patrick Alexander

# Junegrass, Prairie

*Koeleria macrantha*

## Tł'ooléts'ózí



### Alternative Names:

Junegrass

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Cool season

### Special Considerations:

- ✓ Good wildlife and livestock forage

**Prairie junegrass** is a highly variable, moderately long-lived, perennial bunchgrass that grows ½ to 2 feet tall. Its clusters of narrow, markedly veined, light green to bluish-green leaves grow to about seven inches tall.

Active growth begins early in spring, providing good forage for livestock, deer, antelope, and elk. Declining palatability during seed formation rebounds in late summer when there is adequate moisture.



# Junegrass, Prairie

*Koeleria macrantha*

Tł'ooléts'ózí



Matt Lavin




Matt Lavin




Patrick Alexander



Patrick Alexander



Matt Lavin



Patrick Alexander

# Muhly, Mountain

*Muhlenbergia montana*

## Bé'ézhóó'



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

Ponderosa pine/mixed conifer

### Growing Season:

Warm season

### Special Considerations:

- ✓ Good livestock forage

**Mountain muhly** is a perennial bunchgrass with high forage value for cattle. It is a valuable forage plant because of its abundance, though it does not have high palatability. It is grazed most readily when the plants are actively growing. Mountain muhly decreases in response to trampling and heavy grazing.

It is found in grassy parklands, on slopes and foothills, on rocky, dry hillsides, in canyons, and on mesas. Mountain muhly grows during the spring and summer months and becomes semi dormant if there is midsummer drought.

## Muhly, Mountain

*Muhlenbergia montana*

B'é'zhóó'



Patrick Alexander



Patrick Alexander



Patrick Alexander





Frankie Coburn

# Muhly, Ring

*Muhlenbergia torreyi*

## Bé'ézhóó'łichií'í



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

- ▣ Needs rest in the rainy season

**Ring muhly** is a low-growing, fine-leaved, fine-stemmed sodgrass. It tends to grow in rings, which are caused by the center dying out as the plant enlarges. The rings may range in size from several inches to a few feet across.

It appears green to bluish-green, reddish, or purplish. It flowers from July to September on dry ridges, sandy plateaus, and rocky slopes between 4,000 and 7,000 feet. Even when ring muhly is young and growing rapidly, its palatability is low. As the plants mature, palatability drops almost to zero.

Ranges with ring muhly should be managed with other grasses as key species. Continued attempts to obtain even a moderate amount of grazing from the grass will in time drive out all of the desirable forage species and result in consistent weight losses in the animals being grazed. Ranges with an abundance of this grass should be rested during the summer rainy season.



**Muhly, Ring**  
*Muhlenbergia torreyi*  
**Bé'ézhóó'łichíí'í**



Patrick Alexander



Patrick Alexander



Patrick Alexander



Matt Lavin

# Muhly, Sandhill

*Muhlenbergia pungens*

## Béé'ézhóó'



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Controls wind erosion on sand

**Sandhill muhly**, often found in or near blowout or dune areas on sandy sites, is a perennial bunchgrass 4 to 24 inches tall. It often forms large, rounded clumps or rings, dying out in the center. It reproduces by rhizomes and seed. Sandhill muhly has little value as forage for livestock or wild grazers. It is effective in controlling wind erosion in very sandy areas.

**Muhly, Sandhill**  
*Muhlenbergia pungens*  
**Béé'ézhóó'**



Matt Lavin



Arizona State University Herbarium





Max Licher

# Muhly, Spike

*Muhlenbergia wrightii*

## Bé'ézhóó'tsahí



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

Piñon-juniper woodland

### Growing Season:

Warm season

### Special Considerations:

✓ Good for livestock

**Spike muhly** is a perennial bunchgrass reaches heights of 8 to 24 inches. Clumps are generally 1 to 2 feet wide. The grass has a very narrow seedhead that has a distinctive feature: the seeds do not occur continuously, but are sometimes interrupted by bare areas. There is one floret per spikelet.

Spike muhly rates high in palatability for all livestock. It is most abundant on rocky soils and in mountain meadows at elevations from 3,800 to 9,000 feet. Abundant spike muhly on rangeland in good condition needs only moderate grazing for maintenance. Depleted ranges may need summer deferment.

**Muhly, Spike**  
*Muhlenbergia wrightii*  
**Bé'ézhóó'tsahí**



*Russ Kleinman*



*Russ Kleinman*



*Russ Kleinman, Bill Norris, Karen Blisard, & Denise Friedrich*



*Russ Kleinman, Bill Norris, Karen Blisard, & Denise Friedrich*



Patrick Alexander

# Mutton Grass

*Poa fendleriana*

## Tł'ohnikání



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Cool season

### Special Considerations:

- ✓ Good livestock and wildlife forage

**Mutton grass** is a perennial bunchgrass that is rated as excellent forage for cattle and horses. It is also good forage for sheep, elk, and deer. During the winter, seedheads are eaten by pronghorn antelope. Seeds and leaves are also utilized by birds.

The fibrous root system of this species reaches a depth of approximately 10 inches, providing good surface erosion control in arid sites.

It can be used to restore areas where juniper encroachment has depleted the herbaceous understory following juniper removal. The species is drought tolerant and has potential for use in restoration and native plant diversification projects.



Patrick Alexander



Patrick Alexander





Patrick Alexander

# Needle-and-Threadgrass

*Hesperostipa comata subsp. comata*

## Ch'il Bilata dee'ni'ni'



### Alternative Names:

*Stipa cometa*

*Stipa comata*

*Hesperostipa comata*

Spear-grass

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Cool season

### Special Considerations:

✓ Cures well on the stalk

! Sharp awn may injure livestock

**Needle-and-threadgrass** is a native perennial bunch grass. It ranges in height from 1 to 4 feet. It has narrow basal leaves 3 to 12 inches long with the blade usually rolled inward.

The mature spikelet has a long, hairy, spiraling awn.

Needle-and-threadgrass appears in many habitat types, including forest, grassland, and shrub-steppe communities. It reproduces by seed and tillers. Because of a long awn needle-and-thread grass seeds can embed themselves in the soil by a twisting action of the awn in response to daily humidity changes. This sharp awn may injure grazing animals, especially domestic sheep.

Needle-and-threadgrass is grazed by all livestock, particularly in the spring and fall when green, but it also cures well to provide dormant season grazing.

*Description is partially drawn from the USDA USFS Fire Effects Information System.*

**Needle-and-Threadgrass**  
*Hesperostipa comata* subsp. *comata*  
**Ch'il Bilata dee'ni'ni'**



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Matt Lavin

# Needlegrass, Desert

*Pappostipa speciosa*

Tł'ohsahí



## Alternative Names:

Formerly *Stipa speciosa*

*Achnatherum speciosa*

## Plant Type:

Grass

## Habitat:

Brushlands

Grasslands

Wastelands

## Growing Season:

Cool season

## Special Considerations:

! May injure livestock

**Desert needlegrass** is a native, perennial bunchgrass that grows from 1 to 2 feet tall, typically on coarse soils, including alluvial fans, dry, rocky hills, talus slopes, and in canyons. It can tolerate low precipitation and usually occurs in areas that receive 6 to 20 inches annually. The long awns of desert needlegrass seed enable self-planting. Cycles of wetting and drying cause the awn to twist and untwist, which drives the seed into the ground.

Desert needlegrass tolerates light grazing. It is best to graze desert needlegrass before seed develops because the seed has a prominent, sharp callus that can injure the eyes and mouths of grazing animals.



**Needlegrass, desert**

*Pappostipa speciosa*

**Tł'ohtsahí**



Matt Lavin



Matt Lavin



Matt Lavin



Matt Lavin



C.D. Schelz

# Needlegrass, Scribner

*Achnatherum scribneri*

**Tł'ohsahí**



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Piñon-juniper woodland

Ponderosa pine/mixed conifer

## Growing Season:

None

## Special Considerations:

None

**Scribner needlegrass** grows in dense mats or tufts, often on rocky slopes among conifers. The stalks can grow to three feet tall. It gets its name from the needle-like awns that protrude from its seeds. These awns can be ½ to 1 inch long and are slightly fuzzy.

Scribner needlegrass occurs on parts of the Navajo Nation but is not widespread.


## Needlegrass, Scribner

*Achnatherum scribneri*

Tł'ohtsahí



G. J. Goodman



USDA-NRCS PLANTS Database / Hitchcock, A.S. (rev. A. Chase), Manual of the grasses of the United States (1950), USDA Miscellaneous Publication No. 200, Washington, DC



Tracey Slotta, hosted by the USDA-NRCS PLANTS Database





Max Licher

# New Mexico Feathergrass

*Hesperostipa neomexicana*

Tł'ohdit'ódítsoh



## Alternative Names:

Formerly *Stipa neomexicana*

## Plant Type:

Grass

## Habitat:

Brushlands

Grasslands

## Growing Season:

Cool season

## Special Considerations:

- ! Seeds may injure livestock
- ✓ Withstands heavy use in spring and fall

**New Mexico feathergrass** is a perennial bunchgrass. This tufted grass has a long, feathery awn on the seed, which distinguishes it from needle-and-threadgrass. Most common on sandy, gravelly, and rocky sites, it occurs at elevations from 3,100 to 7,200 feet.

New Mexico feathergrass provides moderate to good forage for all livestock. The sharp-pointed seeds, like needle-and-threadgrass seeds, may injure the mouths of animals. The grass tends to decrease under grazing, especially on drier ranges. This grass withstands heavy use in the spring and fall and usually reproduces well if the plants are allowed to mature seed during summer.

## New Mexico Feathergrass

*Hesperostipa neomexicana*

Tł'ohdit'ódítsoh



Amy Smith Muise



Russ Kleinman & Bill Norris



Amy Smith Muise



Russ Kleinman & Bill Norris



Russ Kleinman &amp; Scott Zager

# Redtop

*Agrostis gigantea*

**Tł'ohlátahyíchii'í**



## Alternative Names:

Blackbent

## Plant Type:

Grass

## Habitat:

Grasslands

## Growing Season:

Cool season

## Special Considerations:

- ! Introduced
- ✓ Palatable to cattle and horses

**Redtop** is a rhizomatous perennial grass that makes a coarse but fairly dense turf. Leaves are narrow and sharp and about  $\frac{3}{8}$  inch wide. The stems are slender, growing to 30 or 40 inches tall. The inflorescence is pyramidal and reddish in color—hence the name.

The leaf is harsh to the touch with a prominent ligule. The seedhead is a purplish-red, loose panicle, pyramid-shaped, 4 to 12 inches long. It occurs mostly in mountain meadows and subalpine grasslands where openings in the tree canopy occur, as well as in wet meadows in the western plateau and central plains. Redtop grows at elevations of 5,000 to 10,000 feet, but is most common at 7,500 feet.

Redtop was introduced from Europe as a cultivated species. It is palatable to cattle and horses, and sheep will graze it if necessary. When moisture is adequate, the plants remain green all summer and can be grazed throughout the season. Redtop grows vigorously and forms good sod. Because of this, on its preferred sites, the plant withstands heavy use and serves well as a soil binder for reclaiming gullies and holding slopes and banks. It is well adapted to wet, acid soils.



**Redtop**

*Agrostis gigantea*

**Tł'ohlátahyíchii'**



*Russ Kleinman*



*Russ Kleinman*



*Russ Kleinman*



Debra Moses

# Ricegrass, Indian

*Achnatherum hymenoides*

## Ndídíídi



### Alternative Names:

*Stipa hymenoides*

Formerly *Oryzopsis hymenoides*

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Cool season

### Special Considerations:

- ✓ Good livestock and wildlife forage
- ✓ Edible grain

**Indian ricegrass** is a cool season, native bunchgrass with many tightly rolled, slender leaves growing from the base of the bunch. The nutritious seeds of Indian ricegrass have a long history as a staple food.

Indian ricegrass is highly palatable to livestock and wildlife. It is a preferred feed for cattle, horses, and elk in all seasons. It is considered a preferred feed for sheep, deer, and antelope in spring, and a desirable feed for sheep, deer, and antelope in late fall and winter. It reaches its peak production from mid-June through mid-July. It holds its nutrient value well at maturity.

Indian ricegrass is very winter hardy and has a broad climatic adaptation. It can be found at elevations from 2,000 to 10,000 feet. It grows best in areas with average annual precipitation of 8 inches to 14 inches.

**Ricegrass, Indian**  
*Achnatherum hymenoides*  
**Ndídlííí**



Amy Smith Muise



Amy Smith Muise





Patrick Alexander

# Saltgrass

*Distichlis spicata*

Tł'ohdík'ózhíqtł'oolédich'ízhí



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

River valleys

## Growing Season:

Warm season

## Special Considerations:

✓ Drought tolerant

! Can become invasive

**Saltgrass** is a native perennial from 15 to 35 inches in height. It forms dense mats with rhizomes and sometimes stolons. It is a dioecious species with male and female reproductive parts on separate plants. It is one of the most drought-tolerant species in the dry West.

Blades are firm, the edges often flat at the base and folded or rolled inward meeting in the middle, therefore, appearing attenuate. Blades are generally less than four inches long. Salt crystals may be found on the leaves and stems. The ligules are stiff with a fringe of hair at the tips.

Saltgrass is grazed by both cattle and horses, and it has a forage value of fair to good because it remains green when most other grasses are dry during the drought period. It is resistant to grazing and trampling.

It has been used as a salty seasoning.

## Saltgrass

*Distichlis spicata*

Tł'ohdík'ózhíqtł'oolédich'ízhí



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Macleay Grass Man

# Sandbur, Field

*Cenchrus spinifex*

Dííłihí



## Alternative Names:

Schrader's brome

Formerly *Cenchrus incertus*

## Plant Type:

Grass

## Habitat:

Brushlands

Grasslands

## Growing Season:

Warm season

## Special Considerations:

! Noxious

! May cause injury

**Sandbur** is an annual grass with a spreading rather than upright growth habit. It often grows in areas that are seasonally flooded. The barbed spines on the burs can inflict a painful wound and are easily picked up and transported by clothing, hair, animal fur, and tires. The spikelets are hidden within the armed burs.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*



**Sandbur, Field**

*Cenchrus spinifex*

**Dííłiihí**



*D. Walters and C. Southwick*



*Robert Vidéki, Doronicum Kft.,  
Bugwood.org*



*Joseph M. DiTomaso,  
University of California - Davis*



*Joseph M. DiTomaso,  
University of California - Davis*



Patrick Alexander

# Three-Awns (Annual)

*Aristida spp.*

**Azéé'iilwo'iiyázhí**



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

- ! Fleece damage to sheep
- ! Irritates livestock skin, eyes, nostrils

**Three-awns** are grasses that occur in both annual and perennial forms. On Navajo rangelands, the annual three-awn species are six weeks three-awn (*Aristida adscensionis*) and oldfield three-awn (*Aristida oligantha*). Three-awns are native to North America. They are not considered a valuable forage species, though they do provide some habitat for birds and other small animals. The awns can tend to catch in the wool or irritate the skin of livestock. These bunchgrasses are many-branched toward the base of the clump, and they grow to a height of about three feet. The leaves roll inward. The florets and awns of the inflorescences spread outward, creating a brushy appearance. There are three awns on each floret.

# Three-Awns (Annual)

*Aristida* spp.

Azéé'iilwo'iiyázhí



Patrick Alexander

*Aristida adscensionis*



Patrick Alexander

*Aristida oligantha*



Patrick Alexander

*Aristida oligantha*





Amy Smith Muise

# Three-Awns (Perennial)

*Aristida spp.*

**Azéé'iilwo'iiyázhí**



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Grasslands

Brushlands

## Growing Season:

Warm season

## Special Considerations:

! Fleece damage to sheep

**Three-awns** are bunchgrasses that occur in both annual and perennial forms and have three awns on each floret. On Navajo rangelands, the most common perennial three-awn is purple three-awn (*Aristida purpurea*). Three-awns grow in wastelands and brushlands and are not considered valuable forage species, though they do provide some habitat for birds and other small animals. The awns can tend to catch in the wool or irritate the skin, mouths, or nostrils of livestock.

Many-branched toward the base of the clump, purple three-awn grows to a height of 8 to 16 inches. The leaves are narrow and roll inward. The florets and awns of the inflorescences spread outward, creating a brushy appearance. The florets turn purplish-red.

Three-awn does well on rocky or sandy soils. Presence of the grass may indicate an overgrazed range, but it may also indicate a dry, well-drained soil. It is often found with sagebrush, cheatgrass, needle-and-threadgrass, western wheatgrass, and blue grama. Purple three-awn increases under heavy grazing because it is less palatable than associated grasses. It has excellent drought resistance.

The palatability of this species when green is low to moderate for all livestock. The grass has little value when dry. Although growth begins in the late spring, little forage is produced until the summer rains.

Areas of deep soil supporting extensive stands of red three-awn can be improved through light, winter, or deferred grazing.

## Three-Awns (Perennial)

*Aristida* spp.

Azéé'ilwo'iiyázhí




Patrick Alexander

*Aristida divaricata*, poverty three-awn



Patrick Alexander



Patrick Alexander

*Aristida divaricata*, poverty three-awn



Amy Smith Muike

*Aristida purpurea*



Patrick Alexander

*Aristida pansa*



Patrick Alexander

# Tufted Horse Grass

*Deschampsia cespitosa*

Tł'oh Deits'óózí



## Alternative Names:

Tufted hairgrass

## Plant Type:

Grass

## Habitat:

Grasslands

River valleys

## Growing Season:

Warm season

## Special Considerations:

None

**Tufted horse grass**, often called tufted hairgrass, is native to much of the United States. It is an important livestock forage species, sometimes used for hay. It grows in a wide variety of habitats and elevations. Stalks may grow from 2 to 6 feet in height, forming a feathery seedhead. Basal leaves grow in a dense tuft.



**Tufted Horse Grass**

*Deschampsia cespitosa*

**Tł'oh Deits'óózí**



Patrick Alexander



Patrick Alexander



Patrick Alexander



Franz Xaver [CC BY-SA 3.0 or GFDL],  
from Wikimedia Commons

# Wheatgrass, Crested

*Agropyron cristatum*

Tł'ohdeindtł'izí



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

Brushlands

Grasslands

Wastelands

## Growing Season:

Cool season

## Special Considerations:

- ! Introduced
- ✓ Wildlife and livestock forage

**Crested wheatgrass** is an introduced, long-lived, drought tolerant, and winter hardy grass with an extensive root system.

It produces leaves in the spring about 10 days after bluegrass species and about two weeks earlier than native wheatgrasses. It makes good spring growth, little summer growth, and good fall growth if moisture is available. Livestock and wildlife will graze crested wheatgrass throughout the spring growing season until it becomes too coarse, and again in fall if regrowth occurs. Established stands can withstand very heavy grazing.

In spring, the protein levels can be as high as 18% and decrease to about 4% as the grass matures. Digestible carbohydrates remain high throughout the active growth period. It is commonly utilized for winter forage by cattle and horses, but protein supplements are required to ensure good animal health.

It is not considered a desirable feed for cattle, sheep, horses, deer, antelope, or elk in the summer.

This species was introduced from Asia.



## Wheatgrass, Crested

*Agropyron cristatum*

Tł'ohdeindtł'izi



Public Domain via Wikimedia Commons



Stefan.lefnaer [CC BY-SA 4.0], from Wikimedia Commons



Stefan.lefnaer [CC BY-SA 4.0], from Wikimedia Commons



Stefan.lefnaer [CC BY-SA 4.0], from Wikimedia Commons



Stefan.lefnaer [CC BY-SA 4.0], from Wikimedia Commons





Russ Kleinman

# Wheatgrass, Slender

*Elymus trachycaulus*

**Tł'ooléyilts'oozí**



## Alternative Names:

None

## Plant Type:

Grass

## Habitat:

River valleys

## Growing Season:

Cool season

## Special Considerations:

- ✓ Wildlife and livestock forage
- ✓ Erosion control

**Slender wheatgrass** is a bunchgrass growing to a height of 1 to 2 ½ feet, usually without rhizomes, but tillering freely. It starts growth in mid spring, and seeds mature by August to September. It is native to much of the West, rarely abundant, but common in a number of plant communities. It is found at elevations from 6,000 to 13,500 feet on gentle to steep mountain slopes, valley bottoms, and rolling hills. It occurs on practically all of the mountain and high mountain sites, but only on a few of the upland sites. Under good management, it is moderately drought resistant as well as resistant to long wet periods. It prefers deep, well-drained, medium- to fine-textured soils that are well developed. It is not considered tolerant of salt. Associated species include mountain brome, yarrow, Junegrass, aspen, cottonwood, and big sagebrush.

Its forage value is good, diminishing in palatability in the latter part of the growing season. In general, an excellent forage grass for cattle, horses, sheep, elk, and deer and, valuable as cover and feed for many forms of small mammals and upland game birds as well as songbirds. Later in the growing season, it becomes slightly coarse and stemmy. Grazing management should take into account the fact that it is sensitive to grazing, especially by cattle. Serves as an indicator of trend in associated valuable species, since it withstands moderate grazing well but will decrease under heavy grazing. Erosion control values are excellent, because it will quickly establish from seed.

*\*Description courtesy of Utah State University's Range Plants of Utah.*

**Wheatgrass, Slender**

*Elymus trachycaulus*

**Tł'ooléyilts'oozí**



*Russ Kleinman & Bill Norris*



*Russ Kleinman & Kelly Kindscher*



Patrick Alexander

# Wheatgrass, Western

*Pascopyrum smithii*

## Tł'oolé



### Alternative Names:

Formerly *Agropyron smithii*

### Plant Type:

Grass

### Habitat:

Grasslands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Early spring forage is palatable to all livestock

**Western wheatgrass** is a long-lived species that has coarse blue-green leaves with prominent veins. It is a sod former with very strong, spreading rhizomes. Stems arise singly or in clusters of a few, and reach heights of 1 to 3 feet. Western wheatgrass provides early spring forage and is palatable to all livestock



**Wheatgrass, Western**

*Pascopyrum smithii*

**Tł'oolé**



*Patrick Alexander*



*Russ Kleinman, Bill Norris, Leith Young, & Richard Felger*



*Russ Kleinman, Bill Norris, Leith Young, & Richard Felger*



*Patrick Alexander*



Russ Kleinman & Bill Norris

# Wildrye, Canadian

*Elymus canadensis*

## Tł'ohndtł'izítsoh



### Alternative Names:

Nodding wild rye

### Plant Type:

Grass

### Habitat:

Grasslands

Ponderosa pine/mixed conifer

### Growing Season:

Cool season

### Special Considerations:

None

**Canadian wildrye** is a tall perennial grass with a drooping inflorescence. It grows in moist forested areas. Glumes are well-developed and the lemmas have awns. The rachis does not break apart when mature.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*

**Wildrye, Canadian**

*Elymus canadensis*

**Tł'ohndtł'izítsoh**



Russ Kleinman & Bill Norris



Russ Kleinman





Patrick Alexander

# Wildrye, Russian

*Psathyrostachys juncea*

## Tł'oolétsohndtł'izí



### Alternative Names:

None

### Plant Type:

Grass

### Habitat:

Brushlands

Piñon-juniper woodland

### Growing Season:

Cool season

### Special Considerations:

None

**Russian wildrye** is an introduced, long-lived, perennial bunchgrass. It has an abundance of long, dense, basal leaves that are from 6 to 18 inches long and up to 1¼ inch in width. Plants vary from light to dark green, with many shades of blue-green. The erect, leafless reproductive stems are about 30 to 40 inches tall. The seedhead is a short, dense, erect spike with two or more short-awned spikelets clustered at axis joints. The seed shatters readily at maturity. The seed is about the same size as crested wheatgrass seed. The roots are fibrous and may establish to a depth of 6 to 8 feet. However, about 75% of the roots are in the surface 6 to 24 inches. Russian wildrye roots have an extended horizontal spread and may draw heavily on soil moisture for a distance of up to 4 or 5 feet. Its long season of growth and its vigorous soil-feeding habit make this species an excellent competitor with weeds once the grass is well established.

**Wildrye, Russian**

*Psathyrostachys juncea*


**Tł'oolétsohndtł'izi**



*Patrick Alexander*



*Patrick Alexander*



Wikimedia Commons

# Arrowgrass

*Triglochin maritima*

Tł'ohk'aa'í



## Alternative Names:

Seaside arrowgrass

## Plant Type:

Grasslike

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

! Toxic to livestock

**Arrowgrass** is a wandlike, herbaceous wetland plant 12 to 40 inches tall with rhizomes. It is a grasslike plant in the Juncaginaceae family. Reproduction is from rhizomes and seed. Growth starts in early spring, and flowering occurs June through August. Flowers are green and spaced along a spikelike raceme. Fruits are almost cylindrical, up to ¼ inch long and fall from the plant at maturity. Leaves are linear, 3 to 8 inches long, narrow, and flattened or channeled. Arrowgrass can be found in wet alkaline lowlands, and in meadows and swamps at elevations from 4,200 to 8,800 feet. It is occasionally confused with rushes because it occurs in habitats often dominated by them. It contains hydrocyanic acid, especially when drought- or frost-stressed, which makes it highly toxic to livestock, even in amounts less than ¼ pound. It is a common component of wet meadows that are cut for hay. Arrowgrass in cured hay has been reported to cause poisoning in young animals.

*\*Description based on Utah State University Range Plants of Utah.*



Patrick Alexander

Habitat





Patrick Alexander

# Bulrush

*Schoenoplectus acutus*

## Lók'aat'ohotsoh



### Alternative Names:

Tule

Common tule

Hardstem tule

Tule rush

Hardstem bulrush

Viscid bulrush

### Plant Type:

Grasslike

### Habitat:

Grasslands

River valleys

### Growing Season:

Warm season

### Special Considerations:

- ✓ Provides wildlife habitat and forage
- ! Potentially invasive

**Bulrush** is a grasslike species in the Cyperaceae (sedge) family. Bulrush is a wetland plant which forms dense colonies. It reproduces by seed, dispersed by wind and water. Bulrush seed establishes and germinates best on moist, bare soil. Hardstem bulrush also reproduces vegetatively from rhizomes. The flowering parts are compact and umbellate with a greenish bract extended. It often forms monocultures in marshes throughout its range. It grows best on sites with saturated soil or standing water for most of the year, such as marshes, swamps, seeps, washes, floodplains, lake and stream margins, and wet meadows. Hardstem bulrush can grow in fresh or brackish water where the water table is up to 5 feet above or ½ feet below the soil surface. It is fairly drought tolerant; it can persist through several years of dry conditions.

Hardstem bulrush is an important aquatic or semiaquatic species. It provides cover for numerous birds and mammals. It is a staple food for muskrat and other small mammals. It provides valuable food and nesting for a variety of birds.

It regrows well after removal and is tolerant of fire. It is considered a problem species in some circles because it can be an aggressively invasive plant that closes in bare shorelines and open water in marshes, impeding waterfowl and shorebirds. It is seldom grazed by livestock if other forage is available. If upland forage becomes limited and soil conditions dry, livestock and big game animals may utilize hardstem bulrush.

*\*Based on Utah State University Range Plants of Utah description.*

**Bulrush**

*Schoenoplectus acutus*

**Lók'aatl'ohotsoh**



Patrick Alexander



Patrick Alexander



Matt Lavin

# Rushes

*Juncus spp.*

## Lok'aá bit'zooz



### Alternative Names:

None

### Plant Type:

Grasslike

### Habitat:

Grasslands

River valleys

### Growing Season:

Cool season

### Special Considerations:

- ✓ Wildlife habitat and forage

**Wire rushes** are the primary rush species in Navajo rangelands. They are native to most of North America, and they will grow in wetlands as well as in seasonal wetlands, standing water, and moist grasslands. While not a primary livestock food source, rushes provide food and habitat for many types of wildlife. They also have cultural use as a basket-weaving material and occasional food. They are an important erosion control species due to their dense root systems.

Rushes are not a true grass, but grow in a grasslike form and can be identified by their round leaf cross sections. The wire rush grows in tufts to a height of three feet. The inflorescence may be up to 2¾ inches long and contain 10 to 50 flowers. Seeds are brownish red and the size of grains of sand.

Some people use this simple mnemonic to help distinguish rushes, sedges, and grasses: “Sedges have edges, rushes are round, grasses have knees that bend to the ground.”

An alternate version goes, “Sedges have edges, rushes are round, grasses are hollow right up from the ground.”

Rushes found in the Four Corners region include:

*Juncus arcticus*  
*Juncus bufonius*  
*Juncus ensifolius*  
*Juncus interior*  
*Juncus longistylis*  
*Juncus torreyi*



**Rushes**

*Juncus spp.*

**Lok'aá bit'zooz**



Matt Lavin



Matt Lavin



Matt Lavin



Matt Lavin



Howard F. Schwartz, Colorado State University, Bugwood.org

# Nutsedge, Yellow

*Cyperus esculentus*

## Tłohi'gaí



### Alternative Names:

Chufa

### Plant Type:

Grasslike

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

! Invasive

**Yellow nutsedge** is a native plant found across most of North America. Stems are triangular, growing to two feet in height. Leaves are grasslike and glossy. Yellow-brown flowers occur in a terminal cluster. The tubers (nuts) that give the nutsedge its name can remain in the soil even when the plants are pulled up or eaten by livestock. It can grow thickly, particularly in disturbed wetland areas, and is considered invasive.

Nutsedge is not a true grass, but grows in a grasslike form.



**Nutsedge, Yellow**

*Cyperus esculentus*

**Tłohi'gaí**



Patrick Alexander



Patrick Alexander



Howard F. Schwartz, Colorado State University, Bugwood.org



Howard F. Schwartz, Colorado State University, Bugwood.org





Matt Lavin

# Sedges

*Carex spp.*

## Tééht'oh



### Alternative Names:

None

### Plant Type:

Grasslike

### Habitat:

Grasslands

Ponderosa pine/mixed conifer

River valleys

### Growing Season:

Year-round


### Special Considerations:

None

**Sedges** are similar to grasses and rushes, but can be distinguished from either of these by their (usually) triangular cross section. One way to remember this is with the saying, “Sedges have edges, rushes are round, grasses have knees that bend to the ground” or “Sedges have edges, rushes are round, grasses are hollow right up from the ground.”

Not true grasses, sedges grow in a grasslike form. For more detail about one sedge species, see Yellow Nutsedge.

Species of sedge on the Navajo Endangered Species List: *Carex specuicola*, Navajo sedge ([Group 3](#) Navajo Endangered Species List)



Patrick Alexander

*Carex geophila*



Patrick Alexander

*Carex geophila*



Amy Smith Muise

# Algerita

*Berberis trifoliolata*

## Tsinłitsoi



### Alternative Names:

Agarita  
Agrito  
Currant-of-Texas  
Wild currant

### Plant Type:

Brush

### Flower Color:

Yellow

### Habitat:

Brushlands

### Growing Season:

Cool season

### Special Considerations:

- ✓ Edible berries
- ✓ Good for wildlife
- ! Mechanical injury

**Algerita**, a member of the barberry family, has spiny bluish or gray-green holly-like foliage with masses of fragrant yellow flowers in early spring, followed by attractive juicy red berries. Protected by sharp, spine-tipped leaves, its edible fruit is savored by wildlife and humans. Other similar Southwest native barberries include the larger *B. haematocarpa* and the smaller *B. swaseyi*.



**Algerita**

*Berberis trifoliolata*

**Tsinłitsoi**



Amy Smith Muese



Amy Smith Muese



Amy Smith Muese



Amy Smith Muese



Patrick Alexander

# Antelope Bitterbrush

*Purshia tridentata*

**Tsék'ina'ałch'ízhii**



## Alternative Names:

Buckbrush

Bitterbrush

## Plant Type:

Brush

## Flower Color:

White

Yellow

## Habitat:

Brushlands

Grasslands

## Growing Season:

Warm season

## Special Considerations:

✓ Good livestock forage

**Antelope bitterbrush** is an important native browse shrub for big game and livestock. It also provides shelter for small animals and birds. It is adapted to a wide range of soils with 8 to 34 inches of annual precipitation and occurs at elevations of 4,000 to 8,500 feet. It is normally 2 to 6 feet in height and up to 8 feet in width with wedge shaped, three-lobed leaves (some are persistent in winter). Branches near the soil may layer (develop roots where they touch the soil) providing additional rooting for the plant. Flowering occurs in late spring to early summer with yellow to white blossoms.



## Antelope Bitterbrush

*Purshia tridentata*

Tsék'ina'alch'ízhii



Patrick Alexander



Patrick Alexander

The yellow flowers of Antelope bitterbrush growing among other plants.



Patrick Alexander



Patrick Alexander





Patrick Alexander

# Biscuitroot

*Lomatium spp.*

## Nímasiichilí



### Alternative Names:

Foothill biscuit root

Giant biscuit root

Gray's biscuit root

### Plant Type:

Brush

### Flower Color:

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Provides wildlife forage and habitat

**Biscuitroot** is a perennial that grows up to three feet high and two feet wide. It has a strong, parsley-like odor. Growth begins very early in the spring, and it is an important forage plant for livestock, antelope, deer, and insects early in the season. It is also an important species in sage grouse habitat. The yellow flowers are found in umbels (umbrella-shaped clusters). The seeds are very similar to those of dill, and have papery wings, which are thin and membranous. Leaves of foothill biscuitroot are subdivided into numerous, fine segments, giving the plant a fern-like appearance. Leaves are bright green. Biscuitroots have no real stem. Instead, the leaves and flower stalks arise directly from the crown of an exceptionally thick taproot. Biscuitroots are often found in dry, open, rocky areas and are associated with poison hemlock, cow parsnip, piñon pine, snowberry, big sagebrush, black sagebrush, and Utah juniper. The root can be cooked, dried, and ground into a powder, which can then be mixed with cereal flours or added to soups, etc. Traditionally, it was often eaten in the winter when there was little other food available. The tender young stems can be eaten raw. Biscuitroot may also have been used as an aromatic flavoring in cooked foods.

*\*Based on Utah State University Range Plants of Utah description.*

**Biscuitroot**

*Lomatium* spp.

**Nímasiichilí**



Patrick Alexander



Patrick Alexander



Patrick Alexander



Joshua Tree National Park

# Blackbrush

*Coleogyne ramosissima*

## Ch'iltizhiní



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

Yellow

### Habitat:

Piñon-juniper woodland

Brushlands

### Growing Season:

Warm season

### Special Considerations:

None

**Blackbrush**, an aromatic shrub growing from 1 to 6 feet tall, forms nearly pure stands on large areas. Blackbrush occurs primarily in the transition zone between the Mojave and Great Basin deserts and on the western border of the Sonoran Desert. It occurs along the Colorado and San Juan river drainages in southeastern Utah and in adjacent drainages in the Mojave and Great Basin transition.

The plants are densely branched and spiny, but are browsed by goats and sheep. Its branches are opposite, short and rigid, becoming spiny with age. It flowers March to May and reproduces from seeds. The flowers are bell-shaped and generally yellow with bits of reddish-brown. The fruits contain only one seed, which has a feathery tuft. Fruit and seed formation occur only in years of good to abundant precipitation. Blackbrush can grow in very shallow soils, and it therefore has a shallow root system. One of its dispersal mechanisms is rodents. It is common to find burrows near these plants.

*\*Description based on Utah State University Range Plants of Utah description.*



**Blackbrush**

*Coleogyne ramosissima*

**Ch'illizhiní**



Andrey Zharkikh



Andrey Zharkikh



Stan Shebs [GFDL, CC BY-SA 3.0 or CC BY-SA 2.5], from Wikimedia Commons

# Buckwheat

*Eriogonum spp.*

## Bisnideeshchii'



### Alternative Names:

Sand buckwheat  
Slender buckwheat

### Plant Type:

Brush

### Flower Color:

White  
Yellow  
Pink

### Habitat:

Brushlands

### Growing Season:

Cool season

### Special Considerations:

None

**Buckwheat** is represented on the Navajo Nation by 28 different species. Most have taproots and one main stem (simple or branched and with or without leaves). Most species have their leaves near the base of the plant, but send a flowering stem up higher. Although, as forage, buckwheats are inferior, livestock and wildlife will graze upon the flowers. They seldom form patches or become dominant vegetation.

Buckwheat has been used medicinally to treat a variety of conditions and as a tea.

Below are the scientific and common names of buckwheats found on the Navajo Nation:

*Eriogonum alatum*, winged buckwheat  
*Eriogonum cernuum*, nodding buckwheat  
*Eriogonum clavellatum*, Comb Wash buckwheat  
*Eriogonum corymbosum*, crispleaf buckwheat  
*Eriogonum deflexum*, flatcrown buckwheat  
*Eriogonum divaricatum*, divergent buckwheat  
*Eriogonum effusum*, spreading buckwheat  
*Eriogonum gordonii*, Gordon's buckwheat  
*Eriogonum hieraciifolium*, hawkweed buckwheat  
*Eriogonum hookeri*, Hooker's buckwheat  
*Eriogonum inflatum*, desert trumpet  
*Eriogonum jamesii*, James' buckwheat  
*Eriogonum lachnogynum*, woollycup buckwheat  
*Eriogonum lachnogynum* var. *sarahiae*, Sarah's buckwheat ([Group 4](#) Navajo Endangered Species List)  
*Eriogonum leptocladon*, sand buckwheat  
*Eriogonum leptophyllum*, slenderleaf buckwheat  
*Eriogonum lonchophyllum*, spearleaf buckwheat  
*Eriogonum microthecum*, slender buckwheat  
*Eriogonum ovalifolium*, cushion buckwheat  
*Eriogonum pharnaceoides*, wirestem buckwheat  
*Eriogonum racemosum*, redroot buckwheat  
*Eriogonum scabrellum*, Westwater buckwheat  
*Eriogonum shockleyi*, Shockley's buckwheat  
*Eriogonum subreniforme*, kidneyshape buckwheat  
*Eriogonum umbellatum*, sulphur-flower buckwheat  
*Eriogonum viridulum*, clay hill buckwheat  
*Eriogonum wetherillii*, Wetherill's buckwheat  
*Eriogonum wrightii*, bastardsage



**Brush**

**Buckwheat**

*Eriogonum* spp.

**Bisnideeshchii'**



Patrick Alexander



Matt Lavin

*Eriogonum microthecum*, slender buckwheat



Patrick Alexander



Patrick Alexander



By Eitan F. [CC BY-SA 3.0 or GFDL], from Wikimedia Commons

# Camelthorn

*Alhagi camelorum*

## Ch'ilhoshí



### Alternative Names:

*Alhagi maurorum*

### Plant Type:

Brush

### Flower Color:

Red

Pink

Brown

### Habitat:

Brushlands

### Growing Season:

Cool season

### Special Considerations:

Noxious

**Camelthorn** is a dangerous noxious weed and one of the most difficult to eradicate. Livestock browse on camelthorn and may contribute to its spread.

Pulling or chopping is not an effective method of control, as this can encourage clonal reproduction by fragmenting underground rhizomes.

Its deep and extensive root system allows it to tap into a water table up to 50 feet below the surface. This allows camelthorn to thrive in areas of little rainfall and to out-compete native plants for necessary moisture and nutrients. Preventive measures include promoting the use of certified weed-free hay, not allowing livestock to eat and thus disperse the seeds, and not using heavy equipment in seed-infested areas.

In June and July, camelthorn produces short-stalked flowers, pink to magenta in color. During July and August, seedpods appear on the plant.

Camelthorn is a Middle East native eaten by camels in its native range. The sap contains mannitol and may be used as a laxative.



## Brush


### Camelthorn

*Alhagi camelorum*

### Ch'ilhoshí



Eitan F. [CC BY-SA 3.0 or GFDL], from Wikimedia Commons



Paul Hermann Wilhelm Taubert (1862-1897) [Public domain], via Wikimedia Commons



Lawrence Yazzie

Horse eating camelthorn along a road near Coalmine Canyon, AZ.





Russ Kleinman &amp; Richard Felger

# Chokecherry

*Prunus virginiana*

## Didzédik'ózhí



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

White

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

! Poisonous to livestock

**Chokecherry** is a native, deciduous, thicket-forming erect shrub or small tree. It rarely reaches a height of over 30 feet. The crown is irregular and from 10 to 20 feet wide when mature. The stems are numerous and slender. Reproduction can either be by seed or root rhizomes. Leaves are dark green and glossy above, and paler beneath, measuring 1 to 4 inches long and  $\frac{3}{4}$  to 2 inches wide. They are alternate, simple, glabrous, and oval to broadly elliptic in shape. The margins are toothed with closely-spaced, sharp teeth pointing outward, forming a serrated edge. They turn yellow in autumn.

Chokecherry foliage can be poisonous to grazing livestock. Hydrogen cyanide present in the leaves is converted to the toxic substance prunasin, either as a result of frost damage or as a result of livestock digestive processes. Results of a study indicated that prunasin concentrations are highest (5%) in the new stems and newly initiated leaves of chokecherry. Elevated levels are maintained in the leaves throughout the summer, but prunasin content of new twigs gradually diminishes over the season. The previous season's growth is generally not as toxic.

**Brush**

**Chokecherry**

*Prunus virginiana*

**Didzédik'ózhí**



*Russ Kleinman*



*Russ Kleinman Bill Norris, Richard Felger,  
& Leith Young*



*Russ Kleinman*



*Russ Kleinman & Richard Felger*



Patrick Alexander

# Cliff Fendlerbush

*Fendlera rupicola*

**K'iishzhiní**



## Alternative Names:

None

## Plant Type:

Brush

## Flower Color:

White

## Habitat:

Brushlands

Piñon-juniper woodland

Wastelands

## Growing Season:

Warm season

## Special Considerations:

None

**Cliff fendlerbush** is an upright, open, vase-shaped shrub with showy masses of fragrant flowers in spring. Flowers are white, cross-shaped, and conspicuous. Fendlerbush needs well-drained and rocky soils. It may become somewhat rangey without selective pruning. Its small, paired leaves are sparsely scattered on intricate branches. These plants are long-lived with a strong tap root.



**Cliff Fendlerbush**

*Fendlera rupicola*

**K'iishzhiní**



*Renee Benally*



*Renee Benally*



*Renee Benally*



Gerald Moore

# Cliffrose

*Purshia mexicana*

## Awééts'áál



### Alternative Names:

Formerly *Cowania neomexicana*

### Plant Type:

Brush

### Flower Color:

White

Yellow

### Habitat:

Brushlands

Piñon-juniper woodland

Wastelands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Provides wildlife forage

**Cliffrose** is found on cliffs, hillsides, mesas, and in washes, usually on the hot and dry south and west aspects, at elevations from 2,500 to 8,500 feet. This shrub or small tree is drought resistant, and mature plants are fairly shade tolerant.

Livestock may use it only lightly in spring and summer if deciduous browse species are available. It is an important browse species for mule deer, elk, pronghorn, desert bighorn sheep, livestock, and game birds. It also provides habitat for songbirds. It flowers April to June, and fruits mature September to October. It reproduces from seed, and rarely from sprouting. The flowers are cream to yellow in color.



Renee Benally



Patrick Alexander

# Four-Winged Saltbush

*Atriplex canescens*

## Díwózhiitbéíí



### Alternative Names:

Chamiza

Fourwing saltbush

Fourwinged saltbrush

### Plant Type:

Brush

### Flower Color:

Red

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Good livestock forage

**Four-winged saltbush** loses its leaves in drought. It has grayish-white to pale green leaves. Mature plants range from 1 to 8 feet in height, depending on ecotype, soil, and climate, but it is most common on bottomland sites. Its root system can reach depths of up to 20 feet when soil type allows.

Among the most preferred shrubs of the Southwest, its leaves, stems, flowers, and seeds are used by all livestock except horses. This is a nutritious plant. It has a good rating for sheep and goats; fair for cattle.

Fourwing saltbush often has male and female flowers on separate plants. Male flowers are red to yellow and form dense spikes at the ends of the branches. The female flowers are axillary and nondescript. The seed is contained in a winged sac that turns a dull yellow when ripe and may remain attached to the plant throughout winter.

Species of saltbush on the Navajo Endangered Species List: *Atriplex garrettii* var. *navajoensis*, Navajo Saltbush ([Group 4](#) Navajo Endangered Species List)



## Brush

### Four-Winged Saltbush

*Atriplex canescens*

Díwózhiitbéíí



Amy Smith Muise

Seed in winged sac.



Amy Smith Muise



Amy Smith Muise

Four-winged saltbush in winter.



Amy Smith Muise

Four-winged saltbush in summer.



Matt Lavin

# Greasewood

*Sarcobatus vermiculatus*

## Díwózhiishjiin



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

Green

### Habitat:

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

! Potentially toxic to livestock

**Greasewood** grows 3 to 10 feet tall. It is an important winter browse plant for domestic sheep, cattle, and big game animals. It also gets some use during spring and summer months.

Flowers are green. Fruit are green while growing, turning reddish nearing the end of development, and tan when fully mature.



## Brush

### Greasewood *Sarcobatus vermiculatus* Díwózhiishjiin



Matt Lavin



Gerald Moore



Matt Lavin



Matt Lavin





Elizabeth Sohn

# Mormon Tea

*Ephedra spp.*

## Tł'ohozihii



### Alternative Names:

Tall Mormon tea  
Longleaf ephedra  
Joint Fir  
Green ephedra

### Plant Type:

Brush

### Flower Color:

Purple  
Green  
White

### Habitat:

Brushlands  
Wastelands

### Growing Season:

Cool season

### Special Considerations:

! Potentially toxic to  
sheep and cows

**Mormon tea** is found on dry, rocky, open sites in valleys and washes, and on slopes, alluvial fans, mesas, and foothills. It is typically found at elevations ranging from 3,000 to 7,500 feet. Sites supporting Mormon tea receive average precipitation of 6 to 15 inches. Mormon tea is drought resistant and winter hardy. It grows primarily on sandy, gravelly, or rocky, well-drained, undeveloped soils.

It has numerous parallel stems that point upward resembling a broom, with branchlets clustered around nodes.

It has low forage value, but is frequently browsed by domestic livestock. It is an important browse species for big game, especially on winter range, since green ephedra stems and twigs extend above the snow.

Ephedra can be toxic to both domestic sheep and cows during gestation, even at low doses. It causes ruminal impaction, diarrhea, and in some cases death.

The stems of ephedra were traditionally brewed to make a nonmedicinal beverage as well as a medicinal tea considered to be a remedy for a backache. People have also made flour and a coffee-like beverage from the seeds.

## Brush

### Mormon Tea

*Ephedra* spp.

Tł'ohozihii



Patrick Alexander



Amy Smith Muise



Amy Smith Muise



H. Zell [GFDL or CC BY-SA 3.0], from Wikimedia Commons

# Oregon Grape

*Berberis aquifolium*

Tséch'ilndtł'izíyilt 'áą 'í



## Alternative Names:

Oregon grape holly

Mahonia

Formerly *Mahonia aquifolium*

## Plant Type:

Brush

## Flower Color:

Yellow

## Habitat:

Brushlands

## Growing Season:

Warm season

## Special Considerations:

None

**Oregon grape** is native to parts of the western United States. It is not a grape or a holly, but rather a type of barberry. It grows to a height of 3 to 6 feet and has dark, glossy leaves with spines along the leaf margins. It produces purple berries, which are a source of food for wildlife. The berries were traditionally used for food and dye.



## Oregon Grape

*Berberis aquifolium*

Tséch'ilndtł'izíyilt 'áą 'í



I, Hugo.org [GFDL or CC BY-SA 3.0],  
from Wikimedia Commons



H. Zell [GFDL or CC BY-SA 3.0],  
from Wikimedia Commons



H. Zell [GFDL or CC BY-SA 3.0],  
from Wikimedia Commons



Jean Tosti [GFDL or CC BY-SA 3.0],  
from Wikimedia Commons



Kurt Stüber [GFDL or CC BY-SA 3.0], from Wikimedia Commons



Patrick Alexander

# Pingue

*Hymenoxys richardsonii*

## Ch'ıl̨ı́báhiláhtahtsoi



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ! Very toxic to sheep
- ! Toxic to cattle

**Pingue** is common lower elevation perennial sub-shrub or herb native to much of the west. Pingue is similar to rubberweed in appearance. It grows to a height of just under two feet, with thick, woody stalks. The leaves are mostly basal and are highly dissected. The flowerheads are raised above the leaves on long stems. The inflorescence has yellow rays and a yellow disc, much like sunflowers, to which pingue is related. It flowers June to September. Pingue has bits of wooly growth both at stem bases and near the crown of the plants. It likes rocky hillside habitats at middle elevations.

Pingue is not palatable to livestock, but under forage shortage may be readily eaten by sheep and goats. It is toxic to livestock, particularly sheep, though cattle also can be affected. It is especially toxic to sheep. The best control is good grazing management.

Its stems and leaves have been used as a tea. Its roots have a milky sap that contains latex, and has been used as chewing gum. The stems and leaves have also been used as a poultice for wounds.



**Pingue**

*Hymenoxys richardsonii*

**Ch'illibáhilátahtsoi**



Patrick Alexander



Patrick Alexander

Pingue is the small green plant with yellow flowers on the bare ground in the foreground of this picture.





Patrick Alexander

# Rabbitbrush

*Ericameria nauseosa*

## Ch'ildiilyésiits'óóz



### Alternative Names:

Rubber rabbitbrush

### Plant Type:

Brush

### Flower Color:

Yellow

### Habitat:

Brushlands

### Growing Season:

Cool season

### Special Considerations:

- ✓ Useful for soil stabilization and restoration

**Rabbitbrush** is a rounded shrub generally 2 to 5 feet tall. It is often found growing along roadsides.

Livestock generally forage only lightly on this species, and it is considered to be of little value to all classes of livestock. It can be an important browse species for mule deer, pronghorn, and jackrabbits during fall and winter. It also provides cover for mammals and small nesting birds.

Rabbitbrush thrives in poor conditions and can tolerate coarse, alkaline soils. The species is useful in soil stabilization and restoration of disturbed sites. The deep root system establishes quickly, and plants produce large quantities of leaf litter, helping to bring nutrients to the soil surface from the deeper rooting profile. Its yellow flowers bloom from August to October.

It has been used as a yellow dye, to make a medicinal tea, and for chewing gum.

Species of *Ericameria* on the Navajo Endangered Species List: *Ericameria arizonica*, Rydberg's thistle ([Group 4](#) Navajo Endangered Species List)

## Brush

### Rabbitbrush *Ericameria nauseosa* Ch'ildilyésiits'óóz



Russ Kleinman



Russ Kleinman



Patrick Alexander



Patrick Alexander

# Rabbitbrush, Green

*Chrysothamnus viscidiflorus*

## Ch'ildiilyésiits'óóz



### Alternative Names:

Yellow rabbitbrush

Douglas rabbitbrush

### Plant Type:

Brush

### Flower Color:

Green

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

! Potentially invasive

**Green rabbitbrush** has little value as a forage species.

An erect shrub, branching from near the base, with a rounded crown, 1 to 3½ feet tall, it flowers July to September and reproduces from seed, and vegetatively by vigorous sprouting.

Sheep and cattle occasionally use it for browse when other feed is not available. Deer browse lightly on it in the summer and winter. Elk utilize it in the winter. It can be a somewhat weedy plant, increasing where there has been serious damage to the more desirable forage.

It quickly and aggressively invades disturbed, open sites including burns and overgrazed rangelands.

People sometimes chew the roots of rabbitbrush as gum. It contains rubber, especially when growing in alkali soils.



**Rabbitbrush, Green**

*Chrysothamnus viscidiflorus*



Patrick Alexander



Patrick Alexander



Patrick Alexander



Matt Lavin

# Roundleaf Buffaloberry

*Shepherdia rotundifolia*

Dibédáá'



## Alternative Names:

Silverleaf

## Plant Type:

Brush

## Flower Color:

Yellow

## Habitat:

Piñon-juniper woodland

Ponderosa pine/mixed conifer

## Growing Season:

Year-round

## Special Considerations:

- ✓ Wildlife forage and habitat

**Roundleaf buffaloberry** is found only in southern Utah and northern Arizona. As its name indicates, the foliage is silvery green, and the leaves remain on the small shrub year-round. It provides habitat and fodder for quails and other small animals. The plants are drought tolerant, and can often be found growing on rocky slopes or outcrops.



**Brush**

**Roundleaf Buffaloberry**

*Shepherdia rotundifolia*

**Dibédąą'**



Matt Lavin



Matt Lavin



Matt Lavin





J.S. Peterson, hosted by the USDA-NRCS PLANTS Database

# Sagebrush, Big

*Artemisia tridentata*

## Ts'ahtsoh



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

White

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Excellent wildlife forage

**Big sagebrush** and its subspecies are tall, rounded, native shrubs with short, branched, woody trunks. They are normally about four feet high, but vary from two feet in arid conditions to as high as 15 feet on favorable sites. Big sagebrush is perhaps the most important shrub on western rangelands. Evergreen leaves and abundant seed production provide an excellent winter food source to numerous species of large mammals including mule deer, black-tailed deer, white-tailed deer, elk, pronghorn antelope, bighorn sheep and jackrabbits.

Big sagebrush can be distinguished by its aromatic leaves and stems. The scientific name, *Artemisia tridentata*, provides another useful clue: the three-toothed leaf tips. The leaves are grayish and slightly hairy or fuzzy. The trunks may form interesting and twisty shapes.

Sagebrush prefers dry soils, and will not grow at stream edges. It is used for tea, rope, medicine, and fuel.

**Sagebrush, Big**

*Artemisia tridentata*

**Ts'ahtsoh**



*Patrick Alexander*



*Morris Southward, PhD*



*Patrick Alexander*



*Patrick Alexander*



George Miller

# Sagebrush, Bigelow

*Artemisia bigelovii*

**Ts'ahłibáhí**



## Alternative Names:

Flat sagebrush

## Plant Type:

Brush

## Flower Color:

Green

## Habitat:

Brushlands

Grasslands

## Growing Season:

Warm season

## Special Considerations:

- ✓ Wildlife and  
livestock forage

**Bigelow sagebrush** is adapted to xeric sites and is one of the most drought-tolerant sagebrushes in North America. Bigelow is relatively small compared to other sagebrushes. It grows in canyons, draws, and on washes, plains, hills, and rimrock on well-drained soils (usually sandy or gravelly); it is common on limestone soils. Its overall elevational range is 3,000 to 8,000 feet. Bigelow sagebrush provides valuable winter and spring forage for wildlife and livestock and is an important item in the fall diet of pronghorn in northern Arizona. Palatability and nutritional value are high relative to other sagebrush species, because its twigs are less woody, and its leaves less bitter, with a milder odor, than associated sagebrush taxa. Bigelow sagebrush begins new growth in April. Flowerbuds appear in August, and flowering occurs from August to October. The leaves abscise in winter.



**Sagebrush, Bigelow**

*Artemisia bigelovii*

**Ts'ahlibáhi**



*George Miller*



*George Miller*



*Ken Gishi*



Nick Ashcroft

# Sagebrush, Black

*Artemisia nova*

## Tsétah Ts'ah



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

Yellow

Green

### Habitat:

Brushlands

River valleys

Wastelands

### Growing Season:

Warm season

### Special Considerations:

None

**Black sagebrush** is a low growing evergreen shrub with a flat-topped crown. Mature plants range from 4 to 12 inches tall. The leaves are dark green, sometimes grayish, and triangular in outline with three rounded lobes. The leaf surface is often covered with hair, which distinguishes this species from big sagebrush (*A. tridentata*). These hairs give the leaves a darker appearance.



## Brush

### Sagebrush, Black

*Artemisia nova*

Tsétah Ts'ah



Matt Lavin



Matt Lavin



Matt Lavin



Nick Ashcroft





Matt Lavin

# Sagebrush, Fringed

*Artemisia frigida*

## Tóyikáál

**Alternative Names:**

Prairie sagewort

**Plant Type:**

Brush

**Flower Color:**

Green

**Habitat:**

Grasslands

**Growing Season:**

Warm season

**Special Considerations:**

None

**Fringed sagebrush** is a spreading shrublet ranging from approximately 4 to 16 inches in height. It is pleasantly fragrant, whitish or grayish tomentose, and arising from a tough, woody crown. Tomentose means having dense, velvety, fuzzy hairs. The leaves are also tomentose and abundant, clustered toward the base of the plant and scattered along the stem. The lower leaves are petiolate, about ½ inch long, with the upper leaves becoming sessile. The inflorescence is a panicle with small, greenish flower heads. Fringed sagebrush flowers from July to August. The fruits are dry, smooth, broadly cylindrical achenes.

## Brush

### Sagebrush, Fringed

*Artemisia frigida*

Tóyikáál



Matt Lavin



Matt Lavin



Matt Lavin



Matt Lavin



Patrick Alexander

# Sagebrush, Sand

*Artemisia filifolia*

## Ch'ilzhóó'



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

Brown

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ! Potentially toxic to horses
- ✓ Provides erosion control

**Sand sagebrush** is a freely branching, rounded crown, aromatic shrub growing up to four feet tall. It flowers August to September, and reproduces from seeds. Flowers are numerous and small. The hairy, bluish-green leaves are alternate and found in bundles. The bark eventually exfoliates in thin shreds. Sand sagebrush is found on dunes, sand hills, and deep sands. It is generally considered an indicator of sandy soil and is usually abundant in those areas, at elevations between 3,000 and 5,000 feet.

It is rarely used by livestock and wildlife in grasslands, where more preferred forage species are available, but may be used heavily in arid desert regions. It may cause sage sickness in horses (a temporary, intoxicated state, with loss of coordination).

Sand sagebrush provides hiding or thermal cover for numerous smaller birds and mammals and is extremely effective in preventing wind erosion on light, sandy soils. A canopy of sand sagebrush can afford some protection to grasses such as needle-and-threadgrass on heavily grazed sites.

Traditional uses include preparing a decoction of its leaves to treat intestinal worms and other stomach problems.



## Brush

### Sagebrush, Sand

*Artemisia filifolia*

Ch'ilzhóó'



Patrick Alexander



Patrick Alexander



Amy Smith Muise



Patrick Alexander



George Miller

# Shadscale

*Atriplex confertifolia*

## Dá'ák'óózhdiíoolí



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

Yellow

### Habitat:

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Livestock and wildlife forage

**Shadscale** is a low-growing shrub (1 to 3 feet) with spine tipped branches. It occurs in arid climates at elevations from 4,000 to 7,000 feet. It grows on dry slopes, flat areas, ridges, and valley bottoms.

Shadscale can form almost pure stands in some locations. It is resistant to overgrazing and is drought tolerant. Tolerance to drought is achieved through partial shedding of leaves; this reduces water loss during severe moisture stress.

Shadscale is used by all classes of livestock, as well as by mule deer, for forage. It is used mostly during winter and spring for browse. Late spring or summer grazing, especially if intensive, is injurious. The fruits provide food for game birds and songbirds, and the seeds of shadscale remain on the plant throughout the winter, enhancing its nutritional value.

Traditional uses include grinding the fruits of shadscale into flour.



**Shadscale**

*Atriplex confertifolia*

**Dá'ák'óózhdiíoolí**



*George Miller*



*Nick Ashcroft*



*Nick Ashcroft*





Patrick Alexander

# Skunkbush

*Rhus trilobata*

## Ch'iltichiin



### Alternative Names:

Skunkbrush

### Plant Type:

Brush

### Flower Color:

White

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

None

**Skunkbush** occurs in a variety of habitats at elevations of 4,500 to 8,000 feet, including dry, rocky slopes, along streams and canyon bottoms, waste places, pastures, roadsides, and on sand dunes. It is drought resistant, shade tolerant, and intolerant of flooding. It typically grows where maximum annual precipitation ranges from 10 to 20 inches.

It has spreading woody rhizomes, and it sprouts readily from both the root and crown after disturbance.

Skunkbush's forage value is poor for all classes of livestock. Skunkbush provides some browse for deer, elk, and pronghorn when other more preferred forage is unavailable. In most locations, big game use tends to be heaviest during the winter when food supplies are most limited.

Because the fruit of skunkbush persists through the fall and winter, this species can provide a ready food source for birds and small mammals when other foods are scarce or unavailable.

Traditionally, the fruits were used in food and medicines, and in the preparation of lemonade-like beverages. Pliable young stems were woven into durable baskets. Skunkbush can also be used in making dyes for clothing.

*\*Description adapted from Utah State University's Range Plants of Utah.*

**Brush**

**Skunkbush**

*Rhus trilobata*

**Ch'ililichiin**



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander

# Snakeweed

*Gutierrezia sarothrae*

## Ch'il Diilyésii dzaa



### Alternative Names:

Broom snakeweed

### Plant Type:

Brush

### Flower Color:

Green

Yellow

### Habitat:

Brushlands

### Growing Season:

Year-round

### Special Considerations:

! Potentially toxic to livestock

**Snakeweed** is a woody subshrub with numerous erect ascending branches. It has small, evergreen, linear leaves and tufts of tiny yellow flowers in fall. Though snakeweed was once considered a weed of overgrazed rangelands, some research suggests that its population cycles over time are more related to precipitation patterns than to overgrazing.

Broom snakeweed provides little value to livestock and can be toxic, causing illness, death, or abortion in sheep, goats, and cattle during winter and early spring when it is the only thing green and may be consumed in large quantities.



**Snakeweed**

*Gutierrezia sarothrae*

**Ch'il Diilyésii dzaa**



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Rob Routledge, Sault College,  
Bugwood.org

# Snowberry

*Symphoricarpos albus*

## Didzé



### Alternative Names:

Mountain snowberry

Utah snowberry

Waxberry

Ghostberry

Formerly *Symphoricarpos* spp.

### Plant Type:

Brush

### Flower Color:

White

Pink

### Habitat:

Ponderosa pine/mixed conifer

Brushlands

River valleys

### Growing Season:

Cool Season

### Special Considerations:

- ✓ Early spring forage

**Snowberry** is a low-growing shrub whose habit can be either erect or trailing. Its branches are spreading or arching. It averages 2 to 4 feet in height, but plants on good sites can grow up to five feet. Reproduction is by seed and sometimes by layering. Fruits are small, light green to white berries, up to ½ inch long. Older twigs have reddish-brown, shreddy bark. Buds are light brown.

Snowberry occurs at elevations of 4,800 to 10,500 feet, at the edges of riparian zones, in woodlands, and in moist areas of mountain brush zones. It occurs in sandy loam to clay loam soils and does not tolerate much alkalinity or salinity. Associated species include Ponderosa pine, Douglas fir, aspen, and chokecherry. Because of its abundance and wide distribution, snowberry is an important source of forage on many mountain ranges. Although not highly nutritious or palatable, mountain snowberry is frequently one of the first species to leaf out, making it a highly sought after food in the early spring. Use by livestock and big game is moderate throughout the summer, but declines in fall. Small mammals and birds utilize the fruits. Snowberry's low growth form makes its foliage easily available. Plants withstand browsing well and produce numerous basal sprouts following browsing.

Due to its rhizomatous nature, snowberry is useful for revegetation of disturbed sites such as road cuts. It is also an excellent species for wildlife habitat improvement and landscape or recreational plantings. Its fruits have been used as an emetic and laxative. Roots are steeped to treat colds and stomachaches.

*\*Description courtesy of Utah State University's Range Plants of Utah.*

**Snowberry**  
*Symphoricarpos albus*  
**Didzé**



*Rob Routledge, Sault College, Bugwood.org*



*Rob Routledge, Sault College, Bugwood.org*





Patrick Alexander

# Wax Currant

*Ribes cereum*

K'ínijitahí



## Alternative Names:

None

## Plant Type:

Brush

## Flower Color:

Red

## Habitat:

Brushlands

Ponderosa pine/mixed conifer

## Growing Season:

Warm season

## Special Considerations:

- ✓ Forage and habitat for small wildlife

**Wax currant** occurs on dry, open slopes, ridges, rock outcrops, and mountain shrub communities, as well as at forest edges or in coniferous forests with light canopy cover. In addition, it occurs in the Bebb willow (*Salix bebbiana*) community type in the northwestern third of New Mexico. It is found at elevations of 4,950 to 13,200 feet.

Wax currant provides food and cover for wildlife. It is only fair to poor browse for deer, but it is important on ranges where little else is available.

Chickadees and other birds consume the fruit of wax currant. The fruit of wax currant is used for making jam, jelly, or pie. It has also been used for making pemmican. Wax currant is cultivated as an ornamental. It grows from 1½ to 5 feet tall. Wax currant reproduces mainly by seed.

**Brush**

**Wax Currant**

*Ribes cereum*

**K'ínijitahí**



*Russ Kleinman*



*Russ Kleinman*



*Russ Kleinman*



*Patrick Alexander*



Russ Kleinman &amp; Karen Blisard

# Wild Rose

*Rosa woodsii*

## Chq̄q̄h



### Alternative Names:

None

### Plant Type:

Brush

### Flower Color:

Red

Pink

White

### Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

River valleys

### Growing Season:

Cool season

### Special Considerations:

✓ Erosion control

**Wild rose** is a shrub that grows 2 to 5 feet tall, usually forming thickets. It reproduces from seed, rhizomes, sprouting, and layering. Flowers are found in clusters, with five petals, five sepals, and many stamens. Seeds are contained within an orange-red, round hip, which generally stays on the bush throughout the winter.

Wild rose occurs on prairies, plateaus, dry slopes, and in open woods, ravines, and thickets, growing at elevations from 3,500 to 7,500 feet. It is a common riparian species. It is fairly tolerant of browsing. It can thrive in moderate shade to full sunlight.

Wild rose is adapted to a wide range of soil types and textures. Growth is generally best on moderately fertile, well-drained clay loam, sandy loam, or sandy soils. It is also adapted to a broad range of moisture conditions but tends to favor moist, well-drained soils that are present in riparian ecosystems.

Wild rose is browsed by livestock and big game from spring through fall. They prefer this shrub in the spring when the leaves appear. Porcupines and beavers also browse the leaves. Wild rose hips persist on the plant through much of the winter. Many birds and mammals are sustained by these dry fruits when the ground is covered with snow.

The young shoots have been used as a potherb and the leaves steeped for tea, petals eaten raw, in salads, candied, or made into syrup. The inner bark was smoked like tobacco, and dried petals stored for perfume. Rosehips are a source of vitamin C.

*\*Description courtesy of Utah State University's Range Plants of Utah.*




## Brush

### Wild Rose

*Rosa woodsii*

### Chq̣h



Russ Kleinman & Bill Norris



Russ Kleinman & the Thursday hiking group



Russ Kleinman & Bill Norris



Russ Kleinman



Patrick Alexander

# Winterfat

*Krascheninnikovia lanata*

## Gahtsohdáá'



### Alternative Names:

White sage

*Ceratoides lanata*

*Eurotia lanata*

Formerly *Eurotia ceratoides*

### Plant Type:

Brush

### Flower Color:

White

### Habitat:

Brushlands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Forage for sheep and wildlife

**Winterfat** is a low-growing, long-lived (up to 130 years) subshrub with a woody base and numerous annual branchlets, growing 1 to 3 feet tall. It has a hairy, silvery-white appearance. Winterfat occurs in dry valley bottoms, on flat mesas, and on hillsides, at elevations between 2,400 and 9,300 feet. It is drought resistant and intolerant of flooding, excess water, or acidic soils.

Winterfat is good forage for sheep, pronghorn, elk, mule deer, and many small mammals and birds. It is fair forage for cattle. It is most valued as winter forage. It will decrease under heavy, continuous grazing.

Grazing season can have more influence on winterfat than grazing intensity. Late winter or early spring grazing is most detrimental. Winterfat can tolerate winter use of 50% if it is rested occasionally. Spring and summer use should not exceed 35%. Early winter grazing may actually be beneficial.

It flowers April to September and reproduces from seed and sprouting. Sprouting of the buds near the plant base occurs when the plant is browsed or damaged.

Winterfat has been used as a hair wash, and a decoction from the leaves has been used to treat fevers.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



**Winterfat**

*Krascheninnikovia lanata*

**Gahtsohdáá'**



Amy Smith Muisse



Patrick Alexander



Patrick Alexander



Patrick Alexander

Winterfat flowers are white but occasionally have a red or green tint.





Patrick Alexander

# Wolfberry

*Lycium spp.*

## Haasch'éehdáá'



### Alternative Names:

Water jacket

Pale wolfberry (one species)

Anderson wolfberry  
(another species)

### Plant Type:

Brush

### Flower Color:

Purple

### Habitat:

Brushlands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

None

**Wolfberry** is a spiny, rounded, branched shrub, 1 to 9 feet tall, with numerous slender spines and flattened, thick, fleshy leaves. It may be drought deciduous, meaning it loses foliage in response to low moisture availability. Its fleshy red berries contain many seeds.

Wolfberry frequently provides a source of berries and browse in areas of degraded habitat. The berries also are eaten by humans, who have also used other parts of the plant as medicine and in ceremonials.

In times of famine, the Native Americans of northern Arizona ate the dried berries mixed with saline clay. In this form they were called “food clay” by the Navajo or “potato clay” by the Hopi. As well as being an extender, the clay may have been used to reduce bitterness. With the recent interest in using native plants, more palatable recipes for sauces and jams using pale wolfberry berries are available. Medicinal uses include applying the soaked leaves to cuts and using the ground-up roots for toothaches. These hardy shrubs have also been used as ornamentals.

Stands of pale wolfberry are frequently associated with ancient Anasazi ruins in the Four Corners area. Some stands have been speculated to be deliberate plantings and others the result of inadvertent seed dispersal. After a pueblo is abandoned, moisture tends to collect over decomposing subsurface floors and plazas, and these conditions may make suitable habitat for seeds that were unintentionally dropped by former inhabitants.



Patrick Alexander



Patrick Alexander



Andrey Zharkikh

# Field Bindweed

*Convolvulus arvensis*

Ch'il natł'oi łigai



## Alternative Names:

None

## Plant Type:

Vine

## Flower Color:

Red

Pink

White

## Habitat:

Brushlands

## Growing Season:

Warm season

## Special Considerations:

None

**Field Bindweed** is native to Europe and Asia. It is common in cultivated fields and gardens, along railroads and roadsides, and in disturbed sites and waste places. Outside of agricultural fields, it is most often found in moist locations on tracts once used for agriculture. The competitive ability of field bindweed is due largely to its extensive root system. One plant is able to reduce the available soil moisture in the top 24 inches of soil to below the wilting point. Field bindweed has deep roots that store carbohydrates and proteins. They help field bindweed spread vegetatively and allow it to resprout repeatedly following removal of above-ground growth.



Andrey Zharkikh



Amy Smith Muise





Patrick Alexander

# Puncturevine

*Tribulus terrestris*

Ch'ilhoshíq

Naakaibihosh



## Alternative Names:

Goathead

## Plant Type:

Vine

## Flower Color:

Yellow

## Habitat:

Brushlands

## Growing Season:

Warm season

## Special Considerations:

- ! Toxic to sheep
- ! Invasive

**Puncturevine** (also called goathead), an introduced, invasive plant, is sensitive to competition and will be outcompeted by native species except on repeatedly disturbed areas.

The foliage of this plant is toxic, and grazing animals may eat burs, which cause injuries to the mouth, stomach, and intestines. Ants seem to congregate under the plants, particularly near stem emergence.

The burs are painful to hands or feet of humans and can penetrate shoes or bicycle tires. Clear clothing, shoes, and tires of burs to avoid spreading the plant to new locations.

Stems form a dense mat with small yellow flowers on short stalks. Puncturevine reproduces by seed, so controlling plants prior to seeding is important.

**Puncturevine**

*Tribulus terrestris*

**Ch'ilhoshíq Naakaibihosh**




Patrick Alexander



Amy Smith Muise



Patrick Alexander



Kathryn-Mae Eiland



Russ Kleinman

# Antelopehorn

*Asclepias asperula*

## Azee'ła'dilt'éhé



### Alternative Names:

Green-flowered milkweed

Spider antelope horns

### Plant Type:

Forb

### Flower Color:

White

Green

### Habitat:

Grasslands

### Growing Season:

Warm season

### Special Considerations:

! Poisonous to livestock

**Antelopehorn** is a very common roadside milkweed. The leaves are lanceolate and up to six inches long. The corolla lobes are greenish white and the hoods are purple.

For further description of milkweeds in general, see the milkweed entry.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*



**Antelopehorn**  
*Asclepias asperula*  
**Azee'la'dilt'éhé**



Patrick Alexander



Russ Kleinman



Russ Kleinman



Patrick Alexander

# Beeplant, Rocky Mountain

*Cleome serrulata*

Waa'



## Alternative Names:

Navajo spinach

## Plant Type:

Forb

## Flower Color:

Pink

Purple

## Habitat:

Brushlands

Wastelands

## Growing Season:

Warm season

## Special Considerations:

None

**Beeplant** is an erect, branched annual that grows to about three feet. tall. It is in flower from July to August, and reproduces from seed. Dozens of bright, pink to purplish flowers are crowded into rounded or rather elongate spikes. Fruit is a narrow capsule up to two inches long that bears several to many dark, ovoid seeds. Leaves are in groups of three on long petioles. Rocky Mountain beeplant is often found in wastelands, and in lower mountains. It is frost tender. It can grow in semi-shade or no shade and is often found on sandy soil. It has little to no value as forage for livestock. However, its nectar-filled blossoms nourish pollinators, including bees, butterflies, wasps, and hummingbirds.

Rocky Mountain beeplant has been used traditionally as food and medicine. Seeds can be harvested and eaten raw or cooked, dried and ground into a meal, then used as a mush or mixed with flour to make bread. The plant can be used to treat fevers and stomach disorders. A poultice made from the pounded, soaked leaves has been applied to sore eyes. A decoction of the leaves has been used as a body and shoe deodorant.

A black dye is obtained by boiling down the whole plant. It is used as paint for decorating pottery.

*\*Description based on Utah State University Range Plants of Utah description.*

**Beeplant, Rocky Mountain**

*Cleome serrulata*

**Waa'**



*Patrick Alexander*



*Patrick Alexander*





Russ Kleinman & Karen Blisard

# Birdbeak

*Cordylanthus wrightii*

**Tsésnááá'**



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

Red

Pink

## Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

## Growing Season:

Warm season

## Special Considerations:

- ✓ Provides insect forage

**Birdbeak** is native to Texas, Colorado, Arizona, New Mexico, and Utah, and can be found associated with piñon pine. It is an annual with divided leaves and yellow flowers. (Flowers may also be pink.) It provides forage for bees, butterflies, moths, and flies. The leaves and flowers are covered in fine hairs and never fully open. It is found on dry hillsides at middle elevation. It normally grows from 10 to 30 inches tall. In the Four Corners area, birdbeak blooms in August, September, and October.

**Birdbeak**

*Cordylanthus wrightii*

**Tsésnááá'**



*Russ Kleinman & Karen Blisard*



*Russ Kleinman & Karen Blisard*



*Russ Kleinman & Karen Blisard*



Patrick Alexander

# Bitterweed

*Hymenoxys spp.*

Ch'ilbílatahłitsxoidíchi'i'í



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

## Habitat:

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

River valleys

## Growing Season:

None

## Special Considerations:

- ! Toxic to livestock, especially sheep

**Bitterweeds** are members of the aster family that grow up to five feet tall. There are several species of bitterweed on Navajo rangelands. Like other asters, the flowers have ray flowers (what look like outer petals) and disk flowers (tiny flowers in the dense center of the flower). Both ray flowers and disk flowers are in the yellow to orange spectrum. Bitterweed is toxic to livestock, but affects sheep more than other animals. Due to the bitter taste of the plant, animals generally do not eat it when there are other plants to eat.

*For descriptions of specific species, consult the entries on pingue and rubberweed.*



**Bitterweed**

*Hymenoxys* spp.

**Ch'ilibílatahłitsxoidíchi'i'í**



Patrick Alexander

*Hymenoxys vaseyi*



Patrick Alexander

*Hymenoxys odorata*



Patrick Alexander

*Hymenoxys richardsonii*



Brian Muise



Patrick Alexander

# Blanket Flower

*Gaillardia spp.*

## Ch'ilbílatahózhóón



### Alternative Names:

Blanketflower

### Plant Type:

Forb

### Flower Color:

Red

Pink

Yellow

### Habitat:

Grasslands

### Growing Season:

Warm season

### Special Considerations:

None

**Blanket flower** is named for its resemblance to a brightly patterned blanket. It grows 1 to 3 feet tall, and will often grow in mounds 8 to 18 inches high. It has wiry, branched stems, mostly without leaves. It flowers June through October with bright, daisy-like, single color and bi-color blooms in shades from buff to red to brown. Flowers of certain species can be four inches wide. The ray petals have serrated tips that are usually three-toothed. Some varieties have a fringed look. The base of the flower (pappus) consists of several scales with awns.

The seeds are similar to sunflower seed, but smaller and at least partly covered with long hairs. The leaves are basal and are lance- to linear-shaped. They have a hairy texture, and can have margins which are smooth, toothed, or lobed.

Found on the dry slopes and meadows of uplands and mountains, blanket flower is drought tolerant and capable of growing under very harsh conditions. It prefers full sun.

Not a highly grazed species, when grazed it is most often grazed during the early part of the season.

It is often cultivated and grown in flower beds.

*\*Description based on Utah State University Range Plants of Utah.*



**Blanket Flower**

*Gaillardia* spp.

**Ch'ilbílátahózhóón**



*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*





Russ Kleinman, Bill Norris, Kelly Kindscher, and Danielle Walkup

# Blue-Eyed Grass

*Sisyrinchium* spp.

Azeetł'ohí



## Alternative Names:

Arizona blue-eyed grass

## Plant Type:

Forb

## Flower Color:

Purple

Yellow

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

None

**Blue-eyed grass**, despite its name, is not always blue. The name refers to a number of plants in the iris family, some of which are yellow. Blue-eyed grass has typical iris leaves, which are folded toward the base. The flower is large, with petals over  $\frac{5}{8}$  inch long. This plant lives on rocky slopes at upper elevations.

Species found in the Four Corners region include *Sisyrinchium demissum* (stiff blue-eyed grass), which is blue, and *Sisyrinchium arizonicum* (Arizona blue-eyed grass), which is yellow.

**Blue-Eyed Grass**

*Sisyrinchium* spp.

**Azeetł'ohí**



Russ Kleinman & Richard Felger  
*Sisyrinchium demissum*



Russ Kleinman, Bill Norris, Kelly Kindscher, and Danielle Walkup  
*Sisyrinchium arizonicum* growth habit



Russ Kleinman, Bill Norris, Kelly Kindscher, and Danielle Walkup  
*Sisyrinchium arizonicum* with flower and capsules



Russ Kleinman, Bill Norris, Kelly Kindscher, and Danielle Walkup  
*Sisyrinchium arizonicum*



Patrick Alexander

# Bluebells

*Campanula rotundifolia*

**Dahiitiíhídą́' dootł'izhí**



## Alternative Names:

American bellflower

Bluebell bellflower

Harebell

## Plant Type:

Forb

## Flower Color:

Purple

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

None

**Bluebells** are herbaceous perennials, growing up to 18 inches tall. There are one to several very slender stems, hairless to slightly hairy, and containing a milky sap. The basal stems are creeping. They reproduce by seed and by the spreading of rhizomes, often forming colonies. They flower from May to August. The bell-shaped blue flowers measure  $\frac{1}{2}$  to 1 inch and are slightly nodding, with short lobes. The flower stalk arises six inches above dark green foliage. The flowers may be solitary, but up to 15 flowers may be attached up the stem. Seeds are small, with three capsules. There are two different types of leaves present. Basal leaves are round, forming a rosette. They have a distinct leaf stalk (petiole), are broad, and decrease late in the season. The leaves on the erect part of the stem are long and narrow with alternate attachment. There are 10 to 22 leaves,  $\frac{3}{8}$  to 3 inches long and  $\frac{1}{32}$  to  $\frac{5}{8}$  inch wide; leaf bases and petioles are hairless. All leaves are dark green.

The plant is found from low meadows to subalpine regions ranging from 5,000 to 12,000 feet in elevation. It is adapted to full or partial sunlight and tolerates drought. It grows best in rich, well-drained, moist soil. It can tolerate a range of soil pH, and can thrive in acid and calcareous areas. Bluebells are often used as a plant in flower gardens. This plant does reseed itself quite vigorously, and can become aggressive and weedy.

*\*Description based on Utah State University's Range Plants of Utah description.*



**Bluebells**

*Campanula rotundifolia*

**Dahiitíhídą́' doot'izhí**



Patrick Alexander



Patrick Alexander



Russ Kleinman

# Bundle Flower

*Desmanthus cooleyi*

K'ei'chilí



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

Pink

White

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

None

**Bundle flower** is a prostrate or weakly ascending herb with spherical white flowerheads. The leaves are bipinnately divided, and the stems are unarmed. *Desmanthus cooleyi* is found in dry areas in canyons and mesas. It can be difficult to differentiate *Desmanthus cooleyi* from *Calliandra humilis* (the dwarf stick pea), especially since *Desmanthus cooleyi* dries to a pink purple color similar to *Calliandra*. The fruits are the key—they are much thinner in *Desmanthus*.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*

**Bundle Flower**

*Desmanthus cooleyi*

**K'ei'chilí**



Russ Kleinman



Russ Kleinman



Russ Kleinman





*Ixitxel (eigene Arbeit, selbst fotografiert)  
[GFDL or CC-BY-SA-3.0], via Wikimedia  
Commons*

# California Burclover

*Medicago polymorpha*

Tł'oh azee



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

## Habitat:

Brushlands

Grasslands

## Growing Season:

Warm season

## Special Considerations:

! Noxious

**Burclover** is a shallow-rooted annual legume that generally has numerous prostrate stems branching from the crown and spreading outward 6 to 30 inches. Where thick stands develop, however, stems may become erect, obtaining heights of 18 to 24 inches in favorable years. The leaves are subglabrous and clover-like in appearance, with leaflets normally wedge-shaped and toothed toward the top. The inflorescence is usually quite limited, presenting only a few small, yellow, pea-like flowers. The several-seeded fruit is a flattened, coiled pod, commonly up to ¼ inch in width and fringed with a double row of conspicuous, hooked spines. Well developed plants may produce more than 1,000 pods. The seed is rather large for a legume of this type, usually developing to over ¾ inch in length.

## Forbs

### California Burclover

*Medicago polymorpha*

Tł'oh azee



Forest & Kim Starr [CC BY 3.0], via  
Wikimedia Commons



Forest & Kim Starr [CC BY 3.0], via  
Wikimedia Commons



Forest & Kim Starr [CC BY 3.0], via  
Wikimedia Commons



Forest & Kim Starr [CC BY 3.0], via  
Wikimedia Commons



Nick Ashcroft

# Cocklebur

*Xanthium saccharatum*

## Tá'niits'éhii



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Green

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ! Causes injury
- ! Potentially invasive

**Cocklebur** is an annual plant that infests disturbed areas such as roadsides, ditches, fields, pastures, orchards, riparian areas, and waste areas. Stems grow up to three feet tall. They are tough, with three-branched, yellowish spines measuring ½ to 1 inch long along the leaf axils. The leaves are lanceolate, the underside densely covered with white, wooly hairs. The leaf margins may be entire, toothed, or lobed (with lobes 1 to 4 inches long) with one main vein from the base. Male and female flowers develop in separate heads on the leaf axils of the same plant: Male flower heads, are small and green, and develop in clusters. Female flower heads develop on the leaf axils below the male flower heads singly or in small clusters; spiny female flower heads consist of two flowers that lack corollas and become hardened prickly burs that enclose two seeds at maturity. Manual removal is effective, especially before burs develop.

DO NOT cut and leave the plants with immature burs on the site because they can still develop viable seed. Some herbicides are effective.



**Cocklebur**  
*Xanthium saccharatum*  
**Tá'niits'éhii**



*Nick Ashcroft*



*Nick Ashcroft*



Forest and Kim Starr

# Common Purslane

*Portulaca oleracea*

## Tsiigháįłchíí



### Alternative Names:

Little hogweed

Red root

Pursley

### Plant Type:

Forb

### Flower Color:

Yellow

### Habitat:

None

### Growing Season:

Warm season

### Special Considerations:

✓ Edible leaves

! Weedy

**Purslane** is a common weedy herb frequently found in gardens and on sidewalks. It grows in a spreading, prostrate fashion and has small yellow flowers. The leaves are oval and succulent, on orange to red stems. The flowers are small and yellow. The entire plant is edible and has been used in cuisine around the world. Common purslane can be eaten as a leaf vegetable and can be consumed either raw or cooked.



**Common Purslane**

*Portulaca oleracea*

**Tsiigháįłchíí**



Russ Kleinman



Mourad Louadfel, Bugwood.org



Mourad Louadfel, Bugwood.org



Phil Westra





Max Licher, SEINet - Arizona Chapter

# Copperweed

*Oxytenia acerosa*

K'iiltsoinitł'izí



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

## Habitat:

River valleys

Wastelands

## Growing Season:

Warm season

## Special Considerations:

- ! Poisonous to livestock
- ! Potential skin irritant

**Copperweed** is a woody-based forb, growing 3 to 5 feet tall, with many slender gray stems. The plant turns and stays copper colored through summer. It grows in semidesert climate in summer precipitation areas. It is most likely to be found along streambeds and gullies with extra moisture. It is usually on soils high in salt. Cattle, sheep, and elk may eat copperweed when other feed is scarce. Cattle are more likely to eat this plant in the fall when they are being trailed. Copperweed may cause poisoning in livestock. Symptoms of poisoning include dullness, weakness, and coma followed by death without a struggle. In humans, copperweed may cause skin irritation.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



SEINet - Arizona Chapter, 2018, <http://swbiodiversity.org/seinet/index.php>



Patrick Alexander

# Cow Parsnip

*Heracleum maximum*

## Azee'haagai



### Alternative Names:

Indian celery

Indian rhubarb

Pushki

### Plant Type:

Forb

### Flower Color:

White

### Habitat:

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

✓ Provides wildlife and livestock forage

! Potential skin irritant

**Cow parsnip** is a native, perennial forb that grows from 3 to 10 feet tall and has broad, flat-topped umbels.

Cow parsnip grows throughout the United States and Canada. It occurs in a wide variety of forested habitat types, as well as grassland, shrubland, meadow, alpine, and riparian zones. In the Intermountain West, it is a common understory species in subalpine fir and Engelmann spruce habitat types, as well as in quaking aspen communities. It is commonly found growing in snow-maintained disclimaxes such as avalanche chutes. Cow parsnip is a facultative wetland species; it grows best in moist, shaded areas but can also be found in open woodlands and clearings.

Cow parsnip is a valuable forage species for livestock, deer, elk, moose, and bear. It is rated moderately good for erosion control, short-term revegetation potential, and long-term revegetation potential. It has fair soil stabilization value.

Cow parsnip has traditional medicinal and culinary uses. Some cultures ate the inside of stems raw, and boiled the roots to extract sugar. Others ate the young leaves and stems, and used the roots to treat epilepsy.

Some with sensitive skin develop rashes when contact with cow parsnip is followed by exposure to sunlight.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



**Forbs**

**Cow Parsnip**  
*Heracleum maximum*  
**Azee'haagai**




*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*





Matt Lavin

# Curlycup Gumweed

*Grindelia squarrosa*

## Ch'ibílátahaltsóíí



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Yellow

### Habitat:

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

None

**Curlycup gumweed** is a short-lived perennial or biennial forb averaging 1 to 3 feet high. Numerous branching stems bear alternately arranged leaves, typically 1 to 4 inches long with entire to serrate—or even somewhat lobed—margins. The flower heads are radiate with 25 to 40 yellow rays and a yellow center. The involucre bracts are strongly rolled back, and highly resinous. The fruit is an achene about  $\frac{1}{8}$  inch long bearing 2 to 3 awns. Flowering occurs in mid- to late-summer, typically beginning in July and continuing through August and into September. Occasional plants will be seen with flowers persisting into November.

**Curlycup Gumweed**

*Grindelia squarrosa*

**Ch'ibílátahaltsóí**



Matt Lavin



Matt Lavin



Matt Lavin



Matt Lavin



Montana Statewide Noxious Weed  
Awareness

# Daisy, Rock

*Perityle emoryi*

## Nidíyíliitigaii



### Alternative Names:

Emory's rock daisy

### Plant Type:

Forb

### Flower Color:

Yellow

White

### Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

### Growing Season:

Warm season

### Special Considerations:

None

The rock daisy is native, herbaceous annual that may grow in clumps to a height of two feet. The leaves are spade shaped with toothed margins. The flowers are white with yellow centers.



**Daisy**

*Perityle emoryi*

**Nidíyíliitigaii**



Chris Evans, University of Illinois,  
Bugwood.org



David Stephens, Bugwood.org



Becca MacDonald, Sault College, Bugwood.org



John Ruter, University of Georgia,  
Bugwood.org



Russ Kleinman

# Dandelion

*Taraxacum officinale*

## Bééshyilt'áq'í



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Yellow

### Habitat:

Brushlands

Grasslands

River valleys

### Growing Season:

Cool season

### Special Considerations:

- ✓ Provides wildlife and livestock forage

**Dandelion** is a forb 3 to 12 inches tall, with erect stems that contain a bitter, milky-white juice. It has a fleshy, deep taproot. Flowers are yellow, located on the tip of the leafless stem. It becomes a ball of white, silky bristles at maturity. Seeds of dandelion are topped by a parachute of bristles that aid in dispersal. Basal, deeply lobed leaves are 2 to 16 inches long, and form a rosette. Surfaces of leaves can be lightly pubescent.

Dandelion is widespread throughout North America. It most commonly occurs in disturbed areas such as cut-over or burned forests, avalanche areas, overgrazed ranges, and marshy floodplains, but it may also occur on well-managed ranges. It also occurs on highway and railroad rights-of-way, waste places, old fields, pastures, and lawns. It grows at elevations between 500 and 11,000 feet. Dandelion can reproduce vegetatively by sprouting from the caudex after disturbance. It is found on a variety of soils, but is most common in heavy loams and sandy loams.

Dandelions provide fair to good forage for livestock and wildlife, and are readily eaten because they are relatively succulent. Antelope, as well as sage and forest grouse, use it heavily.

Young leaves can be eaten as spring greens. Roots can be ground and used as a coffee substitute, mild laxative, or to treat heartburn. Tea and wine can be made from the flowers. Flowers can be fried in batter and eaten.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



**Dandelion**

*Taraxacum officinale*

**Bééshyilt'áą'í**



*Russ Kleinman*



*Russ Kleinman*



*Russ Kleinman*





Matt Lavin

# Death Camas

*Zigadenus paniculatus*

## Yííłtsíniitsoh



### Alternative Names:

Foothill deathcamas

### Plant Type:

Forb

### Flower Color:

White

Yellow

### Habitat:

Brushlands

Wastelands

### Growing Season:

Cool season

### Special Considerations:

! Poisonous to livestock

**Death camas** is a forb growing 1 to 2 feet tall, with an underground scaly bulb. Growth begins very early in the spring, before most other plants. Death camas is most abundant in foothill areas. Reproduction is by seed.

Flowers are white to yellowish, with orange stamens that give them a soft, lacy quality. They appear in early summer. Rough brown seeds are formed in a three-celled capsule, which is especially poisonous. Plants have five or six basal, thickened, V-creased leaves with a grasslike appearance.

Death camas is found from the semidesert to mountain climatic zones in a wide range of soils. All parts of this plant contain a poisonous alkaloid at all growth stages. Sheep and cattle often eat death camas in early spring before other plants start producing forage. Sheep are the most commonly poisoned, but cattle deaths have also been reported. Respiratory problems occur in sheep after eating ½ to 2 pounds of death camas. Signs of poisoning include salivation, nausea and vomiting, weakness, coma, and death within a few hours to two days. Pastures containing death camas should not be grazed until late spring when other forage is available. The bulbs, which can be mistaken for onions or sego lily, can cause severe illness in humans.

*\*Courtesy of Utah State University Range Plants of Utah.*

Species of death camas on the Navajo Endangered Species List: *Zigadenus vaginatus*, alcove death camas ([Group 3](#), Navajo Endangered Species List)

**Death Camas**  
*Zigadenus paniculatus*  
**Yíłtsíniitsoh**



Matt Lavin



Matt Lavin



Matt Lavin



Matt Lavin





Patrick Alexander

# Desert Sand Verbena

*Abronia spp.*

K'íneeshdlíshiidaá'



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Blue

Purple

Red

Pink

## Habitat:

Grasslands

Wastelands

## Growing Season:

Warm Season

## Special Considerations:

None

**Sand verbena** is native to the western United States. There are a number of species classified as sand verbena, and they have a range of bloom colors and are generally fragrant. Flowers occur clustered in an umbel. Leaves are slightly fleshy and round rather than elongated.

Desert sand verbena grows on sandy flats, dunes, and roadsides.



Patrick Alexander

*Abronia elliptica*





Patrick Alexander

# Devilweed

*Chloracantha spinosa*

Ni' it tł' hii



## Alternative Names:

Aster

*Aster spinosus*

*Spiny chloracantha*

Buena mujer

## Plant Type:

Forb

## Flower Color:

White

## Habitat:

Grasslands

Ponderosa pine/mixed conifer

## Growing Season:

Warm season

## Special Considerations:

None

**Devilweed** is a member of the aster family, native to the southwestern United States. Like a number of other asters, it has many-branched stems and flowers that actually comprise ray flowers (what look like standard petals) and disk flowers (tiny flowers in the center of the inflorescence). Devilweed has white ray flowers and yellow disk flowers. It can grow to be several (4 to 5, or even 6) feet tall. It grows on a variety of sites from wet areas to semi-arid soils.

**Devilweed**

*Chloracantha spinosa*

**Ni' ı́ t' hii**



*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*



Joy Viola, Northeastern University,  
Bugwood.org

# Evening Primrose

*Oenothera spp.*

Tł'é'yiigáhii



## Alternative Names:

Tufted evening primrose  
Stemless primrose  
Fragrant evening primrose  
White stemless  
evening primrose  
Gumbo lily

## Plant Type:

Forb

## Flower Color:

White  
Yellow

## Habitat:

Piñon-juniper woodland

## Growing Season:

Warm season

## Special Considerations:

None

**Evening primrose's** flowers open late in the day, then turn pink and wilt the following morning. It can grow up to one foot tall, is nearly stemless, and grows from a thick taproot. It blooms April to September.

The flowers are white or yellow with four heart-shaped petals and eight yellow stamens. Flowers can measure up to three inches across and have a wonderful fragrance.

Rough seedpods, about one inch long, form soon after flowering.

Leaves grow in a basal rosette. They are lance-shaped and toothed, growing up to 12 inches in length. The leaves are crinkly, gray-green, and fuzzy.

Evening primrose is found in piñon-juniper woodlands and shadscale scrub communities. It is common on roadsides, and is often abundant on steep, dry slopes. It occurs at elevations of 2,980 to 9,500 feet. It likes well-drained, rocky, sandy soils.

Cattle seem to avoid evening primrose. Young shoots and roots of some evening primroses are eaten by humans.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



## Evening Primrose

*Oenothera* spp.

Tł'é'yiigáhii



Joseph M. DiTomaso,  
University of California - Davis



Karan A. Rawlins, University of Georgia, Bugwood.org



Becca MacDonald, Sault College, Bugwood.org



Robert Vidéki, *Doronicum Kft.*,  
Bugwood.org



Patrick Alexander

# Filaree

*Erodium cicutarium*

## Dahjtiíhídáá'łikaní



### Alternative Names:

Storksbill  
Redstem filaree  
Alfilaria  
Redstem stork's bill

### Plant Type:

Forb

### Flower Color:

Blue  
Purple  
Red  
Pink

### Habitat:

Brushlands  
Grasslands  
Wastelands

### Growing Season:

Cool season

### Special Considerations:

- ✓ Provides wildlife and livestock forage
- ! Invasive
- ! Noxious

**Filaree** is a low and spreading plant, 2 to 5 inches tall, growing from a central taproot. The stems are leafy and hairy. Filaree flowers February to May, and plants usually dry up and disappear quickly after maturity. It is one of the first plants to germinate in late fall or spring. It reproduces from seeds. Borne on hairy stalks in umbrella-shaped clusters, flowers vary in color from pink to purple. Each seed is tipped with an elongated tail, which coils spirally at maturity, assisting the pointed seed in penetrating the soil. In early growth stages, the leaves form only a basal rosette, but later appear on the stems.

Filaree is found in oak woodlands, semidesert grassland, and desert shrublands. It is often found in fields, lawns, and wastelands. It is adapted to a broad range of soil types. It grows in well-drained, clayey, loamy, or sandy soil, and is tolerant of moderately acidic to moderately alkaline soils. It furnishes excellent to good spring forage for cattle, sheep, desert tortoise, and other wildlife. It can also provide winter forage if the seeds germinate following fall rains. It has, however, been reported to cause bloating in livestock.

Young leaves can be eaten raw or cooked. Filaree is reputed to contain an antidote for strychnine. The presence or absence of filaree pollen in fossil records, sediment lakebeds, and artifacts has been used as a dating technique in paleobotany and archeology. Filaree was one of the first exotics to invade North America. It is an aggressive invader of desert ranges under heavy grazing.

*\*Description courtesy of Utah State University Range Plants of Utah.*



## Forbs

### Filaree

*Erodium cicutarium*

Dahjítíhídą́'íkaní



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander





Patrick Alexander

# Four O'Clock

*Allionia incarnata*

K'íneeshdlíshiidáá'łigai



## Alternative Names:

Trailing windmills

Guapile

## Plant Type:

Forb

## Flower Color:

Red

Pink

White

Yellow

## Habitat:

None

## Growing Season:

Warm season

## Special Considerations:

None

**Four o'clock** is a prostrate perennial herb which is glandularly pubescent. The flowers are pink and notched. The leaves are wedge shaped. The three flowers of four o'clock in each whorl all flower at the same time and are very close together, giving the illusion of one large completely symmetrical flower. Actually, the three flowers are each individually bilaterally symmetrical. The complicated fruit of four o'clock has two incurved wings that are frequently triangularly toothed, and it has two rows of glands on its surface. Four o'clock is found in low, dry, rocky areas.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*

Four O'Clock

*Allionia incarnata*

K'íneeshdlíshiidáá'ligai



Patrick Alexander

*Allionia incarnata* var. *villosa*



Patrick Alexander

*Allionia incarnata* var. *incarnata*



Patrick Alexander

*Allionia incarnata* var. *villosa*



Patrick Alexander

*Allionia incarnata* var. *incarnata*



Patrick Alexander

*Allionia incarnata* var. *incarnata*



Patrick Alexander

# Geranium

*Geranium spp.*

## Dahjítííhídáá'tsó



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Purple

Red

### Habitat:

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

River valleys

### Growing Season:

Cool season

### Special Considerations:

None


**Geranium** may be found in grasslands and in wooded areas, growing to a height of four feet. While some geranium species are introduced, there are many natives as well. Species found in Utah and New Mexico include Fremont geranium (*Geranium caespitosum*), sticky geranium (*Geranium viscosissimum*), Richardson's geranium (*Geranium richardsonii*), and sticky purple geranium (*Geranium viscosissimum*). Geraniums provide good livestock forage, particularly for sheep. Deer also graze the plants. Flowers have five parts (petals, stamens) and display a range of colors in the lavender-pink spectrum. The leaves are palmately lobed and hairy. Stems are also hairy. The sticky geranium has edible flowers and leaves. The Cheyenne used the leaves of Richardson's geranium medicinally and made a tea out of the roots.



**Geranium**

*Geranium spp.*

**Dahjítíhídáá'tsó**



Patrick Alexander

*Geranium caespitosum*



Patrick Alexander

*Geranium caespitosum*



Patrick Alexander



Patrick Alexander

*Geranium caespitosum*



Patrick Alexander

# Gilia

*Gilia spp.*

## Azééhaaleeh



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Red

Pink

### Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

River valleys

### Growing Season:

Warm season

### Special Considerations:

- ✓ Provides hummingbird forage

**Gilia** is an herbaceous annual that can grow to heights of five feet, though the plants more commonly grow to about three feet. The plants provide food for hummingbirds and hawkmoths. The tubular flowers vary in color, but shy gilia (*Gilia inconspicua*) has pink blooms, and scarlet gilia (*Ipomopsis aggregata*/*Gilia aggregata*) has flowers in deep shades of salmon and red. The stems are hairy (and sometimes sticky), as are the pinnate leaves. The scarlet gilia has many traditional medicinal uses, including treatment of colds, rheumatism, and skin irritation.

Gilia is a member of the phlox family, and gilia species are native to most of the western United States and a few pockets in the east.

Species of saltbush on the Navajo Endangered Species List: *Aliciella formosa*, Aztec gilia ([Group 4](#) Navajo Endangered Species List)

**Gilia**

*Gilia spp.*

**Azééhaaleeh**



Patrick Alexander

*Ipomopsis aggregata*



Patrick Alexander

*Ipomopsis aggregata*



Matt Lavin

*Gilia inconspicua*



Matt Lavin

*Gilia inconspicua*





Patrick Alexander

# Globemallow

*Sphaeralcea spp.*

## Azeenitł'inii



### Alternative Names:

Scarlet Globemallow

*Sphaeralcea coccinea*

### Plant Type:

Forb

### Flower Color:

Red

### Habitat:

Grasslands

### Growing Season:

Cool season

### Special Considerations:

- ✓ Wildlife and livestock forage

**Globemallow**, a perennial, native forb that grows up to 2½ feet tall. It is commonly eaten by almost all species of herbivores and is an important part of the diets of small mammals, pronghorn, sheep, and cattle.

Scarlet globemallow grows mainly in dry grassland prairies at elevations of 3,500 to 9,000 feet. It is considerably drought resistant and establishes well on disturbed sites. It loses its leaves during times of drought and may actually increase in size in times of drought and overgrazing. Traditional uses include applying the paste to burns, scalds, and external sores as a cooling agent.

It flowers April to August, and reproduces from seeds. Petals are deep orange or brick red to pinkish.

*\*Description courtesy of Utah State University's Range Plants of Utah.*

**Globemallow**

*Sphaeralcea* spp.

**Azeenit'inii**



Patrick Alexander



Gerald Moore



Nick Ashcroft

*Sphaeralcea coccinea*



Nick Ashcroft

*Sphaeralcea coccinea*



Patrick Alexander



Renee Benally



Russ Kleinman

# Groundsel, Threadleaf

*Senecio flaccidus* var. *flaccidus*

## Azeehááldzidí



### Alternative Names:

Wooly Groundsel

Formerly *Senecio longilobus*

### Plant Type:

Forb

### Flower Color:

Yellow

### Habitat:

Brushlands

Grasslands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

Toxic to livestock

**Threadleaf groundsel** typically grows in dry, gravelly, or hardpan soils, and may be found on plains and foothill areas. It flowers in the late summer.

Often called woolly groundsel, it contains pyrrolizidine alkaloids that may poison cattle, horses, and sheep. All parts of these plants are poisonous to all classes of livestock, but younger plants tend to be more palatable to livestock. Young animals are more susceptible.

If liver damage occurs, animals may survive for six months or longer after they have ingested a lethal amount of the plant, and may show no outward symptoms during this period. Often in response to some stressful situation, such as pregnancy or lactation, they may develop liver failure. The best preventive is avoidance of the plant.



**Groundsel, Threadleaf**

*Senecio flaccidus* var. *flaccidus*

**Azeehááldzidí**



Russ Kleinman



Morris Southward, PhD



Matt Lavin

# Halogeton

*Halogeton glomeratus*

## Chil'bit'ááh t'ó



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Green

Red

White

### Habitat:

Brushlands

### Growing Season:

Cool season

### Special Considerations:

- ! Noxious
- ! Invasive
- ! Poisonous, especially to sheep

**Halogeton** is an undesirable weed on rangeland. It was introduced into North America in about 1930 and has rapidly spread.

In larger quantities, it is poisonous, and sheep are the most susceptible. It contains toxic amounts of sodium, potassium, and calcium oxalates. Signs of poisoning include: depression, weakness, reluctance to move, drooling, coma, and death. The first signs of poisoning occur 2 to 6 hours after an animal ingests a fatal amount, and death occurs in 9 to 11 hours.

Halogeton cannot compete with healthy range plants. Therefore, control involves keeping a healthy cover of desirable forage plants.

## Forbs

### Halogeton

*Halogeton glomeratus*

Chil'bit'áąh t'ó



Matt Lavin

Halogeton flowers



Matt Lavin





Patrick Alexander

# Horsemint

*Monarda spp.*

## Kétłoh



### Alternative Names:

Bee balm  
Mountain mint  
Oregano de la Sierra

### Plant Type:

Forb

### Flower Color:

Blue  
Purple

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

- ✓ Provides good insect forage
- ✓ Hummingbirds like it

**Horsemint**, in the mint family, has square stems and attractive flowers. Several species in the *Monarda* genus can be found regionally.

*Monarda punctata* has been used as part of a mixture to treat colds, ground into a powder and snuffed up the nostrils to relieve a headache, or as an infusion to treat fevers. It may be hung up in dwellings for its pleasing odor. Birds, including hummingbirds, are attracted to the nectar.

## Forbs

## Horsemint

*Monarda spp.*

## Kétłoh



Clarence A. Rechenthin @ USDA-NRCS  
PLANTS Database, Courtesy of USDA  
NRCS Texas State Office



Patrick Alexander



Patrick Alexander

# Indian Paintbrush

*Castilleja spp.*

## Dahiitjįhídą́ą'tsoh



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Green

Yellow

### Habitat:

Wastelands

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

None

**Indian paintbrush**, is admired for its aesthetic value, due to its showy bracts. It grows 9 to 18 inches tall.

Palatability of Indian paintbrush has been rated as fair to good for sheep, poor to fair for cattle, and poor for horses.

Indian paintbrush is widespread, occurring from the desert floors to the subalpine meadows. Some species are known to be partial root parasites, especially with sagebrush. It reproduces by seed and tillering.



**Indian Paintbrush**

*Castilleja spp.*

**Dahiitį́hídą́'tsoh**



Nick Ashcroft

*Castilleja linarifolia*



Patrick Alexander

*Castilleja integra*



Nick Ashcroft

*Castilleja integra*



Nick Ashcroft

*Castilleja linarifolia*



Russ Kleinman

# Iris

*Iris missouriensis*

## Bílatahyígaii



### Alternative Names:

Blue flag  
Flag lily  
Liver lily

### Plant Type:

Forb

### Flower Color:

Purple  
White  
Yellow

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

None

**Western blue flag**, the iris species found in Navajo rangelands, is a perennial herb, usually evergreen and growing from a creeping, tuberous rhizome. Rhizomes on blue flag are between 1 to 1½ inches in diameter, quite large for a native iris. The leaves are ½ to ¾ inch wide, long and linear with parallel venation. They are sometimes purplish. Stems are 8 to 20 inches tall and sometimes branched. Blossoms of this iris species are pale lilac to whitish with lilac-purple veins.

The roots have been used to treat toothaches and to make a green dye.



**Iris**

*Iris missouriensis*

**Bílátahyígaii**



*Russ Kleinman*



*Russ Kleinman & Bill Norris*



*Russ Kleinman & Bill Norris*



*Russ Kleinman & Bill Norris*



*Russ Kleinman & Bill Norris*





Russ Kleinman, Richard Felger &  
Kelly Kindscher

# Jacob's Ladder

*Polemonium foliosissimum*

**Azee'naachii'**



## Alternative Names:

Jacobs ladder

## Plant Type:

Forb

## Flower Color:

Purple

Yellow

## Habitat:

Grasslands

Ponderosa pine/mixed conifer

## Growing Season:

Warm season

## Special Considerations:

None

**Jacob's ladder** is an upper elevation herb similar except for flower color to *Polemonium flavum*, yellow Jacob's ladder, which is found in upper elevation mixed conifer forests. It is upright and has a compound leaf, the lanceolate leaflets. The flowers are lavender.

**Jacob's Ladder**  
*Polemonium foliosissimum*  
**Azee'naachii'**



*Russ Kleinman, Richard Felger, & Kelly Kindscher*



*Russ Kleinman, Richard Felger, & Kelly Kindscher*



*Russ Kleinman, Richard Felger, & Kelly Kindscher*



*Russ Kleinman, Richard Felger, & Kelly Kindscher*



Patrick Alexander

# Jimmyweed

*Isocoma pluriflora*

## K'iłtsołizhiní



### Alternative Names:

Goldenweed

Rayless goldenrod

Southern jimmyweed

Southern goldenbush

*Haplopappus heterophyllus*

*Haplopappus pluriflorus*

*Isocoma wrightii*

### Plant Type:

Forb

### Flower Color:

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ! Poisonous to livestock
- ! May cause illness in humans from meat or milk

**Jimmyweed** is native to Colorado, Arizona, New Mexico, and Texas. It is poisonous to livestock, and it may also cause illness in humans who drink milk or eat meat from poisoned animals. Jimmyweed is common on roadsides, disturbed areas, and pastures. The plant is a member of the aster family. It grows from 1 to 2½ feet in height, with about the same width. The multiple stems and flowers on each plant give it a shrubby appearance. Flowers appear in summer.



**Jimmyweed**  
*Isocoma pluriflora*  
**K'íiltsoilizhiní**



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Russ Kleinman

# Jimsonweed

*Datura spp.*

## Ch'óhojilééh



### Alternative Names:

Sacred thorn-apple

### Plant Type:

Forb

### Flower Color:

Blue

Purple

White

### Habitat:

Grasslands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

- ! Poisonous to humans and livestock
- ! Potential skin irritant

**Jimsonweed** is a large forb, 1¼ to 4 feet tall, often with a similar spread. It often falls over from its own weight. It germinates readily and self sows under most conditions. Flowers are showy white to purple trumpets 2 to 4 inches long. Flowers open for only one evening, but new ones continue to open throughout the summer and autumn.

Mature leaves are elliptical to egg-shaped, uneven at the margins. They are 3 to 8 inches long, six inches wide, on long, stout petioles. Margins of leaves have a few large triangular teeth. Jimsonweed prefers warm areas. It competes aggressively for water and grows rapidly. It is a common weed in pastures, barnyards, roadsides, and waste places.

Jimsonweed and its derivatives have several medicinal uses. At low doses, it is used to treat asthma, muscle spasms, and symptoms of Parkinson's disease. At higher doses it causes hallucinations. It is occasionally grown as an ornamental, and attracts bees, butterflies, and moths.

All parts of jimsonweed are very poisonous. Cattle and sheep have died from eating it. Symptoms include dilated pupils, thirst, fever, loss of coordination, confusion, rapid pulse, labored respiration, hallucinations, convulsions, and eventual coma. Even inhaling the sweet fragrance of the flowers can cause headaches and dizziness. The sap can cause a skin rash. Species of *Datura* on the Navajo Nation include *D. wrightii* and *D. stramonium*.



**Jimsonweed**  
*Datura stramonium*  
**Ch'óhojilééh**



*Russ Kleinman*



*Patrick Alexander*



*Russ Kleinman & Richard Felger*



*Russ Kleinman*



*Jeanne Gleason*





Gerald Moore

# Knapweed

*Centaurea spp.*

## Ch'it Bilat'a dootłizhigí



### Alternative Names:

Spotted knapweed

Yellow starthistle

### Plant Type:

Forb

### Flower Color:

Red

Pink

Yellow

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ! Invasive
- ! Noxious
- ! Spiny

**Knapweed** is represented on Navajo rangelands by several species, including Russian knapweed, diffuse knapweed, and squarrose knapweed, which are described in their own entries.

Other species of *Centaurea* include yellow starthistle (*Centaurea solstitialis*) and spotted knapweed (*Centaurea biebersteinii*). All knapweeds are introduced species in the Four Corners region, and they are often invasive or noxious.

Spotted knapweed is a multi-stemmed, herbaceous perennial that can grow from 1 to 3 feet in height. It produces bright pink blooms in the summer. Two things make it a particularly invasive plant: it produces many seeds whose viability can last for years, and it also grows very thickly, crowding out other species. It also has a degree of fire resistance due to its taproot and hardy seeds. It does, however, provide forage for domestic and wild animals.

Yellow starthistle is likewise a problematic species. The rosette of yellow starthistle lies close to the ground and produces 6 to 28 deeply lobed leaves that range from 1 to 8 inches long and are generally less than two inches wide. The rigidly branched, winged flowering stems average about two feet tall, but range from four inches to five feet, depending on environmental conditions. Flower heads are solitary on the ends of short stems and have many bright yellow flowers. The bracts of the flower heads are armed with stout, straw-colored spines 1 to 2 inches long that radiate from the flowerhead in a star shape.

**Knapweed**

*Centaurea* spp.

**Ch'it Bilat'a dootlizhigí**



Gerald Moore



Matt Lavin



Matt Lavin



Patrick Alexander

*Centaurea solstitialis*



Patrick Alexander

*Centaurea solstitialis*



Matt Lavin

# Knapweed, Diffuse

*Centaurea diffusa*

## Ch'il lat'á dei nínigí



### Alternative Names:

White knapweed  
Spreading knapweed  
Tumble knapweed

### Plant Type:

Forb

### Flower Color:

White

### Habitat:

Brushlands  
Grasslands  
Piñon-juniper woodland  
Ponderosa pine/mixed conifer

### Growing Season:

Warm season

### Special Considerations:

- ! May injure livestock
- ! May decrease range value
- ! Introduced
- ! Noxious

**Diffuse knapweed** is a nonnative, annual, biennial, or short-lived perennial forb. During the juvenile stage, diffuse knapweed is a rosette with deeply divided basal leaves borne on short stalks on a central crown with a taproot. At maturity, diffuse knapweed produces one upright stem, rarely two. Stems are 1 to 3 feet tall, with numerous, spreading branches that give the plant a ball-shaped appearance and tumbleweed mobility when broken off. Flowerheads are solitary or borne in clusters of two or three at the ends of branches. Diffuse knapweed flowerheads are  $\frac{1}{8}$  to  $\frac{2}{8}$  inch (3 to 6 mm) in diameter and  $\frac{5}{16}$  to  $\frac{7}{16}$  inch long, excluding spines and flowers. Bracts are edged with a fringe of spines, with a longer ( $\frac{1}{16}$  to  $\frac{3}{16}$  inch), erect spine at the tip.

Diffuse knapweed replaces traditional wildlife and livestock forage on rangelands and pasturelands. While it is not poisonous, the presence of diffuse knapweed in hay or on rangeland can decrease feeding value to livestock and wildlife species. In situations of overgrazing or drought, when fewer forage species are available, the flower shoots are sometimes grazed while immature rosettes are not. Although rosettes of the first year's growth are nutritious and edible, they are difficult for cattle to eat because they are so close to the ground. Mature knapweed plants are coarse and fibrous, and the spines on the bracts can be very irritating or may even cause injury to the mouths and digestive tracts of grazing animals. Diffuse knapweed is grazed by deer and domestic sheep, and by elk and cattle, at least through the bolting stage.



**Knapweed, Diffuse**

*Centaurea diffusa*

**Ch'il lat'á dei nínigí**



Matt Lavin



Matt Lavin



Matt Lavin



Patrick Alexander

# Knapweed, Russian

*Acroptilon repens*

## Ch'ildích'l'itibáhí



### Alternative Names:

Turkestan thistle

Creeping knapweed

Mountain bluet

Russian cornflower

Hardheads

Formerly *Centaurea picris*

Formerly *Rhaponticum*

*repens*

### Plant Type:

Forb

### Flower Color:

Green, pink, and purple

### Habitat:

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

! Toxic (horses); potentially toxic to other livestock

! Introduced

! Noxious

**Russian knapweed** is an introduced, long-lived, creeping perennial. This invasive weed is the most widely distributed of the various knapweed species. It is widespread in northern states, including Colorado, Montana, and Wyoming, and is currently becoming more common in New Mexico and Arizona. Russian knapweed can be distinguished from other knapweeds by its scaly, brown to black, spreading rhizomes and by its unique flowering bract features.

Livestock (including cattle, sheep, and goats) normally will not graze Russian knapweed due to its bitter flavor; however, animals may graze the weed lightly during early growth. The weed is especially toxic to horses and should not be grazed by them. The time of grazing preferred for pastures infested with Russian knapweed is late summer, early fall, or winter. Use grazing to encourage perennial grass growth and competition against Russian knapweed. Reduce grazing pressure in early spring when grasses are first starting to grow, and allow grasses to tiller and produce seed. Utilization of knapweed by livestock should be carefully monitored, and heavy grazing should be avoided.





Patrick Alexander



Patrick Alexander



Patrick Alexander





Matt Lavin

# Kochia

*Bassia scoparia*

## Ch'il nilt'qlí



### Alternative Names:

*Kochia scoparia*

### Plant Type:

Forb

### Flower Color:

Green

### Habitat:

Grasslands

River valleys

Wastelands

### Growing Season:

Warm season

### Special Considerations:

- ! Potentially toxic to livestock
- ! Introduced
- ! Noxious

**Kochia**, native to southern and eastern Russia, was introduced to North America from Europe. Plants growing on dry or infertile sites typically average two feet tall. The leaves are generally green but change to yellow, red, and brown as the plant ages and dies. Dead plants break off at the base and are blown by the wind as tumbleweeds that scatter seeds.

Kochia and common ragweed look very similar from a distance, and the two sometimes occur together. Because kochia is extremely efficient at using water, it can thrive in warm, low rainfall environments. The plants have a wide tolerance of soil types and are often found on saline/alkaline soils. Plants grow on grasslands, pastures, prairies, roadsides, ditch banks, wastelands, floodplains, riparian habitats, and cultivated fields.

Kochia is readily grazed by livestock, but sometimes contains high nitrate levels and can be toxic.

**Kochia**

*Bassia scoparia*

**Ch'il nilt'ólí**



Patrick Alexander



Matt Lavin



Matt Lavin



Russ Kleinman, Bill Norris and the  
Range Vegetation class of autumn  
2009 at WNMU

# Lambsquarters

*Chenopodium album var. missouriense*

Tł'oh dei hałgaí



## Alternative Names:

*Chenopodium missouriense*

Aellen

Lamb's quarters

## Plant Type:

Forb

## Flower Color:

Blue

Purple

## Habitat:

Wastelands

## Growing Season:

Cool season

## Special Considerations:

None

**Lambsquarters** are an erect herb with dentate leaves. The glomerules are densely packed and  $\frac{3}{16}$  to  $\frac{4}{16}$  inch in diameter. There are five sepals which are farinose (dusty in appearance), keeled, and enclose the seed. The pericarp (seed covering) is densely adherent such that the seed tears before the pericarp will separate. This variety of *Chenopodium album* is native to North America and is distinguished by stiff panicles, dense glomerules without interrupted areas, and purplish nodes. *Chenopodium album var. missouriense* is found at roadsides.

*\*Description based on description from Western New Mexico University's Vascular Plants of the Gila Wilderness.*




## Forbs


### Lambsquarters

*Chenopodium album* var. *missouriense*

Tł'oh dei hałgai



Russ Kleinman, Bill Norris, and the Range Vegetation class of autumn 2009 at WNMU



Russ Kleinman, Bill Norris, and the Range Vegetation class of autumn 2009 at WNMU



Russ Kleinman, Bill Norris, and the Range Vegetation class of autumn 2009 at WNMU



Russ Kleinman, Bill Norris, and the Range Vegetation class of autumn 2009 at WNMU



Patrick Alexander

# Larkspur

*Delphinium spp.*

## Tádíííndootł'izh



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Blue

Purple

### Habitat:

Grasslands

River valleys

### Growing Season:

Warm season

### Special Considerations:

! Potentially toxic to livestock

**Larkspur** grows 3 to 8 feet tall, with stems that are somewhat straw-colored and hollow at the base, but a darker bluish color above. It grows from a deep, vertical, woody taproot, beginning growth in late spring, flowering July to August, with seeds maturing from August to September.

Flowers are bluish-purple in color with a spur-like projection, which looks like a duncecap, on the calyx, hence its name.

Flowers grow in dense clusters at the top of the stem. The leaves are hairy and divided into sharp, pointed segments (palmately lobed) that resemble geranium leaves in pre bloom stage. The fruits/seeds are highly poisonous.

Tall larkspur occurs in meadows, thickets, stream banks, around springs, and in moist meadows and open woods. It is most abundant where the snow pack persists. It is usually associated with aspen and spruce groves.

Larkspur provides fair to good forage for sheep and some wildlife. It is palatable to cattle even when other plants are available. Horses seldom eat it. Larkspur is poisonous to cattle until after blossoming. It contains alkaloids, which act on the nervous system. Death may result following paralysis of breathing; bloat is common.

Traditional uses of larkspur included crushing the plants and applying them to hair to control lice and other insects.

*\*Description courtesy of Utah State University's Range Plants of Utah.*





Patrick Alexander

*Delphinium hansenii*



Patrick Alexander

*Delphinium scopulorum*



Patrick Alexander

*Delphinium geraniifolium*



Patrick Alexander

*Delphinium scopulorum*





William M. Ciesla, Forest Health  
Management Intern

# Leafy Spurge

*Euphorbia esula*

## Ch'il abci tsoh



### Alternative Names:

Faitours-grass

Wolf's milk

### Plant Type:

Forb

### Flower Color:

Green

Yellow

### Habitat:

Brushlands

Grasslands

River valleys

Wastelands

### Growing Season:

Warm season

### Special Considerations:

! Noxious

! Poisonous

! Introduced

! Toxic to cattle

! Skin and eye irritation  
in humans

**Spurge** is an herbaceous perennial that is adapted to many soil types and habitats. It typically invades disturbed and undisturbed areas such as pastures, rangelands, abandoned croplands, roadsides, wetlands, woodlands, floodplains, riparian areas, mountain ridges, and prairies. Its milky sap may cause skin and eye irritation in humans and be toxic to cattle if ingested in large quantities.

**Leafy Spurge**  
*Euphorbia esula*  
**Ch'il abci tsoh**



Patrick Alexander



Norman E. Rees, USDA Agricultural  
Research Service



Patrick Alexander



Patrick Alexander



Patrick Alexander

# Loco, Ellis

*Astragalus praelongus* var. *ellisiae*

## Ch'ilaghání'dool ghasí



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Green

Yellow

### Habitat:

Brushlands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

### Growing Season:

Warm Season

### Special Considerations:

! Poisonous

**Ellis loco** has narrowly egg- or football-shaped pods,  $\frac{3}{4}$  to  $1\frac{1}{2}$  inches long, and  $\frac{1}{4}$  to  $\frac{3}{8}$  inch in diameter.

It grows on bluffs, clay knolls, plains, bottomlands, and some forest and woodland areas. Ellis loco can be found on selenium-rich soils and alkaline soils of shale or volcanic parent material at 2,750 to 7,100 feet.

Like some locoweeds, Ellis loco is poisonous/toxic to livestock. Not all locoweeds are poisonous or toxic.

Locoweed species on the Navajo Nation may include:

- *Astragalus amphioxys*
- *Astragalus beathii* ([Group 4](#) Navajo Endangered Species List)
- *Astragalus ceramicus* E. Sheldon var *ceramicus*
- *Astragalus cronquistii* ([Group 3](#) Navajo Endangered Species List)
- *Astragalus cutleri* ([Group 2](#) Navajo Endangered Species List)
- *Astragalus heilii* ([Group 4](#) Navajo Endangered Species List)
- *Astragalus humillimus* ([Group 2](#) Navajo Endangered Species List)
- *Astragalus humistratus*
- *Astragalus lonchocarpus*
- *Astragalus lentiginosus* Douglas ex Hooker var *australis* Barnely
- *Astragalus missouriensis*
- *Astragalus mollisium*
- *Astragalus naturitensis* ([Group 3](#) Navajo Endangered Species List)
- *Astragalus nuttallianus*
- *Astragalus praelongus* Sheldon var *ellisiae*
- *Oxytropis lambertii*



**Loco, Ellis**

*Astragalus praelongus* var. *ellisiae*

**Ch'ilaghání'dool ghasí**




Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Amy Smith Muise

# Loco, Rattleweed

*Astragalus allochrous*

## Dá'ághálii



### Alternative Names:

Rattleweed

Half-moon milkvetch

### Plant Type:

Forb

### Flower Color:

Red

Pink

### Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

Wastelands

### Growing Season:

Warm season

### Special Considerations:

! Toxic to livestock

**Rattleweed loco** is native to Texas, New Mexico, and Arizona. This plant contains swainsonine, which is toxic to livestock.

To recognize it in the field, look for green leaves; small, bluish or purplish flowers; pods that are inflated and bladderlike; and sparse hairs near the base of the plant.

The plants are usually annual, sometimes persisting into a second season, with several stems 4 to 20 inches tall.

Locoweeds species on the Navajo Nation may include:

- *Astragalus amphioxys*
- *Astragalus beathii* ([Group 4](#) Navajo Endangered Species List)
- *Astragalus ceramicus* E. Sheldon var *ceramicus*
- *Astragalus cronquistii* ([Group 3](#) Navajo Endangered Species List)
- *Astragalus cutleri* ([Group 2](#) Navajo Endangered Species List)
- *Astragalus heilii* ([Group 4](#) Navajo Endangered Species List)
- *Astragalus humillimus* ([Group 2](#) Navajo Endangered Species List)
- *Astragalus humistratus*
- *Astragalus lonchocarpus*
- *Astragalus lentiginosus* Douglas ex Hooker var *australis* Barnely
- *Astragalus missouriensis*
- *Astragalus mollisium*
- *Astragalus naturitensis* ([Group 3](#) Navajo Endangered Species List)
- *Astragalus nuttallianus*
- *Astragalus praelongus* Sheldon var *ellisiae*
- *Oxytropis lambertii*



## Forbs

### Loco, Rattleweed

*Astragalus allochrous*

### Dá'ágháíí



Amy Smith Muise




Amy Smith Muise



Amy Smith Muise





Russ Kleinman

# Loco, Wooly

*Astragalus mollissimus*

## Ch'ilaghání



### Alternative Names:

Patterson loco

### Plant Type:

Forb

### Flower Color:

Purple

Red

### Habitat:

Grasslands

Brushlands

### Growing Season:

Warm season

### Special Considerations:

! Poisonous

**Wooly loco** is very silvery in appearance and has relatively large clusters of purple flowers. It is found at lower to middle elevations in fields and dry areas in canyons. It flowers early in the spring and can be found mixed in with ashen milkvetch (*Astragalus tephrodeand*) and rattleweed loco (*Astragalus allochrous*) in disturbed areas.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*

Locoweeds species on the Navajo Nation may include:

- *Astragalus amphioxys*
- *Astragalus beathii* ([Group 4](#) Navajo Endangered Species List)
- *Astragalus ceramicus* E. Sheldon var *ceramicus*
- *Astragalus cronquistii* ([Group 3](#) Navajo Endangered Species List)
- *Astragalus cutleri* ([Group 2](#) Navajo Endangered Species List)
- *Astragalus heilii* ([Group 4](#) Navajo Endangered Species List)
- *Astragalus humillimus* ([Group 2](#) Navajo Endangered Species List)
- *Astragalus humistratus*
- *Astragalus lonchocarpus*
- *Astragalus lentiginosus* Douglas ex Hooker var *australis* Barnely
- *Astragalus missouriensis*
- *Astragalus mollisium*
- *Astragalus naturitensis* ([Group 3](#) Navajo Endangered Species List)
- *Astragalus nuttallianus*
- *Astragalus praelongus* Sheldon var *ellisiae*
- *Oxytropis lambertii*

**Loco, Wooly**  
*Astragalus mollissimus*  
**Ch'ilaghání**



Russ Kleinman



Russ Kleinman



Kelly Allred

# Red Stem Peavine

*Astragalus emoryanus*

## Ch'ilna'át'ó'í



### Alternative Names:

Locoweed

Emory's milkvetch

### Plant Type:

Forb

### Flower Color:

Purple

### Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

### Growing Season:

Warm season

### Special Considerations:

! Toxic to livestock

**Red stem peavine** has stems in loose mats, leaflets less than ½ inch long, and pods that are straw-colored and falling from the stem, splitting open while on the ground.

The plants are annual or winter-annual, seldom persisting into a second year. The stems spread, forming tangled mats, and flower clusters are spreading or drooping. Stems have 1 to 12 flowers with pink-purple petals. Pods are spreading to drooping, narrow, often curved in a half-moon shape, falling from the stem and splitting open from both ends while on the ground. Pods are ½ to 1 inch long and ⅜ to ⅝ inch wide. Flowering March to June, red stem peavine grows in desert grassland, desert scrub, and juniper forest on gravel to clay-loam soils at an elevation of 2,000 to 7,000 feet.

This toxic species is very similar to *Astragalus nuttallianus*, which apparently is not toxic and may provide valuable forage. Not all locoweeds are poisonous or toxic. The main difference between the two is that the pods of *A. nuttallianus* persist on the stem and split open from only the tip while on the stem, whereas the pods of *A. emoryanus* readily fall from the stem and split open from both ends while on the ground. Its principal toxins are nitro-compounds and swainsonine.



**Red Stem Peavine**

*Astragalus emoryanus*

**Ch'ilna'át'ó'í**



*Kelly Allred*



*Kelly Allred*



Patrick Alexander

# Lupine

*Lupinus spp.*

Łį́dą́ą



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Blue

Purple

White

## Habitat:

Brushlands

River valleys

## Growing Season:

Cool season

## Special Considerations:

- ! Poisonous (sheep and horses)

**Lupine** is a complex group of erect to ascending forbs, 10 to 24 inches tall, with one to several stems growing from a branching taproot. Flowers are small and spurred at the base.

Lupine is widespread from valley bottoms to high mountain areas. Because this plant is a legume, nodules on its roots form a symbiotic relationship with bacteria that fix nitrogen. It thus aids in building soil fertility.

Lupine is poor forage for cattle and fair for sheep before the legume fruit develops. It provides fair to good forage for elk and deer. Cattle may be attracted to the legumes and graze them selectively. Lupine is poisonous, especially to sheep and horses. Alkaloids are concentrated in the seeds and occasionally in the young plants. Plants are poisonous either green or dry, but poisoning seldom occurs when other forage is adequate. Poisoning can cause blindness within 10 minutes. The most characteristic symptoms of lupine poisoning are: 1) excitement, leading to running about and butting into other objects; 2) convulsions, with accompanying attacks of dyspnea; and 3) heavy and labored breathing. In sheep, symptoms may not appear for 1 to 24 hours after eating the plant. Poisoning can also cause fetus deformities in pregnant animals. Some lupines can cause crooked calf disease if consumed between the 40th and 70th day of gestation.

A drug has been extracted from lupine for management of cardiac arrhythmias.

*\*Description courtesy of Utah State University's Range Plants of Utah.*

## Forbs

### Lupine *Lupinus spp.* Łį́dą́ą



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander





Patrick Alexander

# Milkweed

*Asclepias spp.*

## Ch'ilabe'é



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Blue

Purple

Green

White

### Habitat:

Grasslands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

! Poisonous to livestock

✓ Good insect forage

**Milkweed** is an important forage plant for monarch butterflies and bees. Numerous milkweeds grow in the southwest United States. Some are native, and others are introduced. The stems and leaves produce a milky sap when broken. The plants grow to heights of about 1 to 3 feet. Their most distinctive feature is the form of the flowers, which are almost orchidlike. After flowering, the plants form a pod filled with white fibers and brown seeds.

Species found in Navajo rangelands include antelopehorn (*Asclepias asperula*) and broadleaf milkweed (*Asclepias latifolia*).

**Forbs**

**Milkweed**

*Asclepias* spp.

**Ch'ilabe'é**




Russ Kleinman

*Asclepias latifolia*



Russ Kleinman

*Asclepias latifolia*



Russ Kleinman

*Asclepias latifolia*



Russ Kleinman

*Asclepias latifolia*



Russ Kleinman

*Asclepias latifolia*



Russ Kleinman

*Asclepias latifolia*



Russ Kleinman

# Mullein

*Verbascum thapsus*

Bijhyiljaa'í



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

## Habitat:

Wastelands

## Growing Season:

Warm season

## Special Considerations:

None

**Mullein** is a tall, unmistakable plant. The leaves are mostly basal, and can reach very large size. They are woolly. The inflorescence is a long crowded spike of yellow flowers. The stamen filaments are very hairy, and the top of the style is flattened. Mullein grows at lower to middle elevations on dry soil and on roadsides. The branching of the multicellular leaf cilia is quite interesting. The flowers mature into two-chambered, stellate-pubescent capsules partially enclosed by leafy bracts. The capsules initially are green, and then turn brown. After the capsules turn brown, they split open from the pointed apex toward the base revealing hundreds of minute, rugose brown seeds.

*\*Based on description at Western New Mexico University's Vascular Plants of the Gila Wilderness.*



## Forbs

### Mullein

*Verbascum thapsus*

Bij̣hyiljaa'í



Russ Kleinman



Amy Smith MuiSe



Russ Kleinman



Russ Kleinman



Patrick Alexander

# Mustard

*Brassica spp.*

## Oostse'



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Purple

White

Yellow

### Habitat:

Grasslands

Ponderosa pine/mixed conifer

### Growing Season:

Cool season

### Special Considerations:

! Potentially toxic to livestock

**Mustard** plants can grow to about three feet in height. Various species from the mustard family can be found in the Navajo Nation. The most common are tansy mustard (*Descurainia pinnata*), which is native to North America, and London rocket (*Sisymbrium irio*), which is introduced. While many members of this family are edible, some, such as tansy mustard, are poisonous to livestock. Most members of the mustard family have yellow or white flowers with four floral parts (petals, stamens, etc.). Many, many species of mustard exist on the Navajo Nation.




## Forbs

## Mustard

*Brassica spp.*

## Oostse'



Patrick Alexander

*Descurainia pinnata*



Patrick Alexander

*Descurainia pinnata*



Patrick Alexander

*Descurainia pinnata*



Patrick Alexander

*Sisymbrium irio*



Patrick Alexander

*Sisymbrium irio*



Patrick Alexander

*Sisymbrium irio*





Patrick Alexander

# Navajo Tea

*Thelesperma megapotamicum*

## Ch'ilgohwéhí'deí



### Alternative Names:

Rayless greenthread

### Plant Type:

Forb

### Flower Color:

Yellow

### Habitat:

Brushlands

Pinon-juniper woodlands

### Growing Season:

Warm season

### Special Considerations:

■ Useful to humans

**Navajo tea** is native to Arizona, New Mexico, Utah, Colorado, and Nevada up through Idaho. It is a member of the aster family. The plants grow to a height of about one foot. The leaves are primarily basal, thin and long, growing in a tuft. Leaves are smooth in texture. Like other asters, the flowers have outer florets (called ray flowers) and a central disk with tiny flowers (called disk flowers).

*Thelesperma subnudum*, a close relative of *Thelesperma megapotamicum*, is also known as Navajo tea.

This plant was traditionally used to make tea and dye. Navajo tea flowers May to September and makes fruit July to September.

# Navajo Tea

*Thelesperma megapotamicum*

Ch'ilgohwéhí'deí



Patrick Alexander



Patrick Alexander



Nick Ashcroft

*Thelesperma megapotamicum*



Nick Ashcroft

*Thelesperma megapotamicum*



Patrick Alexander



Patrick Alexander

# Navajo Tobacco

*Rumex crispus*

## Dziłnát'oh



### Alternative Names:

Curly dock

### Plant Type:

Forb

### Flower Color:

Green

Yellow

Red

Pink

Brown

### Habitat:

Grasslands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

! Introduced

! Toxic to poultry

**Navajo tobacco**, is a perennial plant that grows 1½ to 4 feet high, and reproduces by seed. It generally flowers from June to October.

Young seedlings vary in color from entirely green to red tinged in cooler months. Small flowers occur on the upper portion of the stem, and are yellowish-green at first, but become rosy, then reddish-brown. Flowering is delayed by shoot removal.

Seeds are borne in a large, loose, branching cluster, 1½ to 2 feet long at the top of the stems. Leaves are 3 to 12 inches long (or more including the stalk), with noticeably wavy or curly edges. Smooth, fleshy stems die back each fall, and new ones arise each spring. Stems have swollen nodes and are often reddish in color. The plant has a large, deep taproot, yellowish-orange in color.

Common along roadsides and waste places, it is an aggressive weed in permanent pastures, irrigation ditches, and cultivated fields, especially in association with standing water. Seedlings have a low competitive ability and find it difficult to become established in closed vegetation.

Navajo tobacco prefers deep, moist soil. Seeds and vegetation are toxic to poultry and can cause problems for cattle, sheep and horses. It is an alternate host to many crop diseases.

People have used the plant to treat chronic skin complaints such as psoriasis. It can be useful for constipation. It promotes the flow of bile and has been used to treat jaundice.

*\*Description courtesy of Utah State University's Range Plants of Utah.*





*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*



Patrick Alexander

# Navajo Wild Carrot

*Rumex hymenosepalus*

## Chaasht'ezhiitsoh



### Alternative Names:

Sand dock  
Dock root  
Canyaigre  
Wild rhubarb

### Plant Type:

Forb

### Flower Color:

Green  
Red  
Pink

### Habitat:

Brushlands  
Grasslands  
River valleys

### Growing Season:

Warm season

### Special Considerations:

■ Useful to humans

**Navajo wild carrot** is not a carrot but rather a member of the buckwheat family. It is native to the southwest United States. It is known colloquially as a carrot, perhaps because the root can be used to make a burnt orange dye. Wild carrot has also been used medicinally. The plants grow to a height of three feet and have tall, reddish stems. The leaves are basal, elliptic in shape, and sometimes wavy at the margin. Seeds are reddish-brown. Plants are found on sandy roadsides and fields at lower to middle elevations. The stems are reddish with an interior that is somewhat spongy with airspaces.

**Navajo Wild Carrot**

*Rumex hymenosepalus*

**Chaasht'ezhiitsoh**



Patrick Alexander



Patrick Alexander





Patrick Alexander

# Nightshade

*Solanum spp.*

Ma'iidaá'



## Alternative Names:

Silverleaf nightshade

Western horsenettle

## Plant Type:

Forb

## Flower Color:

Purple

White

## Habitat:

Brushlands

Grasslands

Wastelands

## Growing Season:

Warm season

## Special Considerations:

! Toxic to livestock

! May cause injury

**Nightshades** found on the Navajo rangelands include horsenettle and silverleaf nightshade. Both are native species, but are toxic to livestock as well as to humans despite being related to tomatoes, potatoes, and eggplants. The plants rarely grow to a height of more than three feet. The horsenettle has small thorns, which may cause mechanical injury. Flowers have five petals and are shaped like stars. The petals lie relatively flat, allowing the five yellow stamens to stand out.

**Nightshade**

*Solanum* spp.

**Ma'iidaá'**



Patrick Alexander



Patrick Alexander

*Solanum elaeagnifolium*, silverleaf nightshade



Patrick Alexander



Amy Smith Muise

*Solanum elaeagnifolium*, silverleaf nightshade



Patrick Alexander

# Penstemon

*Penstemon spp.*

## Tsédídeéh



### Alternative Names:

Rocky Mountain penstemon

Beardtongue

Formerly *Penstemon strictus*

### Plant Type:

Forb

### Flower Color:

White

Red

Pink

Purple

### Habitat:

Brushlands

Grasslands

### Growing Season:

None

### Special Considerations:

✓ Good for wildlife

**Penstemon** species are native to the United States, with different species occupying different geographical areas. The plants are common in the West, where they provide forage for wildlife, particularly deer, antelope, and birds. They are considered of minimal grazing value for livestock, but they provide erosion control and diversity in grass and range ecosystems. Penstemons display a variety of colors, but red, purple, and pink are the most common in the Navajo rangelands. The flowers can be distinguished by their consistent shape, which is tubular with a two-lobed upper lip and a three-lobed lower lip. The flowers have five stamens, four of which are fertile (pollen-bearing). The common name “beardtongue” derives from the fuzz or hair often found on the sterile stamen, though not all species of penstemon display this trait.

Species of penstemon on the Navajo Endangered Species List: *Penstemon navajoa*, Navajo penstemon ([Group 3](#), Navajo Endangered Species List)



**Penstemon**

*Penstemon* spp.

**Tsédídeéh**




Patrick Alexander

*Penstemon thurberi*



Patrick Alexander

Hummingbird drinking nectar from *Penstemon barbatus*



Patrick Alexander

*Penstemon fendleri*



Patrick Alexander

*Penstemon fendleri*



Max Licher

# Pepperweed

*Lepidium alyssoides*

## Os si tsóh



### Alternative Names:

Tall whitetop

Giant whiteweed

Perennial pepperweed

Broadleaved pepperweed

Clasping pepperweed

Formerly *Lepidium virginicum*

### Plant Type:

Forb

### Flower Color:

White

### Habitat:

Grasslands

River valleys

### Growing Season:

Warm season

### Special Considerations:

! Invasive

**Pepperweed** is a highly invasive non-native herbaceous perennial. It can invade a wide range of habitats including riparian areas, wetlands, marshes, and floodplains. It adapts readily to natural and disturbed wetlands. As it establishes and expands, the plants create large stands that displace native plants and animals.

In addition to impacting alfalfa and pasture production, it has been reported to adversely affect food quality and nesting habitat for native birds.



**Pepperweed**  
*Lepidium alyssoides*  
**Os si tsón**



Max Licher



Amy Smith Muise



Jeff Brasher



Amy Smith Muise





Russ Kleinman, Andy Anderson, &  
Richard Felger

# Poison Ivy

*Toxicodendron rydbergii*

## K'ishishjiin



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

White

Green

### Habitat:

Grasslands

River valleys

### Growing Season:

Warm season

### Special Considerations:

! Causes rashes

**Poison ivy** can assume different morphologies depending on its location. The viney growth pattern occurs as it grows over rocks. More commonly, it is erect and few-branched up to a few feet tall with trifoliate, shiny, dentate leaves. Large stands with this morphology occur along creeks and rivers in the area. The flowers are inconspicuous and are in small, axillary clusters. The fruit is a yellow drupe.

Most people have an allergic reaction to poison ivy, becoming sensitized by repeated exposure. The reaction is a contact dermatitis, with a severe form consisting of weeping, open blisters. The more severe reactions require treatment by a physician, while milder ones can be treated with over-the-counter medications. The offending agent in the plant is the chemical urushiol. The appearance of the rash may be delayed by a few days after exposure in some people so that it may be difficult to figure out the cause of the reaction.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*

## Forbs

### Poison Ivy

*Toxicodendron rydbergii*

### K'ishishjiin



Russ Kleinman



Russ Kleinman



Russ Kleinman



Russ Kleinman



Patrick Alexander

# Rubberweed, Bitter

*Hymenoxys odorata*

Né'éshjaa'yilkee'é



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

## Habitat:

Brushlands

Grasslands

River valleys

## Growing Season:

None

## Special Considerations:

- ! Toxic to livestock, particularly sheep

**Rubberweed** is native to the southwest United States. It grows in a variety of habitats, including rangelands and riverine areas. It is toxic to livestock, particularly sheep. Cattle will eat it when other green feed is not available. Toxicity increases with maturity of the plant. The plants grow to a height of two feet in a moundlike form. Flowers have a diameter of  $\frac{3}{8}$  to  $\frac{1}{2}$  inch. Pingue is similar to rubberweed.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*





Patrick Alexander



Patrick Alexander



Patrick Alexander

# Sego Lily

*Calochortus nuttallii*

## Gáagiibit'ohchin



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

White

Yellow

### Habitat:

Ponderosa pine/mixed conifer

Brushlands

### Growing Season:

Cool season

### Special Considerations:

None

**Sego lily** is a perennial, native monocot with a single stem and a single, large, terminal flower. This plant is one of the most conspicuous and beautiful early-blooming flowers of the semidesert. They dry up shortly after blossoming. The flower is white with cherry red and yellow markings at the base. Goblet shaped, satiny flowers can be up to three inches across. Fruit is a capsule, which splits open along partitions, with numerous flattened seeds inside. Sego lilies have a few basal, grasslike leaves, and leafy stems. It can be distinguished from death camas (prior to flowering) by the rounded, trough-like cross section of their U-shaped leaves, as opposed to the sharply V-shaped leaves of death camas.

Sego lily occurs on open sagebrush foothills and in valleys, as well as in open ponderosa pine stands at moderate elevations. It thrives on rather dry, sandy soils. The forage value of sego lily is limited, chiefly due to the small amount of herbage and the sparseness of the stands. When eaten, the palatability of the herbage is good for sheep and fair for cattle. Horses do not graze it. The bulbs are eaten, and also gathered and stored by pocket gophers and other rodents.


Edible to humans, the bulb-like roots of sego lily are considered a great delicacy.

*\*Description courtesy of Utah State University's Range Plants of Utah.*


**Sego Lily**  
*Calochortus nuttallii*  
**Gáagiibit'ohchin**



*Patrick Alexander*



*Patrick Alexander*



*Patrick Alexander*





Patrick Alexander

# Senecio

*Senecio spp.*

## Ch'ilbílátahalsoi



### Alternative Names:

Ragwort

Groundsel

### Plant Type:

Forb

### Flower Color:

Yellow

### Habitat:

Brushlands

Grasslands

### Growing Season:

Warm season

### Special Considerations:

- ! Toxic to humans and livestock

**Senecio** plants have simple or branched upright growth with leaves alternate on the stem. Flowers are yellow and appear from April to October.

Plants in the genus *Senecio* produce pyrrolizidine alkaloids, which cause liver damage in livestock and humans when these compounds are ingested. *Senecio* is poisonous to cattle and horses and toxic to humans, such as when plant parts are harvested, either intentionally or unintentionally, for use in herbal teas.

Some *Senecio* species (e.g., *S. vulgaris*) reduce crop yield. Others affect pasture productivity, are poisonous to livestock, or affect native plant communities.

**Senecio**

*Senecio Spp.*

**Ch'ilbílátahalsoi**



Patrick Alexander

*Senecio warnockii*



Patrick Alexander

*Senecio warnockii*



Patrick Alexander

*Senecio wootonii*



Patrick Alexander

*Senecio warnockii*



Rob Routledge, Sault College,  
Bugwood.org

# Sneezeweed

*Helenium autumnale*

Nií'ii'nił



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

## Habitat:

Wastelands

Grasslands

## Growing Season:

Warm season

## Special Considerations:

! Potentially toxic to  
livestock and humans

✓ Butterfly forage

**Sneezeweed** is native to the United States. It is not a significant food source for livestock or large animals, but it does attract butterflies. It can be toxic to livestock (particularly sheep) and humans if consumed in quantity. The plants grow to a height of 4 to 5 feet. Stems are winged. Leaves are hairy. The flower's center is globular and protruding from the petals or ray flowers. Flower diameter is about two inches, and each of the petals have three lobes. Leaves are lance shaped and about six inches long. The plants grow on roadsides and show a preference for moist soils.



**Sneezeweed**  
*Helenium autumnale*  
**Níí'íí'nił**



*Beverly Turner, Jackson Minnesota, Bugwood.org*



*Rob Routledge, Sault College, Bugwood.org*



*Rob Routledge, Sault College, Bugwood.org*



Patrick Alexander

# Spreading Wallflower

*Erysimum repandum*

**Bist'á azéé tsoh**



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

! Noxious

**Spreading wallflower** was introduced to North America from Europe and has become naturalized. It is a member of the mustard family. It grows in disturbed areas and fields. Flowers of this plant are small and yellow with four petals. Multiple flowers grow on each stem in a terminal cluster. The plant later forms pods 1¼ to 3¼ inch long with seeds.

**Spreading Wallflower**

*Erysimum repandum*

**Bist'á azéé tsoh**



*Patrick Alexander*



*Patrick Alexander*





Steve Dewey, Utah State University,  
Bugwood.org

# St. Johnswort

*Hypericum perforatum*

Biłhaazhch'ih



## Alternative Names:

None

## Plant Type:

Forb

## Flower Color:

Yellow

## Habitat:

Grasslands

## Growing Season:

Warm season

## Special Considerations:

- ! Toxic to livestock
- ✓ Pollen source for bees and wasps

**Common St. Johnswort** is an introduced species that can quickly crowd out other plants in disturbed areas, but does not readily colonize undisturbed areas. It is poisonous to livestock, but it provides a pollen source for wasps and bees. St. Johnswort grows in a variety of dry, sunny habitats, particularly grasslands, rangelands, or areas disturbed by fire. Plants grow to a height of three feet with many flowers on each plant. Flowers are yellow, with five petals, and a diameter of  $\frac{3}{4}$  to 1 inch. Leaves are narrow and oblong.

**St. Johnswort**  
*Hypericum perforatum*  
**Biłhaazhch'ih**



Norman E. Rees,  
USDA Agricultural Research Service



William M. Ciesla,  
Forest Health Management Intern



William M. Ciesla,  
Forest Health Management Intern



Eric Coombs, Oregon Department of Agriculture, Bugwood.org



Steve Dewey, Utah State University,  
Bugwood.org

# Sulphur Cinquefoil

*Potentilla recta*

**Azee tsoxíí,  
Chil di tsoxíí**

**Alternative Names:**

Rough-fruited cinquefoil

**Plant Type:**

Forb

**Flower Color:**

Yellow

**Habitat:**

None

**Growing Season:**

Warm season

**Special Considerations:**

! Noxious

**Sulphur cinquefoil** was introduced to North America from Europe. It is considered invasive, and it readily takes to disturbed areas and roadsides. It is similar in appearance to other cinquefoils, growing to about two feet in height. The flowers are approximately  $\frac{1}{2}$  to  $\frac{3}{4}$  inch in diameter with five petals. The leaves are palmate, with five to seven elongated leaflets meeting at a central point. Leaf edges are toothed. The plant grows at elevations of up to 5,000 feet.



**Sulphur Cinquefoil**

*Potentilla recta*

**Azee tsoxíí, Chil di tsoxíí**




Montana Statewide Noxious Weed Awareness



Ohio State Weed Lab , The Ohio State University, Bugwood.org



Ohio State Weed Lab, The Ohio State University, Bugwood.org



Steve Dewey, Utah State University, Bugwood.org



Lesley Ingram, Bugwood.org

# Sunflower

*Helianthus annuus*

## Nidíyílii



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Yellow

### Habitat:

Grasslands

River valleys

### Growing Season:

Warm season

### Special Considerations:

- ✓ Food for wildlife and humans

**The common sunflower** is native to North America, but it has the potential to behave like an invasive, occasionally growing in dense stands. It is grown as a food crop for humans, birds, and other wildlife. It can grow in a range of conditions from roadsides to fields. It grows quite tall, to heights of 3 to 10 feet. The flowerheads are usually 3 to 6 inches in diameter. Leaves are rough in texture with toothed edges. Compared to the giant sunflower, *Helianthus giganteus*, the leaves of this sunflower are more oval shaped.

Other species of sunflower found on the Navajo Nation may include:


*Helianthus anomalus*

*Helianthus maximiliani*

*Helianthus nuttallii*

*Helianthus petiolaris*

**Sunflower**  
*Helianthus annuus*  
**Nidíyílii**



*Richard Floyd, Creative Ideas LLC,  
Bugwood.org*



*William M. Ciesla, Forest Health  
Management Intern*



*Julia Scher,  
Cut Flower Exports of Africa, USDA*





Amy Smith Muise

# Thistle, Musk

*Carduus nutans*

## Azee'okani'whooshí, Azee'okani'deniní



### Alternative Names:

Nodding plumeless thistle

### Plant Type:

Forb

### Flower Color:

Blue

Purple

Green

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ! Invasive
- ! May cause injury
- ! Noxious

**Musk thistle** is a colony-forming weed, growing up to six feet tall, with a spiny stem and long, fleshy taproot. Musk thistle reproduces solely by seed. Musk thistle invades pasture, range, and forestlands. It is often found on roadsides, waste areas, ditch banks, stream banks and in grain fields. It spreads rapidly, forming extremely dense stands that crowd out desirable forage. Musk thistle is shade intolerant. It is adapted to a broad range of soils, but establishment is best on bare soil. Musk thistle is tolerant of saline and acidic soils and can be a noxious weed. Chemical control is effective. An introduced biological control agent, the musk thistle weevil, feeds on the seeds and can limit the spread of this plant, but will not eliminate it. Musk thistle is not palatable to livestock because of its long sharp spines. Musk thistle does provide a source of nectar for high quality honey.

*\*Description based on description at Utah State University's Range Plants of Utah.*

## Forbs

### Thistle, Musk

*Carduus nutans*


Azee'okani'whooshí, Azee'okani'denini



Amy Smith Muise



Amy Smith Muise



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander

# Thistle, Canada

*Cirsium arvense*

## Azéé hakaní yázhí



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

Blue

Purple

### Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

### Growing Season:

Warm season

### Special Considerations:

! Noxious

! Invasive

**Canada thistle** is a non-native, invasive, perennial plant that can be very difficult to kill. Its roots spread aggressively and form dense colonies on roadsides, crop fields, pastures, and rangelands, generally on disturbed ground. It emerges from its root system (or from seed) in the spring and forms rosettes. Flowers occur in groups of one to five, varying in color from white to purple. Individual flowers are about half an inch in diameter. Leaves are bright green and shiny with a hairy undersurface. Canada thistle prefers moist soil but can be found in a variety of soil types, at elevations up to 12,000 feet. Seeds survive in the soil up to 20 years, so it is wise to treat new infestations (with grazing, mowing and/or chemical control) before plants set seed. Pulling up or tilling Canada thistle spreads root fragments that can form new plants. Prevention, by maintaining healthy pastures, riparian areas, and rangelands, is the best control.

Cattle, goats, and sheep will graze on Canada thistle when plants are young.

Species of thistle on the Navajo Endangered Species List: *Cirsium rydbergii*, Rydberg's thistle ([Group 4](#) Navajo Endangered Species List)




**Thistle, Canada**  
*Cirsium arvense*  
**Azéé hakaní yázhí**



Udo Schmidt



Patrick Alexander



Patrick Alexander



Patrick Alexander



Superior National Forest



Amy Smith Muiise

# Thistle, Scotch

*Onopordum acanthium*

**Zéé hókaniít béeí,  
Whosh Waa'**



## Alternative Names:

Scotch cottonthistle

## Plant Type:

Forb

## Flower Color:

Purple

## Habitat:

Piñon-juniper woodland

## Growing Season:

Warm season

## Special Considerations:

! Noxious

**Scotch thistle** is a noxious, biennial weed that grows up to 12 feet tall. It has spiny wings along the stem. The leaves have a grayish appearance due to dense hairs on the surface. Flowers are reddish to violet. Scotch thistle is often associated with waste places and roadways, as well as rivers, streams, canals, or other waterways,

It can also be abundant in dry pastures, fields, and rangeland, and thrives in light, well-drained, and sandy or stony soils.



# Thistle, Scotch

*Onopordum acanthium*

Zéé hókaníít béí, Whosh Waa'



Amy Smith Muise



Leslie J. Mehrhoff, University of Connecticut, Bugwood.org



Lesley Ingram, Bugwood.org



Leslie J. Mehrhoff, University of Connecticut, Bugwood.org



Leslie J. Mehrhoff, University of Connecticut, Bugwood.org



Leslie J. Mehrhoff, University of Connecticut, Bugwood.org





Nick Ashcroft

# Tumbleweed

*Salsola tragus*

## Ch'ildeeníní



### Alternative Names:

Russian thistle  
Prickly Russian thistle  
Slender thistle  
Formerly *Salsola kali*

### Plant Type:

Forb

### Flower Color:

Blue  
Purple  
White

### Habitat:

Brushlands  
Grasslands  
Wastelands

### Growing Season:

Warm season

### Special Considerations:

- ! Sign of unhealthy rangeland
- ! May cause injury

**Russian thistle** (*Salsola tragus*), barbwire Russian thistle (*Salsola paulsenii*), and spineless Russian thistle (*Salsola collinea*) are common annual weeds found in the southwestern United States. These plants are problematic in a wide range of areas, including agricultural, roadside, pasture, urban, and residential areas. They increase fire hazard and thrive in areas of high disturbance. Prevention or reduction of disturbance is critical in preventing establishment of these weeds!

Russian thistle grows from 1 to 3 feet tall. Its branched and rounded shape means it can be up to five feet wide. Found on nearly all soil types, it is salt resistant and grows on alkali soils. Russian thistle provides fair forage for cattle and sheep in the early spring, when leaves are soft and succulent, becoming worthless with maturity because of the sharp-pointed leaves. If it turns black at maturity and is softened by moisture it is good winter feed. Livestock can suffer injury from the leaves, which may accumulate nitrates, and may contain oxalates. It is an indicator of overused and unhealthy rangeland. It does not compete well with other plant species. When the plant is mature, it breaks off at the base and blows away, hence the name "tumbleweed." This is how the seed is spread.

Young shoots of Russian thistle can be used as a potherb and seeds can be ground into meal. Russian thistle is sometimes harvested for hay and silage.


**Tumbleweed**

*Salsola tragus*

**Ch'ildeeníní**



Patrick Alexander



Nick Ashcroft

*Salsola nigra*



Patrick Alexander



Patrick Alexander



Mary Ellen (Mel) Harte, Bugwood.org

# Water Hemlock

*Cicuta douglasii*

## Shashdáá'



### Alternative Names:

None

### Plant Type:

Forb

### Flower Color:

White

### Habitat:

River valleys

### Growing Season:

Cool season

### Special Considerations:

- ! Poisonous,  
especially its roots

**Water hemlock** is an erect forb, growing 3 to 7 feet tall, usually swollen at the base. The stem has cross partitions at the base. Roots also have partitions inside, and a musky odor. It begins growth in the spring, and flowers in late spring and early summer. Reproduction is by seed and tuber branching. Flowers are small, white, and grouped in umbrella-shaped clusters.

The fruit is kidney-shaped, tea-colored, with corky ridges. Leaves are alternate, narrow to broadly lance-shaped, with toothed margins. A distinguishing characteristic is that the veins in the leaf run to the notches between the teeth.

Water hemlock is a wetland plant and is especially common in pastures or untilled areas. It is found along mountain streams and valley ditch banks.

It occurs on wet, fertile soils at the water's edge. It is most common in deep loams, clay loams, or clays. Associated species include Nebraska sedge, redtop, bulrush, and timothy.

Water hemlock contains highly poisonous alkaloids. The roots are the most toxic part of the plant. When roots become exposed and are eaten by livestock, death occurs in 1 to 12 hours. The leaves and stems lose most of their toxicity as they mature. Sheep do not seem to be as affected as cattle. Symptoms of poisoning include muscle twitch, rapid pulse, rapid breathing, tremors, convulsions, excessive salivation or frothing at the mouth, and dilation of the pupils.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



## Forbs

### Water Hemlock

*Cicuta douglasii*

Shashdáá'



Mary Ellen (Mel) Harte, Bugwood.org



Mary Ellen (Mel) Harte, Bugwood.org



Mary Ellen (Mel) Harte, Bugwood.org



Mary Ellen (Mel) Harte, Bugwood.org



Patrick Alexander

# Western Yarrow

*Achillea lanulosa*

## Tééhch'íhóózhood



### Alternative Names:

Wooly yarrow

### Plant Type:

Forb

### Flower Color:

White

### Habitat:

Brushlands

### Growing Season:

Warm season

### Special Considerations:

- ✓ Wildlife and livestock forage

**Western yarrow** is a native, herbaceous perennial in the aster family. It is a very common wildflower that grows erect from creeping rootstocks, to a height of 10 to 36 inches.

Western yarrow is a food source for bighorn sheep, pronghorn antelope, and deer. Sage grouse (especially chicks) and other upland birds rely heavily on the foliage of western yarrow as a food source. Domestic sheep and goats derive a fair amount of forage value from western yarrow, while cattle and horses mostly graze the flowerhead. The volatile oils, alkaloids, and glycosides are considered toxic, but the plant is seldom overgrazed by foraging animals. Western yarrow has been used as a tea to cure stomach ailments, a poultice on infected wounds, and as a mosquito repellent.



Patrick Alexander



Patrick Alexander

# Whitetop

*Lepidium draba*

Chil bi la t'ahłigaí



## Alternative Names:

Hoary cress

Formerly *Cardaria draba*

## Plant Type:

Forb

## Flower Color:

White

## Habitat:

Grasslands

River valleys

## Growing Season:

Warm season

## Special Considerations:

! Invasive

**Whitetop** was introduced to North America from Europe and Asia, and is now a widespread invasive. It grows in a wide variety of habitats and can infiltrate grazing lands. It can grow to be two feet tall and has grayish-green leaves and clustered white flowers with four petals.



## Forbs

### Whitetop

*Lepidium draba*

Chil bi la t'ahłigai



Patrick Alexander



Patrick Alexander



Russ Kleinman



Russ Kleinman



Patrick Alexander

# Woolly Indian Wheat

*Plantago patagonica*

Azee'haatl'inii



## Alternative Names:

None

## Plant Type:

Forb

## Habitat:

Brushlands

Grasslands

## Growing Season:

Warm season

## Special Considerations:

None

**Woolly Indian wheat** is the only member of this family which is found away from disturbed areas and in the wilderness. It is fuzzy and gray in appearance, as opposed to the bright green appearance of *Plantago major* and *Plantago lanceolata*, both of which require considerably more water to survive. *Plantago patagonica* can achieve the same size as the other two species, but more commonly it is quite a bit smaller.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*



**Wooly Indian Wheat**

*Plantago patagonica*

**Azee'haatl'inii**



Patrick Alexander



Patrick Alexander



Patrick Alexander



Patrick Alexander





H. Zell [GFDL or CC BY-SA 3.0], from Wikimedia Commons

# Wormsweed

*Dysphania ambrosioides*

## Tsé'ázhííh



### Alternative Names:

Epazote

### Plant Type:

Forb

### Flower Color:

Green

### Habitat:

Wastelands

### Growing Season:

Warm season

### Special Considerations:

✓ Edible

**Wormsweed** is not native, but has become naturalized. It is commonly known as epazote, and it is used in Mexican, Central American, and South American cuisine. It grows in disturbed areas. The plant can grow to be four feet tall. The leaves vary in shape, from ovate to lanceolate, and can have small to deep serrations on the leaf edges. The small green flowers occur in clusters, and later form red-brown seeds.



Forest & Kim Starr [CC BY 3.0], via Wikimedia Commons



Forest & Kim Starr [CC BY 3.0], via Wikimedia Commons



Matt Lavin

# Cholla

*Cylindropuntia spp.*

## Hoshdítsáhii



### Alternative Names:

Walking stick cholla

Cane cholla

### Plant Type:

Cactus

### Flower Color:

Purple

### Habitat:

Grasslands

Wastelands

### Growing Season:

Warm season

### Special Considerations:

! Causes injury

**Cholla** is a tree-like plant growing to eight feet tall with cylindrical joints and branching stems with numerous one-inch spines. Flowers mature into yellow fruit which drops to the ground when ripe, producing new plants. Cholla also can reproduce from terminal joints that fall to the ground and root when moisture is available.

Cholla often becomes a problem on rangeland when the grass cover is depleted by drought and over utilization. After establishment, the cactus encroaches upon valuable range until dense thickets develop. These thickets hinder livestock operations and compete with needed forage for moisture and nutrients. Cows may develop a cholla-eating habit, especially when pastured among thick stands.

Cholla species found on the Navajo Nation may include:

*Cylindropuntia versicolor*

*Cylindropuntia imbricata*

*Cylindropuntia whipplei*



**Cholla**

*Cholla spp.*

**Hoshdítsáhíi**



Amy Smith MuiSe



Matt Lavin

Cholla browsed by cattle.



Ken Gishi



Amy Smith MuiSe





Patrick Alexander

# Prickly Pear

*Opuntia spp.*

## Hoshniteelí



### Alternative Names:

Plains prickly pear

*Opuntia phaeacantha*

### Plant Type:

Cactus

### Flower Color:

Purple

Pink

Yellow

### Habitat:

Brushlands

Grasslands

Piñon-juniper woodland

Ponderosa pine/mixed conifer

Wastelands

### Growing Season:

Cool season

### Special Considerations:

! Potentially invasive

✓ Forage for wildlife

**Prickly pears** produce familiar pink-red fruits and pads, both very thorny. These plants provide shelter for wildlife, but may become invasive on rangelands because livestock do not find them highly palatable.

The plants are mainly 1 to 2 feet tall and can spread into wide colonies by layering and sprouting from fallen segments. The flowers are large, with numerous yellow, and sometimes pink or violet petals. Prickly pear may displace desirable vegetation if not properly managed. It is often a problem on overgrazed rangeland. The spines of prickly pear make it and other desirable vegetation growing underneath unavailable to livestock. However, it is palatable to livestock once the spines are removed.

Species that grow in or near Navajo rangelands include:

*Opuntia polyacantha*, the plains prickly pear, which provides protection and cover for small wildlife.

*Opuntia chlorotica*, pancake prickly pear or clockface prickly pear, which has a trunk like a tree.

*Opuntia engelmannii*, Engelmann prickly pear, which has larger paddles than other species.

*Opuntia macrocentra*, purple prickly pear.

*Opuntia phaeacantha*, brown-spined prickly pear, which grows in a sprawling formation along the ground.

*Opuntia pottsii*, Pott's prickly pear, whose fruits appear elongated compared to other species.

Prickly Pear

*Opuntia* spp.

Hoshniteelí



Patrick Alexander

*O. phaeacantha* and *O. polyacantha* are very common on the Nation



Nick Ashcroft

*Opuntia macrorhiza*



Patrick Alexander



Nick Ashcroft

*Opuntia macrorhiza*





Nick Ashcroft

# Yucca

*Yucca spp.*

## Tsá'ászeh



### Alternative Names:

Banana yucca (one species)

Narrowleaf yucca  
(another species)

*Yucca angustissima*

*Yucca baccata*

### Plant Type:

Cactus

### Flower Color:

White

Yellow

### Habitat:

Brushlands

Wastelands

### Growing Season:

Cool season

### Special Considerations:

✓ Edible stalks, buds, flowers

✓ Edible fruit (some species)

**Yuccas** are deep rooted and long lived, with individual plants living hundreds of years. Leaves stand out protectively and are armed with sharp points. Most have loose, thread-like fibers that curl from their edges.

Flowers produced in May and June are highly palatable and used by livestock and wildlife.

Yucca has diverse uses. Stalks, buds, flowers, and some fruits have served as food. Roots can be used for soap and as a laxative, and leaf fibers can be used as cordage, weaving material, and to make sandals.

The fruits of banana yucca are fleshy and succulent, roughly like short, fat, green bananas. (Most other yuccas have dry, hard fruits.) Banana yucca fruits are traditional foods. They are prepared by roasting or baking, stripping out the seeds, pounding the remaining flesh into a pulp, forming the pulp into flat cakes, and sun-drying for later use. The resulting product is nutritious, sweet, and delicious. The fruits are often picked before maturity and ripened off the plant to keep wildlife from eating them.



**Yucca**  
*Yucca spp.*  
**Tsá'ászeh**



Patrick Alexander

*Yucca baccata*



Patrick Alexander



Patrick Alexander

*Yucca baileyi*



Patrick Alexander



Nick Ashcroft

*Yucca baileyi*



Russ Kleinman, Bill Norris & Mark Donnell

# Alder

*Alnus spp.*

## K'ish



### Alternative Names:

Arizona alder

### Plant Type:

Tree

### Flower Color:

Green

Red

Pink

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

None

**Alder** is usually a streambank tree, although it can also grow as a large shrub. The leaves are simple, doubly serrate, and heavy-veined. The male flowers are structured in catkins and the female flowers are surrounded by bracts which harden into a woody remnant (called by some an ament, and by others a strobilus) which resembles a small cone. The young branches can be longitudinally ridged.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*



**Alder**  
*Alnus spp.*  
**K'ish**



*Russ Kleinman*



*Russ Kleinman*



*Russ Kleinman*



*Russ Kleinman, Bill Norris, Scott Zager,  
Maggie Knox, Leith Young, &  
Bonnie Arfsten*





Amy Smith Muise

# Colorado Piñon

*Pinus edulis*

## Chá'ot



### Alternative Names:

Piñon pine  
Common pinyon  
Twoneedle pinyon  
Colorado pinyon

### Plant Type:

Tree

### Flower Color:

Green

### Habitat:

Piñon-juniper woodland

### Growing Season:

None

### Special Considerations:

✓ Provides wildlife forage

**Colorado piñon** often grows as a low, bushy tree with an irregularly rounded, spreading crown. The trunk is generally short and crooked, with several large, crooked branches. It may grow to 40 inches in diameter. Height is typically 26 to 56 feet. It is a slow-growing, long-lived tree. It can survive more than 500 years and may reach 800 to 1,000 years of age.

Piñon nuts are a preferred food for turkeys, piñon jays, woodrats, bears, and other wildlife, and they are a common food for deer, particularly during harsh winters with deep snows. Piñon ranks first among the native nut trees that are not also cultivated. The nuts are commonly sold and eaten after roasting in the shell, but small quantities are sold raw. They were once a staple food.

**Colorado Piñon**

*Pinus edulis*

**Chá'ot**



*Patrick Alexander*



*Patrick Alexander*



*Amy Smith Muise*



*Patrick Alexander*



Patrick Alexander

# Coyote Willow

*Salix exigua*

K'ei'łibáhí



## Alternative Names:

Sandbar willow

Narrowleaf willow

## Plant Type:

Tree

## Flower Color:

Yellow

White

## Habitat:

River valleys

## Growing Season:

Cool season

## Special Considerations:

- ✓ Provides wildlife and livestock forage and habitat
- ✓ Provides erosion control

**Coyote willow** usually forms a thicket with its long, slender stems. It seldom exceeds 15 feet in height, but has been known to reach heights of 26 feet. It reproduces by seed and rhizomes (forming clones). Regeneration may also occur through broken pieces of stems and roots that sprout after being transported and deposited by floodwaters.

It has caterpillar-like catkins, which are long spikes containing many small flowers. The wood is light, soft, and weak, susceptible to decay. Coyote willow occupies banks of major rivers and smaller streams, lakes and ponds, marshy areas, alluvial terraces, and ditches, at elevations from 2,700 to 8,500 feet. It characteristically forms zones immediately adjacent to the water's edge. Coyote willow may also occur on moist, well-drained benches and bottomlands. It is a pioneer species, one of the first to colonize gravelly and sandy flood deposits. It is also well adapted to continued survival on sites that are regularly flooded. It is an important food source for many wildlife species.

Stands of coyote willow should be maintained because they help stabilize streambanks and protect them from erosion. Once degradation occurs, streambanks may erode rapidly. All willows produce salicin, which chemically is closely related to acetylsalicylic acid, commonly known as aspirin. Various preparations from willows have been used to treat toothache, stomachache, diarrhea, dysentery, and dandruff. Stems have been made into baskets and bows, and the bark into fabric.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



**Coyote Willow**

*Salix exigua*

**K'ei'tibáhi**



Patrick Alexander



Patrick Alexander



Brian Muise

# Fremont Cottonwood

*Populus fremontii*

## T'iis



### Alternative Names:

Rio Grande cottonwood

Valley cottonwood

### Plant Type:

Tree

### Flower Color:

Green

Yellow

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

None

**Fremont cottonwood** is a native tree growing in riparian areas near streams, rivers, and wetlands. Its height ranges from 40 to 115 feet, and trunk diameter ranges from 1 to 4 feet. The bark is smooth in younger trees, becoming deeply furrowed with whitish, cracked bark with age. The trees bloom from March to April. The wind-dispersed fruit is attached to silky hairs, which gives the appearance of patches of cotton hanging from the limbs, thus the name “cottonwood.”

Cottonwood leaves have been used for soothing poultices. Its inner bark provides a source of Vitamin C. Cottonwood often grow from multiple large, spreading trunks.

## Fremont Cottonwood

*Populus fremontii*

T'iis



Amy Smith Muiise



Patrick Alexander



Patrick Alexander



Amy Smith Muiise





Patrick Alexander

# Gambel Oak

*Quercus gambelii*

## Che ch'il



### Alternative Names:

None

### Plant Type:

Tree

### Flower Color:

Green

Yellow

Brown

### Habitat:

Ponderosa pine/mixed conifer

### Growing Season:

Warm season

### Special Considerations:

- ✓ Acorns are edible
- ! Poisonous to cattle at high dosages

**Gambel oak** is ecologically important, providing food and shelter for many wildlife species. It occurs as clones of shrubs in dense patches 3 to 20 feet tall, often with a central thicket rising above the others. Widely spaced trees grow up to 75.5 feet tall. Bark of gambel oak is rough and furrowed.

Gambel oak may contribute up to 50% of cattle diet without cattle showing any ill effects, with poisoning occurring at higher percentages, and death often resulting when more than 75% of the diet is gambel oak. Freezing enhances toxic properties of gambel oak browse; young foliage turned black by freezing is extremely toxic.



Patrick Alexander



Patrick Alexander

# Juniper

*Juniperus spp.*

## Gad



### Alternative Names:

None

### Plant Type:

Tree

### Flower Color:

Green

### Habitat:

Piñon-juniper woodland

### Growing Season:

Warm season

### Special Considerations:

None

**Juniper** of several species grow on the Navajo Nation. They are small, native, evergreen trees growing up to 40 feet, often with an irregular crown. They occur in mixed or pure stands of open scrub woodland at elevations of 5,000 to 7,500 feet, often on dry, rocky ridges but also in canyons or on middle-elevation foothills.

Juniper does well in cultivation, is adapted to a wide range of soils and moisture conditions, and is winter hardy, slow growing, and very long lived.

Piñon-juniper woodlands provide good habitat for mule deer, bighorn sheep, bison, wild horses, pronghorns, coyotes, bobcats, badgers, porcupines, rabbits, mice, voles, woodrats, squirrels, and numerous birds. Juniper foliage and berries provide food for many species of birds and mammals, including humans.

Across the West, junipers have expanded their historical range, especially into sagebrush-grass communities below areas of traditional piñon-juniper. Overgrazing, fire suppression, and climatic change have been identified as potential causes of juniper invasion. In the absence of fire or other disturbances, trees eventually dominate a site and crowd out herbaceous and shrub species.

Species found on the Navajo Nation include:

- *Juniperus scopularum* (Rocky Mountain juniper)
- *Juniperus monosperma* (one-seed juniper)
- *Juniperus osterosperma* (Utah juniper).



**Juniper**  
*Juniperus spp.*  
**Gad**



Patrick Alexander



Patrick Alexander



Patrick Alexander





Patrick Alexander

# Mountain Mahogany

*Cercocarpus montanus*

Tsé'ásdaazii



## Alternative Names:

Birchleaf mountain mahogany

## Plant Type:

Tree

## Flower Color:

White

Yellow

## Habitat:

Piñon-juniper woodland

## Growing Season:

Warm season

## Special Considerations:

- ✓ Provides forage for wildlife and livestock
- ✓ Provides wildlife habitat

**Mountain mahogany** is a shrub or small tree that grows from 2 to 20 feet tall, with upright or spreading branches. It flowers from May to June, and its fruits mature from July to September. It reproduces from seed and by root sprouts. Seed production is sporadic—the plants go as long as 10 years with no seed production. Mountain mahogany occurs on rocky bluffs, mountainsides, rimrock, breaks, and in canyons and open woodlands. It is common in swales where snow persists during the winter. It grows at elevations between 5,000 and 7,000 feet. It is heat and drought tolerant. It is also somewhat shade tolerant, but grows better without a forest canopy.

It is good to excellent forage for cattle, sheep, and goats. It is a good source of cover for livestock, big game, and many small mammals and birds. It is extremely valuable as winter browse for deer and bighorn sheep.

The twigs are palatable all year long, and are grazed heavily. However, the leaves may contain cyanogenic glycoside, which may cause hydrocyanic poisoning.

Wood from mountain mahogany has been used to make tools and war clubs, and its bark to make a reddish-brown dye for leather.

## Mountain Mahogany

*Cercocarpus montanus*

Tsé'ásdaazii



Patrick Alexander



Patrick Alexander



Patrick Alexander

# Ponderosa Pine

*Pinus ponderosa*

Índíshchíí'



## Alternative Names:

Big heavy  
Black jack  
Bull pine  
Ponderosa white

## Plant Type:

Tree

## Flower Color:

Pink  
Yellow

## Habitat:

Ponderosa pine/mixed conifer

## Growing Season:

Warm season

## Special Considerations:

- ✓ Good wildlife forage
- ! Potential danger to cattle

**Ponderosa pine** is a large, long-lived, native forest tree common throughout the western United States. The height of mature trees ranges from 55 to 90 feet. Ponderosa pine seeds are consumed by a great many birds and small mammals such as mice, chipmunks, and tree squirrels. Snags in the mature pine forest provide a large number of species with nesting and roosting sites. Big game, such as deer and elk, also use the pine forests for food and shelter. Ponderosa pines are fire-adapted and may grow in an open, park-like structure.

When the needles are eaten during gestation, they can cause abortion in cattle.



## Ponderosa Pine

*Pinus ponderosa*

Índíshchíí'



Patrick Alexander




Patrick Alexander



Patrick Alexander



Patrick Alexander



Russ Kleinman, Richard Felger,  
& Carey Anne Laffert

# Quaking Aspen

*Populus tremuloides*

## Ts'iisbéíí



### Alternative Names:

None

### Plant Type:

Tree

### Flower Color:

Red

Pink

White

### Habitat:

Ponderosa pine/mixed conifer

River valleys

### Growing Season:

Cool season

### Special Considerations:

- ✓ Good forage for sheep
- ✓ Provides watershed protection

**Quaking aspen** is an attractive deciduous tree, growing up to 40 feet tall. The trunk is generally long and slender, but can be up to nearly three feet in diameter. The leaves change from green to bright yellow or yellowish-orange in the fall. Quaking aspen grows on moist uplands, dry mountainsides, high plateaus, mesas, avalanche chutes, talus, parklands, gentle slopes near valley bottoms, alluvial terraces, and along watercourses. It is most common at elevations between 6,000 and 10,000 feet. Aspens seldom reproduce from seed. Most reproduction of aspen is by basal sprouting or root sprouting, many trees in a grove being connected together by a common root system in what are referred to as "clones." Quaking aspen grows on a range of soils: shallow, rocky soil; deep, loamy sands; and heavy clays. Aspen is good to excellent forage for sheep, and fair for cattle. The twigs, bark, and buds are browsed by wildlife, and birds eat the seeds. Grazing sprouts, especially by cattle and elk, is a growing concern in the maintenance of aspen stands.

Wild and domestic ungulates use quaking aspen for summer shade. Seral quaking aspen communities provide excellent hiding cover for moose, elk, and deer. Deer use quaking aspen stands for fawning grounds. Well-stocked quaking aspen stands provide excellent watershed protection. Aspen bark contains salicin (similar to the active ingredient in aspirin) and traditionally was used internally as an expectorant and externally as a counterirritant.

*\*Description courtesy of Utah State University's Range Plants of Utah.*



## Quaking Aspen

*Populus tremuloides*

Ts'iisbéíí



Russ Kleinman, Bill Norris, &  
Mark Donnell



Patrick Alexander



Patrick Alexander



Russ Kleinman



Russ Kleinman





Paul Wray, Iowa State University,  
Bugwood.org

# Red Plum

*Prunus americana*

## Didzétsoh dík'ózhígíí



### Alternative Names:

American plum

### Plant Type:

Tree

### Flower Color:

White

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

✓ Edible fruit

**Red plum** is a shrub or small tree that can form small groves in moist middle elevation creekbeds and riparian areas. The small umbels of flowers bloom in the spring before the plant leafs out. The bark is gray and obviously peeling away on larger trunks. The trunks can be up to 8¼ inches in diameter. The flowers are white, and the petals are about ½ inch long. The leaves are serrate and mostly glabrous.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*

**Red Plum**

*Prunus americana*

**Didzétsoh dík'ózhígíí**



David Stephens, Bugwood.org



T. Davis Sydnor, The Ohio State University, Bugwood.org



T. Davis Sydnor, The Ohio State University, Bugwood.org



William M. Ciesla, Forest Health Management Intern



Nick Ashcroft

# Russian Olive

*Elaeagnus angustifolia*

## Tsin łibáhá



### Alternative Names:

Silverberry

Oleaster

Wild olive

Persian olive

### Plant Type:

Tree

### Flower Color:

Yellow

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

! Noxious

**Russian olive** is a shrub or small tree, usually 12 to 45 feet tall. It can grow up to six feet per year. It forms a dense, rounded crown. Near the ground, its branches spread from 10 to 20 feet. Unpruned trees have five or six main stems starting near the ground. The twigs are flexible, coated with a gray, scaly pubescence, and have a thorn at the end. Bark is reddish-brown and thin, with shallow fissures. It exfoliates into long strips.

Russian olive is commonly found growing along floodplains, riverbanks, stream courses, marshes, and irrigation ditches at elevations from 4,500 to 6,000 feet. It is tolerant of considerable amounts of salinity or alkalinity, and can survive considerable droughts. In addition to reproducing from seed, it sprouts from the root crown and sends up suckers. It thrives under a wide range of soil textures, from sand to heavy clay, and withstands flooding and silting. It grows best in deep sandy or loamy soils with only slight salt and alkali content.

Russian olive has no forage value for livestock or big game, although both browse the foliage at times. In general, Russian olive-dominated communities provide inferior wildlife habitat to that of native riparian vegetation. Russian olive rapidly colonizes lowland fields, often chokes irrigation ditches, and damages tires and equipment. Once established, Russian olive is difficult to control and nearly impossible to eradicate. The most effective combination of control efforts is cutting trees, followed by either spraying or burning the stumps.



**Russian Olive**  
*Elaeagnus angustifolia*  
**Tsin łibáhá**



Amy Smith Muise



Amy Smith Muise



Z. Rauch

# Saltcedar

*Tamarix spp.*

## K'eilichii'its'óóz



### Alternative Names:

None

### Plant Type:

Tree

### Flower Color:

Red

White

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

✓ Forage for bees

! Invasive

**Saltcedar** is a perennial, introduced, deciduous tree. It lives 50 to 100 years, and grows 6 to 26 feet tall. Saltcedar has a deep taproot and extensive laterals. It reproduces from seeds, as well as root sprouts. The flowers are white to pink, each 1/16-inch long. They grow in narrow drooping clusters 1 to 2 inches long.

Saltcedar commonly occurs along floodplains, riverbanks, stream courses, salt flats, marshes, and irrigation ditches. It often forms pure thickets that extend for miles. Once established, saltcedar can tolerate both drought and flooding. By shedding its leaves and halting growth, it can withstand lengthy drought periods. It can also survive inundation by water. Saltcedar has a deep taproot that can penetrate the soil to depths of 30 feet or more. It also has extensive laterals. Its dense roots and rhizomes spread out and slow river flow, which increases deposition, and can cause rechanneling of streams and rivers. Leaf drop increases the salinity of the surface soil, causing a white color, thus the name saltcedar.

Saltcedar provides nesting for birds, and is an important pollen source for honeybees. It is also used by blacktailed jackrabbits as a major food source. It is relatively unpalatable to most classes of livestock and wildlife.

Medicinally, saltcedar species have been used to treat stomach troubles, diarrhea, rheumatism, rickets, and as an astringent and antiseptic.

There are two species of saltcedar in the region: *Tamarix ramosissima* and *Tamarix parviflora*, extremely similar in appearance, and both noxious.



**Saltcedar**

*Tamarix spp.*

**K'eiłichii'its'óóz**



Russ Kleinman & Richard Felger  
*Tamarix ramosissima*



Gerald Moore



Amy Smith Muese  
*Tamarix ramosissima*



Renee Benally  
*Tamarix ramosissima*



Z. Rauch  
*Tamarix ramosissima*





Russ Kleinman

# Siberian Elm

*Ulmus pumila*

## Naasts'ósí bit'iis



### Alternative Names:

None

### Plant Type:

Tree

### Flower Color:

Green

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

! Noxious

**Siberian elm** is an introduced, fast-growing tree, from 50 to 70 feet in height. Its leaves are alternate, oblong in shape, 1 to 3 inches long, and usually have serrate (saw-toothed) margins. The flowers are greenish and clustered, with short pedicels. They appear with or before the leaves, from March through April. The bark is a light gray-brown with irregular furrows and is often streaked with stains caused by bacterial wetwood. The fruit, a samara, ripens from April to May, and consists of a dry, compressed nutlet surrounded by a thin, membranous wing.

**Siberian Elm**

*Ulmus pumila*

**Naasts'ósi bit'iis**



Russ Kleinman



Russ Kleinman



Russ Kleinman



Russ Kleinman



Russ Kleinman



Matt Lavin

# Subalpine Fir

*Abies lasiocarpa subsp. Arizonica*

## Ch'oh deeníní



### Alternative Names:

None

### Plant Type:

Tree

### Flower Color:

None

### Habitat:

Ponderosa pine/mixed conifer

### Growing Season:

Year-round

### Special Considerations:

None

**Subalpine fir** is a native, evergreen growing to 65 feet tall. It has a sharp, spire-like crown, the upper several feet of which are often less than 1 foot in diameter. The plants often are reduced to a prostrate shrub on exposed sites near the timberline.

There are several subspecies of subalpine fir, but the one most likely to be seen on the Navajo Nation is *Abies lasiocarpa subsp. Arizonica*. General characteristics of the species are smooth, grayish-white bark with resin blisters. In subspecies *Arizonica*, the trunk is more likely to have softer, corky bark. The bark becomes furrowed as the tree reaches one foot in diameter. Bark on the branches splits to reveal a reddish-brown layer. The needles are 11/16 to 1 3/16 inches long, flattened, grooved, and a waxy, bluish-green on the upper surface. Needles tend to turn upward so that the foliage of a particular branch appears flattened and as though no leaves were attached to the lower sides of the twigs. The seed cones are 2 3/8 to 4 3/4 inches long and 3/4 to 1 1/2 inches wide. Cones are dark purple, erect, and only on the uppermost branches. The common name refers to the distribution of the species in the subalpine zone.



## Subalpine Fir

*Abies lasiocarpa* subsp. *arizonica*

Ch'oh deeníní



Bryan Ungard from Santa Fe,  
New Mexico, USA (DSC\_1324)  
[CC BY-SA 2.0], via Wikimedia Commons



quinn.anya on Flickr (Flickr) [CC BY-SA 2.0], via Wikimedia Commons

Cones of *Abies lasiocarpa* of a different subspecies. Cones of subspecies *arizonica* are similar in appearance.



Russ Kleinman

# Tree of Heaven

*Ailanthus altissima*

## T'iis Nattóí



### Alternative Names:

None

### Plant Type:

Tree

### Flower Color:

White

Yellow

### Habitat:

None

### Growing Season:

Warm season

### Special Considerations:

! Noxious

**Tree of heaven** is found in middle elevations growing at the roadside and in nearby swales. It is generally an unwanted interloper. It has pinnately compound leaves which are coarsely dentate. Portions of the tree take on a reddish hue by mid-summer because the samaras become partially red in color. The winter twigs are very thick and light brown, with a light brown pith. There are around 8 or 9 bundle scars in the leaf scars. The buds are hemispheric.

*\*Description courtesy of Western New Mexico University's Vascular Plants of the Gila Wilderness.*

**Tree of Heaven**

*Ailanthus altissima*

**T'iis Nattói**



Russ Kleinman



Russ Kleinman



Russ Kleinman



Russ Kleinman





Patrick Alexander

# Utah Serviceberry

*Amelanchier utahensis*

## Dídzéi dit'odi



### Alternative Names:

Shadbush

Sarvis

Juneberry

### Plant Type:

Tree

### Flower Color:

White

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

✓ Edible berries

**Utah serviceberry** grows as a shrub or small tree, generally 6 to 13 feet tall, with white flowers in spring. Fruits are small, brown or black, apple-like, and edible to humans and animals. Serviceberry provides both forage and fruit for wildlife.

**Utah Serviceberry**

*Amelanchier utahensis*

**Dídzéi dit'odi**



Russ Kleinman



Patrick Alexander



Patrick Alexander



Russ Kleinman



Stan Shebs [GFDL, CC BY-SA 3.0 or CC BY-SA 2.5], from Wikimedia Commons

# Water Birch

*Betula occidentalis*

## K'ishchíí



### Alternative Names:

Rocky Mountain birch

### Plant Type:

Tree

### Flower Color:

Green

Yellow

### Habitat:

River valleys

### Growing Season:

Warm season

### Special Considerations:

- ✓ Stabilizes streambanks
- ✓ Provides wildlife habitat


**Water birch** is native to western North America. It is an important species for streambank stabilization and wildlife habitat. It grows along streams and in wet canyon bottoms, where it is frequently found with cottonwoods, alders, willows, and other water-loving plants. This is a small tree distinguished by catkins hanging from its twigs in the spring and by its rough, reddish-brown bark patterned with small lines called lenticels. The leaves are dark green and glossy through the growing season. The leaves are ovate with dentate edges.



## Water Birch

*Betula occidentalis*

K'ishchíí



Robert H. Mohlenbrock (USDA NRCS Plants Profile: *Betula occidentalis*) [Public domain], via Wikimedia Commons



Stan Shebs [GFDL, CC BY-SA 3.0 or CC BY-SA 2.5], from Wikimedia Commons



Stan Shebs [GFDL, CC BY-SA 3.0 or CC BY-SA 2.5], from Wikimedia Commons



Thayne Tuason [CC BY-SA 4.0], from Wikimedia Commons



Patrick Alexander

# Horsetail

*Equisetum arvense*

## Ch'ilabe'éts'óóz



### Alternative Names:

None

### Plant Type:

Other

### Flower Color:

None

### Habitat:

Grasslands

River valleys

### Growing Season:

Cool season

### Special Considerations:

! Can poison horses in hay

**Horsetail** has aerial, jointed stems, which occur in two different forms: A single, simple, cone-bearing stem grows in early spring, and a vegetative, non fertile stem grows after the first. This second stem has many whorls of slender, green-jointed branches. Roots are tuber-bearing and rhizomatous. Horsetail lacks flowers, but has a single cone,  $\frac{3}{4}$  to  $1\frac{1}{2}$  inches long. It reproduces by spores, which look like a light yellow powder. Leaves are small and scale-like, often not green, whorled, and united at the base to form a sheath around the stem.

Horsetail occurs in woods, fields, meadows and swamps, and moist soils alongside streams, rivers, and lakes, and in disturbed areas. It usually occurs on moist sites but can also be found on dry and barren sites such as roadsides, borrow pits, and railway embankments. Horsetail is sensitive to moisture stress; drought conditions result in a reduction in the production of new shoots. Associated species include rushes and sedges.

Horsetail is not an important range forage for livestock, and excessive amounts (more than 20%) in hay can cause scours, paralysis, and death in horses. Usually animals avoid the plant.


Horsetail has been used as a cough medicine for horses and as a diuretic tea. It has been made into dyes for clothing. The stems of the plant can be used for scouring and polishing. The young shoots can be eaten either cooked or raw.



**Horsetail**  
*Equisetum arvense*  
**Ch'ilabe'êts'óóz**



Mary Ellen (Mel) Harte, Bugwood.org



Mary Ellen (Mel) Harte, Bugwood.org



Patrick Alexander



Mary Ellen (Mel) Harte, Bugwood.org



Patrick Alexander





Matt Lavin

# Horsetail, Smooth

*Equisetum laevigatum*

## Ch'ilabe'éts'óóz



### Alternative Names:

None

### Plant Type:

Other

### Flower Color:

None

### Habitat:

River valleys

### Growing Season:

Year-round

### Special Considerations:

None

**Smooth horsetail** is a native cryptogam, a plant that reproduces without flowers or seeds. Horsetail prefers moist and sandy soil. The stems reach heights from 1 to 6 feet and have one dark band near the tip. The round-tipped cone at the top of the stem is yellow with varying amounts of black.

**Horsetail, Smooth**

*Equisetum laevigatum*

**Ch'ilabe'êts'óóz**



Matt Lavin



Matt Lavin



Jim Pisarowicz -  
<http://www.nps.gov/wica/photos>



Amy Smith Muise

# Lichen

*Lichen*

## Ni'hadláád



### Alternative Names:

None

### Plant Type:

None

*These aren't actually plants!*

### Flower Color:

Green

Red

Pink

White

### Habitat:

None

### Growing Season:

Year-round

### Special Considerations:

✓ Good for wildlife

**Lichens** are not a single organism but rather a symbiotic organism composed of fungi and algae. Lichens grow in a wide variety of habitats including on trees, on rocks, and in the soil. They have the ability to absorb a great deal of moisture. They fulfill a vital ecosystem function because they have the ability to gather atmospheric nitrogen and convert it into a form that plants can use. This is particularly important in the desert where there is very little nitrogen available to plants. Lichens also slowly break down rocks to create soil. Lastly, lichens do not thrive in situations with environmental pollution, and they are therefore an indicator of pollution levels. Both wildlife and humans use lichens. Deer browse on lichens, and birds use lichens as nesting material. Humans eat some lichens and also use them for dye. Lichens display diverse colors and morphology, filling the spectrum from black to red to white. Lichens can be flat or crusty or scaly. Some even grow in the form of tiny shrubs.



## Lichen

*Lichen*

Ni'hadláád



Amy Smith Muiise



Amy Smith Muiise



Amy Smith Muiise



Amy Smith Muiise

# Acknowledgments

This project was initiated by Gerald Moore, Coordinating Extension Agent, Navajo tri-state Federally-Recognized Tribes Extension Program. This handbook and the accompanying mobile-accessible web database (<http://navajorange.nmsu.edu>) were prepared by New Mexico State University (NMSU) Department of Innovative Media Research and Extension.

Support for this project was provided by the Western SARE program through USDA-NIFA. Project Director was Gerald “Jerry” Hawkes of New Mexico State University Department of Extension Animal Sciences and Natural Resources. Gerald Moore, Coordinating Extension Agent of Navajo tri-state Federally-Recognized Tribes Extension Program, had the initial vision for this project and led it in every stage.

Our deepest thanks go out to individuals and organizations who provided images, descriptions, plant identification, user testing, consultation, and support.

These include but are not limited to (in alphabetical order):

Patrick Alexander (BLM)	Matt Lavin (Montana State University)	Kathryn Thomas (Southwest Biological Science Center)
Chris Allison (NMSU)		
Nick Ashcroft (NMSU)	Andrea Long (Range Conservationist, Navajo Soil and Water Conservation District)	Lillis Urban (BLM)
Renee Benally (Dept. Rangeland, BIA)		Utah State Extension and its online tool <i>Range Plants of Utah</i>
Casey Francisco (Dept. Rangeland Management Specialist, BIA)	Felix Nez, Jr. (USDA-NRCS District Conservationist)	Judy Willetto (Range Conservationist, Navajo Dept. of Agriculture)
Ken Gishi (USDA)	Felix Nez (Diné College)	
Russ Kleinman (WNMU) and his online tool <i>Vascular Plants of the Gila Wilderness</i>	Carole Palmer (NWF Tribal Partnership Outreach)	Lawrence Yazzie (BIA).
	Nora Talkington (Navajo Natural Heritage Program)	

NMSU is an equal opportunity/affirmative action employer and educator. NMSU and the U.S. Department of Agriculture cooperating.

If you are an individual in need of an auxiliary air or accessibility service please contact [mmedia@nmsu.edu](mailto:mmedia@nmsu.edu) or (575) 646-5003 within a month of needing this aid.



# Credits

## Principal Investigators

Gerald “Jerry” Hawkes, PhD  
Jeanne Gleason, EdD  
Gerald Moore

## Content Advisors

Gerald Moore  
Nick Ashcroft, PhD  
Chris Allison, PhD  
Renee Benally  
Nora Talkington  
Kathryn Thomas, PhD

## Executive Producer

Jeanne Gleason, EdD

## Studio Production Director

Pamela N. Martinez, EdD

## Project Director

Jerry Hawkes, PhD

## Project Managers

Elizabeth Sohn  
Kathryn-Mae Eiland  
Amy Smith Muise

## Artists and Illustrator

Gerritt McGill  
Kathryn-Mae Eiland

## PDF Design and Layout

Kathryn-Mae Eiland  
Elizabeth Sohn

## Programmer

John “CC” Chamberlin

## Navajo Language Review

Tom Seaton

## Navajo Written Language

Cuyler Frank  
Arnold Clifford  
Tennell Nez  
Lorene B. Legah  
Gerald Moore

## Navajo Voice Talent

Gerald Moore  
Varian Begay  
Wilfred Big  
Cuyler Frank

## Audio Editing

Cuyler Frank

## Content Support

Levi Luchini  
J.D. Ho

## Text Work and Editorial

Amy Smith Muise  
Levi Lucchini  
J.D. Ho  
Timothy Hretz

## Website Developers

John “CC” Chamberlin  
Elizabeth Sohn

## User Testing

Amanda Armstrong  
Gerald Moore  
Jeanne Gleason, EdD  
Kathryn-Mae Eiland  
Amy Smith Muise  
Elizabeth Sohn

## Research and Evaluation

Gerald Moore  
Jeanne Gleason, EdD  
Amy Smith Muise

## Studio Support

Esther Aguirre  
Luis Barraza  
Amy Chacon  
Patricia Clark

## Special Thanks To:

Window Rock focus  
group participants

Copyright 2018 New Mexico State University. Partially funded by the Western Sustainable Agriculture Research and Education Program ([westernsare.org](http://westernsare.org); 435.797.2257), project EW15-023. Programs and projects supported by Western SARE are equally open to all people. NMSU is an equal opportunity/affirmative action educator and employer.

Our gratitude and appreciation to Tom Seaton and all the elders who have kept Navajo language and plant knowledge alive.



## References

- Allison, C. D., Turner, J. L., & Wenzel, J. C. (2016). *Poisonous plants of New Mexico rangelands* (Circular 678). Retrieved from New Mexico State University, College of Agricultural, Consumer, and Environmental Sciences website: [https://aces.nmsu.edu/pubs/\\_circulars/CR678/](https://aces.nmsu.edu/pubs/_circulars/CR678/)
- Center for Invasive Species and Ecosystem Health. (n.d.) Invasive and Exotic Species of North America. *Invasive.org*. Retrieved from <https://www.invasive.org/>
- Fox, W. E. III, Allred, K. W., & Roalson, E. H. (2010). *A guide to the common locoweeds and milkvetches of New Mexico* (Circular 557). Retrieved from New Mexico State University, College of Agricultural, Consumer, and Environmental Sciences website: [https://aces.nmsu.edu/pubs/\\_circulars/CR557/](https://aces.nmsu.edu/pubs/_circulars/CR557/)
- Heil, K., O'Kane, S. Jr., Reeves, L. M., & Clifford, A. (2013). *Flora of the Four Corners Region: Vascular Plants of the San Juan River Drainage: Arizona, Colorado, New Mexico, and Utah* (Monographs in Systemic Botany: Vol. 124). St. Louis: Missouri Botanical Garden Press.
- Kleinman, R. (n.d.). *Vascular Plants of the Gila Wilderness*. Retrieved from <https://wnmu.edu/academic/nspages/gilaflora/index.html>
- Ralphs, M. (n.d.) *Rangeland plants poisonous to livestock*. Retrieved from <https://www.ars.usda.gov/ARSUserFiles/20801500/.../Dr.MichaelRalphsLecture.ppt>
- United States Department of Agriculture. (2018, September 24). *Fire Effects Information System (FEIS): Syntheses about fire ecology and fire regimes in the United States*. Retrieved from <https://www.feis-crs.org/feis/>
- United States Department of Agriculture Natural Resources Conservation Service, National Plant Data Team. (2018, October 11). *The PLANTS Database*. Retrieved from <http://plants.usda.gov>
- University of California, Berkeley. (2018). Jepson eFlora. The Jepson Herbarium. Retrieved from <http://ucjeps.berkeley.edu/eflora/>
- University of Texas at Austin. (n.d.) Find Plants. Lady Bird Johnson Wildflower Center. Retrieved from <https://www.wildflower.org/plants/>
- Utah State University Extension. (n.d.) *Range Plants of Utah*. Retrieved from <https://extension.usu.edu/rangeplants/>

# Glossary

## **Navajo Nation/Diné Bikéyah**

Covers 27,425 square miles in northeastern Arizona, southeastern Utah, and northwestern New Mexico.

Navajo people have strong family bonds and enjoy a rich cultural, spiritual, and daily life, often based on small-scale farming and ranching. These human strengths, traditional lifestyles, knowledge, values, and resources are the foundation of the Navajo people (Diné).

## **Rangeland**

Open country or woodland used for grazing animals.

## **Official definition from USDA APHIS**

Any plant or plant product that can directly or indirectly injure or cause damage to crops, livestock, poultry or other interests of agriculture, irrigation, navigation, the natural resources of the United States, the public health, or the environment.

For more information about introduced, invasive, and noxious plants of the U.S.:

<https://plants.usda.gov/java/noxiousDriver#invasive>

## **Toxic/Poisonous**

When eaten or touched, may cause harm or death.

A poisonous or toxic plant may be harmful or fatal to some animals and not to others, or it may be harmful only at certain quantities or when eaten in combination with certain other plants. Some are only toxic at certain times of the year or at a certain growing stage of the plant. Stress from drought, freezing, or physical damage can also sometimes cause certain plants to be poisonous.

## **Introduced**

A plant growing outside its historical range.

## **Invasive**

A plant that tends to spread beyond its historical range, often in response to disturbance.

## **Noxious**

A plant that is particularly troublesome.

## **Annual plants**

A plant that lives for one year, during which it completes its life cycle.

## **Biennial plants**

A plant that grows for two years, producing its fruit and seed in the second year.

## **Perennial plants**

A plant that lives for three years or more. This includes plants that live just a few years and those that live many, many years.

## **Habitat Types**

### ***Brushlands***

Dominated by shrubs and bushes, such as sagebrush, four-winged saltbush, blackbrush, greasewood, etc.

### ***Grasslands***

Mainly grassy; may include occasional shrubs, bushes, trees, or cacti.

### ***Piñon-Juniper Woodland***

Patchy forested area of mainly shorter trees. May also include shrubs and grasses.

### ***Ponderosa Pine / Mixed Conifer***

Fully developed forest dominated by tall trees. May also include shrubs and grasses.

## Glossary

*continued*

### Habitat Types: *(continued)*

#### **River Valley**

Area around a watercourse, which may include sections dominated by grasses, trees, shrubs, or wetland plants.

#### **Wastelands**

Barren or disturbed areas of sparse or weedy vegetation; may include previously cultivated land.

#### **Cool Season Plants**

Plants which begin growth early in the spring and make most of their growth during the cool weather of spring and early fall. Examples are western wheatgrass and Indian ricegrass.

#### **Warm Season Plants**

Plants which begin growth and make most of their growth during the hot summer months. Examples are blue grama and sand dropseed.

### **Plant Terms**

#### **Stem**

Supports a plant above ground and connects roots to leaves.

#### **Node**

Growth point along a plant's stem, often where a leaf stalk connects.

#### **Prostrate**

A plant whose branches lie directly on the ground or just above it.

#### **Upright**

A plant that grows more or less vertically.

#### **Rhizomes**

Underground stems that put out shoots and roots.

#### **Stolons**

Creeping stems that can root into the ground.

### **Navajo Endangered Species List Definitions**

From the Navajo Nation Division of Natural Resources Department of Fish and Wildlife

#### **Group 1:**

Those species or subspecies that no longer occur on the Navajo Nation.

#### **GROUP 2 (G2) & GROUP 3 (G3):**

"Endangered" - Any species or subspecies whose prospects of survival or recruitment within the Navajo Nation are in jeopardy or are likely within the foreseeable future to become so.

**G2:** A species or subspecies whose prospects of survival or recruitment are in jeopardy.

**G3:** A species or subspecies whose prospects of survival or recruitment are likely to be in jeopardy in the foreseeable future.

#### **Group 4:**

Any species or subspecies for which the Navajo Nation Department of Fish and Wildlife (NNDFW) does not currently have sufficient information to support their being listed in G2 or G3 but has reason to consider them. The NNDFWL will actively seek information on these species to determine if they warrant inclusion in a different group or removal from the list.

#### **Related to stems:**

**Alternate:** Arrangement of leaves on stems where one leaf or stem forms at one node and the next from another node on the other side of the stem.

**Basal:** Growth emerges from the base of the plant.

**Opposite:** Arrangements where two plant stems or leaves form at the same node on opposite sides of the stem.



## Glossary

*continued*

### Related to stems: *(continued)*

**Whorled:** Arrangement where three or more stems or leaves are clustered around the same node.

### Related to leaves:

**Compound:** a leaf that is actually made up of leaflets joined together.

**Dentate:** toothed.

**Leaflet:** a small leaflike structure that is actually part of a larger unit called a compound leaf.

**Lobed:** (of a leaf) having rounded sections.

**Lanceolate:** (of a leaf's shape) long and narrow, but wider in the middle.

**Linear:** long and very narrow.

**Margin:** the outer edge of a leaf.

**Palmate:** palm shaped with lobes or leaflets radiating from a central point.

**Palmately compound:** a leaf composed of leaflets that radiate from a central point.

**Palmately lobed:** having rounded sections that radiate from a central point at the base of the leaf.

**Pinnately lobed:** having rounded sections that are arranged along the central axis of the leaf.

**Serrate:** much like being toothed except that the teeth all face in a particular direction.

**Simple:** describes a leaf that is not divided into leaflets. What most people think of as a leaf is a simple leaf.

**Oblong:** almost like a rectangle but with rounded edges.

**Ovate:** egg-shaped with the wider part of the leaf near where it joins the stem.

**Tomentose:** having dense, velvety, fuzzy hairs

### Related to grasses:

**Bunchgrass:** grass with a compact (clumping), rather than spreading, growth habit.

**Culms:** stems or stalks of grasses and sedges, usually hollow and jointed.

**Floret:** just like other plants, grasses have flowers called florets, which are very small and inconspicuous. They later turn into individual seeds.

**Glabrous:** having a smooth surface

**Glumes:** bracts (see definition for bracts under flower) around the spikelet.

**Inflorescence:** refers to the entire portion of the grass that bears flowers or seeds. In some cases, the inflorescence is small. In other cases, quite large. For a good example of the latter, see the entry on Uruguayan pampas grass.

**Panicle:** a grass seedhead that has a very branched structure: a main stem with branches that in turn have smaller stalks (spikelets) with seeds on each stalk.

**Pubescent:** covered with fine hairs

**Raceme:** a central stalk with small branches with spikelets attached to them.

**Rachis:** shaft or stem.

**Spike:** a grass seedhead where the spikelets are directly attached to the main stem.

**Spikelet:** grass flowers and seeds are arranged along stems ranging in size and location in the plant's overall structure. A spikelet is something like a twig in relation to a tree. It is a very small "stem" that provides an axis along which flowers grow.

**Tillers:** new shoots that spring up off of stolons.

**Related to flowers:**

**Anther:** the part of the stamen that contains the pollen; it usually has the appearance of an oval, yellow sac or ball.

**Bract:** a leaflike structure at the base of a flower. Bracts differ from sepals because they are attached directly to the stem, and they are separate from the flower or inflorescence.

**involucre:** a whorl of bracts at the base of a flower; involucre is quite noticeable in sunflowers and other members of the aster family.

**Pistil:** female part of a flower that receives pollen.

**Sepal:** sepals can often be confused with bracts because they also can be leaflike, and they enclose the petals of a flower. Sepals are particularly noticeable on roses because they come up to cup the flower.

**Stamen:** male part of the flower comprising a filament (like a little stem) with a large amount of pollen at the end of it.

# Index

## Common Name

agarita 110–111  
 agrito 110–111  
 alder 319–320, 350  
 alfalfa 277–278  
 alfilaria 211–212  
 algerita 110–111  
 American bellflower 185–186  
 American plum 336–337  
 antelope bitterbrush 112–113  
 antelopehorn 173–174, 261  
 Arizona alder 319–320  
 arrowgrass 101  
 ashen, milkvetch 255–256  
 Aster 207–208  
 banana yucca 317–318  
 beard grass 12  
 beardtongue 275–276  
 bee balm 225–226  
 beeplant, Rocky Mountain 175–176  
 big heavy 332–333  
 birchleaf mountain mahogany 330–331  
 birdbeak 177–178  
 biscuitroot 114–115  
 bitterbrush 112–113  
 bitterweed 179–180  
 blackbent 77–78  
 blackbrush 116–117  
 black jack 332–333  
 blanket flower 181–182  
 blanketflower 181–182  
 bluebell bellflower 185–186  
 bluebells 185–186  
 blue-eyed grass 183–184  
 blue-eyed grass, Arizona 183–184  
 blue-eyed grass, stiff 183–184  
 blue flag 229–230  
 bluegrass, Kentucky 10–11  
 bluestem, little false 12  
 bobtail barley 44–45  
 bottlebrush squirreltail 13  
 broadleaved pepperweed 277–278  
 brome, California 18–19  
 brome, field 14–15  
 brome, Japanese 16–17  
 brome, mountain 18–19, 93  
 brome, red 20–21, 42  
 brome, ripgut 22–23  
 brome, Schrader's 83–84  
 brome, smooth 24–25  
 broom snakeweed 158–159

buckbrush 112–113  
 buckwheat 118–120, 271  
 buckwheat, sand 118–120  
 buckwheat, slender 118–120  
 buena mujer 207–208  
 bull pine 332–333  
 bulrush 102–103, 305  
 bundle flower 187–188  
 California burclover 189–190  
 camelthorn 121–122  
 cane cholla 313–314  
 canyagra 271–272  
 chamiza 128–129  
 cheatgrass 26–27, 42, 87  
 chokecherry 123–124, 160  
 cholla 313–314  
 chufa 106–107  
 clasping pepperweed 277–278  
 cliff fendlerbush 125–126  
 cliffrose 127  
 cocklebur 191–192  
 Colorado piñon 321–322  
 Colorado pinyon 321–322  
 common pinyon 321–322  
 common purslane 193–194  
 common tule 102–103  
 copperweed 195  
 cordgrass, alkali 28–29  
 cottonwood 93, 350  
 cow parsnip 114, 196–197  
 coyote willow 323–324, 350  
 curlycup gumweed 198–199  
 curly dock 269–270  
 currant-of-Texas 110–111  
 daisy, rock 200–201  
 dandelion 202–203  
 death camas 204–205, 283  
 death camas, alcove 204  
 desert sand verbena 206  
 devilweed 207–208  
 dock root 271–272  
 dropseed, giant 30–31  
 dropseed, hairy 32–33  
 dropseed, pine 32–33  
 dropseed, sand 34–35  
 dropseed, spike 36–37  
 dwarf stick pea 187  
 Emory's milkvetch 257–258  
 Emory's rock daisy 200–201  
 epazote 312  
 evening primrose 209–210  
 faitours-grass 249–250

fescue, Arizona 40–41  
 fescue, six weeks 42–43  
 field bindweed 170  
 filaree 42, 211–212  
 flag lily 229–230  
 foothill biscuit root 114–115  
 foothill deathcamas 204–205  
 four o'clock 213–214  
 fourwinged saltbrush 128–129  
 four-winged saltbush 128–129  
 fourwing saltbush 128–129  
 foxtail barley 44–45  
 fragrant evening primrose 209–210  
 Fremont cottonwood 325–326  
 galleta 46–47  
 galleta, James' 46–47  
 Gambel oak 327  
 geranium 215–216  
 geranium, Fremont 215–216  
 geranium, purple 215  
 geranium, Richardson's 215  
 geranium, sticky purple 215  
 ghostberry 160–161  
 giant biscuit root 114–115  
 giant whiteweed 277–278  
 gilia 217–218  
 gilia, scarlet 217–218  
 gilia, shy 217–218  
 globemallow 219–220  
 goatgrass, jointed 48–49  
 goathead 171–172  
 goldenweed 233–234  
 grama, black 50–51  
 grama, blue 52–53, 87  
 grama, sideoats 54–55  
 Gray's biscuit root 114–115  
 greasewood 42, 130–131  
 green ephedra 132–133  
 green-flowered milkweed 173–174  
 groundsel 285–286  
 groundsel, threadleaf 221–222  
 guapile 213–214  
 gumbo lily 209–210  
 half-moon milkvetch 253–254  
 halogeton 223–224  
 hardheads 241–242  
 hardstem bulrush 102–103  
 hardstem tule 102–103  
 harebell 185–186  
 hoary cress 308–309  
 horsemint 225–226  
 horsetail 352–353



## Index

*continued*

### Common Name *continued*

- horsetail, smooth 354–355  
Indian celery 196–197  
Indian paintbrush 227–228  
Indian rhubarb 196–197  
iris 229–230  
Jacob's ladder 231–232  
Jacobs ladder 231–232  
jimmyweed 233–234  
jimsonweed 235–236  
johnsongrass 56–57  
joint fir 132–133  
juneberry 348–349  
junegrass 58–59, 93  
junegrass, prairie 58–59  
juniper 68, 114, 328–329  
knapweed 237–238  
knapweed, creeping 241–242  
knapweed, diffuse 239–240  
knapweed, Russian 241–242  
knapweed, spotted 237–238  
knapweed, spreading 239–240  
knapweed, tumble 239–240  
knapweed, white 239–240  
kochia 243–244  
lamb's quarters 245–246  
lambsquarters 245–246  
larkspur 247–248  
leafy spurge 249–250  
lichen 356–357  
little hogweed 193–194  
liver lily 229–230  
loco, Ellis 251–252  
loco, Patterson 255–256  
loco, rattleweed 253–254, 255  
locoweed 257–258  
loco, woolly 255–256  
London rocket 265–266  
longleaf ephedra 132–133  
lupine 259–260  
mahonia 134–135  
milkweed 173–174, 261–262  
milkweed, broadleaf 261–262  
Mormon tea 132–133  
mountain bluet 241–242  
mountain mahogany 330–331  
mountain mint 225–226  
mountain snowberry 160–161  
muhly, mountain 60–61  
muhly, ring 62–63  
muhly, sandhill 64–65  
muhly, spike 66–67  
mullein 263–264  
mustard 265–266  
mustard, tansy 265–266  
mutton grass 68  
narrowleaf willow 323–324  
narrowleaf yucca 317–318  
Navajo spinach 175–176  
Navajo tea 267–268  
Navajo tobacco 269–270  
Navajo wild carrot 271–272  
needle-and-threadgrass 69–70, 75, 87  
needlegrass, desert 71–72  
needlegrass, scribner 73–74  
New Mexico feathergrass 75–76  
nightshade 273–274  
nodding plumeless thistle 297–298  
nutsedge, yellow 106–107, 108  
oleaster 338–339  
oneseed juniper 328–329  
oregano de la Sierra 225–226  
Oregon grape 134–135  
Oregon grape holly 134–135  
pale wolfberry 168–169  
penstemon 275–276  
penstemon, Navajo 275  
pepperweed 277–278  
perennial pepperweed 277–278  
Persian olive 338–339  
pingue 136–137, 281  
piñon pine 114, 321–322  
plains prickly pear 315–316  
poison ivy 279–280  
ponderosa pine 160, 332–333  
ponderosa white 332–333  
prairie sagewort 150–151  
prickly pear 315–316  
prickly pear, brown spined 315–316  
prickly pear, Engelmann 315  
prickly pear, pancake 315  
prickly pear, Pott's 315  
prickly pear, purple 315  
puncturevine 171–172  
pursley 193–194  
pushki 196–197  
quaking aspen 93, 160, 247, 334–335  
rabbitbrush 138–139  
rabbitbrush, Douglas 140–141  
rabbitbrush, green 140–141  
rabbitbrush, yellow 140–141  
ragweed 42, 243  
ragwort 285–286  
rattleweed 253–254  
rayless goldenrod 233–234  
rayless greenthread 267–268  
red plum 336–337  
red root 193–194  
redstem filaree 211–212  
red stem peavine 257–258  
redstem stork's bill 211–212  
redtop 77–78, 305  
ricegrass, Indian 79–80  
Rio Grande cottonwood 325–326  
Rocky Mountain birch 350–351  
Rocky Mountain penstemon 275–276  
rough-fruited cinquefoil 293–294  
roundleaf buffaloberry 142–143  
rubber rabbitbrush 138–139  
rubberweed, bitter 136, 281–282  
rushes 104–105, 352  
Russian cornflower 241–242  
Russian olive 338–339  
sacaton, alkali 38–39  
sacaton, giant 30–31  
sacred thorn-apple 235–236  
sagebrush 87, 227  
sagebrush, big 42, 93, 114, 144–145, 148  
sagebrush, bigelow 146–147  
sagebrush, black 114, 148–149  
sagebrush, flat 146–147  
sagebrush, fringed 150–151  
sagebrush, sand 152–153  
saltcedar 340–341  
salt grass 38–39  
saltgrass 81–82  
sandbar willow 323–324  
sandbur, field 83–84  
sand dock 271–272  
sarvis 348–349  
scarlet globemallow 219–220  
Scotch cottonthistle 301–302  
seaside arrowgrass 101  
sedges 108–109, 305, 352  
sego lily 204, 283–284  
senecio 285–286  
shadbush 348–349  
shadscale 154–155  
Siberian elm 342–343  
silverberry 338–339  
silverleaf 142–143  
silverleaf nightshade 273–274

## Index

*continued*

### Common Name *continued*

- six weeks fescue 42–43
- skunkbrush 156–157
- skunkbush 156–157
- snakeweed 158–159
- sneezeweed 287–288
- snowberry 114, 160–161
- southern goldenbush 233–234
- southern jimmyweed 233–234
- spear-grass 69–70
- spider antelope horns 173–174
- spreading wallflower 289–290
- spruce 247
- spurge 249–250
- squirreltail barley 44–45
- stemless primrose 209–210
- St. Johnswort 291–292
- storksbill 42, 211–212
- subalpine fir 344–345
- sulphur cinquefoil 293–294
- sunflower 295–296
- sunflower, giant 295–296
- tall Mormon tea 132–133
- tall whitetop 277–278
- thistle, barbwire Russian 303–304
- thistle, Canada 299–300
- thistle, musk 297–298
- thistle, prickly Russian 303–304
- thistle, Russian 303–304
- thistle, Scotch 301–302
- thistle, slender 303–304
- thistle, spineless Russian 303–304
- three-awn, oldfield 85
- three-awn, purple 87
- three-awn, red 87
- three-awns (annual) 85–86
- three-awn, six weeks 85
- three-awns (perennial) 87–88
- trailing windmills 213–214
- tree of heaven 346–347
- tufted evening primrose 209–210
- tufted hairgrass 89–90
- tufted horse grass 89–90
- tule 102–103
- tule rush 102–103
- tumbleweed 303–304
- Turkestan thistle 241–242
- twoneedle pinyon 321–322
- Utah serviceberry 348–349
- Utah snowberry 160–161
- valley cottonwood 325–326
- viscid bulrush 102–103
- walking stick cholla 313–314
- water birch 350–351
- water hemlock 114, 305–306
- water jacket 168–169
- waxberry 160–161
- wax currant 162–163
- western blue flag 229–230
- western horsenettle 273–274
- western yarrow 93, 307
- wheatgrass, crested 91–92
- wheatgrass, slender 93–94
- wheatgrass, western 95–96
- white sage 166–167
- white stemless evening primrose 209–210
- whitetop 308–309
- wild currant 110–111
- wild olive 338–339
- wild rhubarb 271–272
- wild rose 164–165
- wildrye, Canadian 97–98
- wildrye, nodding 97–98
- wildrye, Russian 99–100
- winterfat 166–167
- wolfberry 168–169
- wolfberry, Anderson 168
- wolfberry, pale 168
- wolf's milk 249–250
- wooly Indian wheat 310–311
- wooly yarrow 307
- wormsweed 312
- woul 87–88
- yellow Jacob's ladder 231
- yellow starthistle 237–238
- yucca 317–318

## Index

*continued*

### Flower Color

#### Blue

desert sand verbena 206  
field sandbur 83–84  
filaree 211–212  
horsemint 225–226  
jimsonweed 235–236  
lambsquarters 245–246  
larkspur 247–248  
lupine 259–260  
milkweed 261–262  
rushes 104–105  
thistle, Canada 299–300  
thistle, musk 297–298  
tumbleweed 303–304

#### Brown

camelthorn 121–122  
Gambel oak 327  
johnsongrass 56–57  
Navajo tobacco 269–270  
nutsedge, yellow 106–107  
rushes 104–105  
sagebrush, sand 152–153  
saltgrass 81–82  
sedges 108–109

#### Green

alder 319–320  
antelopehorn 173–174  
arrowgrass 101  
bluegrass, Kentucky 10–11  
bluestem, little false 12  
bottlebrush squirreltail 13  
brome, Japanese 16–17  
brome, mountain 18–19  
brome, ripgut 22–23  
brome, smooth 24–25  
cocklebur 191–192  
Colorado piñon 321–322  
dropseed, giant 30–31  
dropseed, pine 32–33  
dropseed, sand 34–35  
dropseed, spike 36–37  
fescue, six weeks 42–43  
foxtail barley 44–45  
Fremont cottonwood 325–326  
Gambel oak 327  
grama, black 50–51  
grama, blue 52–53  
grama, sideoats 54–55  
greasewood 130–131  
halogeton 223–224  
Indian paintbrush 227–228  
James' galleta 46–47

johnsongrass 56–57  
jointed goatgrass 48–49  
junegrass, prairie 58–59  
juniper 328–329  
knapweed, Russian 241–242  
kochia 243–244  
leafy spurge 249–250  
lichen 356–357  
loco, Ellis 251–252  
milkweed 261–262  
Mormon tea 132–133  
muhly, mountain 60–61  
muhly, ring 62–63  
muhly, sandhill 64–65  
muhly, spike 66–67  
mutton grass 68  
Navajo tobacco 269–270  
Navajo wild carrot 271–272  
needle-and-threadgrass 69–70  
needlegrass, desert 71–72  
needlegrass, scribner 73  
New Mexico feathergrass 75–76  
poison ivy 279–280  
rabbitbrush, green 140–141  
ricegrass, Indian 79–80  
sacaton, alkali 38–39  
sagebrush, bigelow 146–147  
sagebrush, black 148–149  
sagebrush, fringed 150–151  
sedges 108–109  
Siberian elm 342–343  
snakeweed 158–159  
thistle, musk 297–298  
three-awns (annual) 85–86  
tufted horse grass 89–90  
water birch 350–351  
wheatgrass, crested 91–92  
wheatgrass, slender 93–94  
wheatgrass, western 95–96  
wildrye, Canadian 97–98  
wildrye, Russian 99–100  
wooly Indian wheat 310–311  
wormsweed 312

#### Pink

alder 319–320  
beeplant, Rocky Mountain 175–176  
birdbeak 177–178  
blanket flower 181–182  
buckwheat 118–120  
bundle flower 187–188  
camelthorn 121–122

desert sand verbena 206  
field bindweed 170  
field sandbur 83–84  
filaree 211–212  
four o'clock 213–214  
gilia 217–218  
knapweed 237–238  
knapweed, Russian 241–242  
lichen 356–357  
loco, rattleweed 253–254  
Navajo tobacco 269–270  
Navajo wild carrot 271–272  
penstemon 275–276  
ponderosa pine 332–333  
prickly pear 315–316  
quaking aspen 334–335  
redtop 77–78  
rushes 104–105  
snowberry 160–161  
wild rose 164–165

#### Purple

beeplant, Rocky Mountain 175–176  
bluebells 185–186  
blue-eyed grass 183–184  
cholla 313–314  
desert sand verbena 206  
field sandbur 83–84  
filaree 211–212  
geranium 215–216  
horsemint 225–226  
iris 229–230  
Jacob's ladder 231–232  
jimsonweed 235–236  
knapweed, Russian 241–242  
lambsquarters 245–246  
larkspur 247–248  
loco, wooly 255–256  
lupine 259–260  
milkweed 261–262  
Mormon tea 132–133  
mustard 265–266  
nightshade 273–274  
penstemon 275–276  
prickly pear 315–316  
red stem peavine 257–258  
rushes 104–105  
thistle, Canada 299–300  
thistle, musk 297–298  
thistle, Scotch 301–302  
tumbleweed 303–304  
wolfberry 168–169



## Index

*continued*

### Flower color *continued*

#### Red

alder 319–320  
birdbeak 177–178  
blanket flower 181–182  
camelthorn 121–122  
desert sand verbena 206  
field bindweed 170  
field sandbur 83–84  
filaree 211–212  
four o'clock 213–214  
four-winged saltbush 128–129  
geranium 215–216  
gilia 217–218  
halogeton 223–224  
knapweed 237–238  
lichen 356–357  
loco, rattleweed 253–254  
loco, woolly 255–256  
Navajo tobacco 269–270  
Navajo wild carrot 271–272  
penstemon 275–276  
quaking aspen 334–335  
redtop 77–78  
rushes 104–105  
saltcedar 340–341  
scarlet globemallow 219–220  
wax currant 162–163  
wild rose 164–165

#### White

antelope bitterbrush 112–113  
antelopehorn 173–174  
buckwheat 118–120  
bundle flower 187–188  
chokecherry 123–124  
cliff fendlerbush 125–126  
cliffrose 127  
cow parsnip 196–197  
coyote willow 323–324  
daisy, rock 200–201  
death camas 204–205  
devilweed 207–208  
evening primrose 209–210  
field bindweed 170  
four o'clock 213–214  
halogeton 223–224  
iris 229–230  
jimsonweed 235–236  
knapweed, diffuse 239–240  
lichen 356–357  
lupine 259–260  
milkweed 261–262  
Mormon tea 132–133

mountain mahogany 330–331  
mustard 265–266  
nightshade 273–274  
penstemon 275–276  
pepperweed 277–278  
poison ivy 279–280  
quaking aspen 334–335  
red plum 336–337  
sagebrush, big 144–145  
saltcedar 340–341  
sego lily 283–284  
skunkbush 156–157  
snowberry 160–161  
tree of heaven 346–347  
tumbleweed 303–304  
Utah serviceberry 348–349  
water hemlock 305–306  
western yarrow 307  
whitetop 308–309  
wild rose 164–165  
winterfat 166–167  
yucca 317–318

#### Yellow

algerita 110–111  
antelope bitterbrush 112–113  
birdbeak 177–178  
biscuitroot 114–115  
bitterweed 179–180  
blackbrush 116–117  
blanket flower 181–182  
blue-eyed grass 183–184  
brome, field 14–15  
brome, red 20–21  
buckwheat 118–120  
bundle flower 187–188  
California burclover 189–190  
cheatgrass 26–27  
cliffrose 127  
common purslane 193–194  
copperweed 195  
cordgrass, alkali 28–29  
coyote willow 323–324  
curlycup gumweed 198–199  
daisy, rock 200–201  
dandelion 202–203  
death camas 204–205  
dropseed, pine 32–33  
dropseed, sand 34–35  
dropseed, spike 36–37  
evening primrose 209–210  
fescue, Arizona 40–41  
four o'clock 213–214

four-winged saltbush 128–129  
Fremont cottonwood 325–326  
Gambel oak 327  
grama, black 50–51  
grama, blue 52–53  
groundsel, threadleaf 221–222  
Indian paintbrush 227–228  
iris 229–230  
Jacob's ladder 231–232  
jimmyweed 233–234  
knapweed 237–238  
leafy spurge 249–250  
loco, Ellis 251–252  
mountain mahogany 330–331  
mullein 263–264  
mustard 265–266  
Navajo tea 267–268  
Navajo tobacco 269–270  
nutsedge, yellow 106–107  
Oregon grape 134–135  
ponderosa pine 332–333  
prickly pear 315–316  
puncturevine 171–172  
rabbitbrush 138–139  
rabbitbrush, green 140–141  
ricegrass, Indian 79–80  
roundleaf buffaloberry 142–143  
rubberweed, bitter 281–282  
Russian olive 338–339  
sagebrush, big 144–145  
sagebrush, black 148–149  
sagebrush, sand 152–153  
saltgrass 81–82  
sego lily 283–284  
senecio 285–286  
shadscale 154–155  
skunkbush 156–157  
snakeweed 158–159  
sneezeweed 287–288  
spreading wallflower 289–290  
St. Johnswort 291–292  
sulphur cinquefoil 293–294  
sunflower 295–296  
tree of heaven 346–347  
water birch 350–351  
yucca 317–318

## Index

*continued*

### Growing Season

#### Cool Season

algerita 110–111  
bluegrass, Kentucky 10–11  
bottlebrush squirreltail 13  
brome, field 14–15  
brome, Japanese 16–17  
brome, mountain 18–19  
brome, smooth 24–25  
buckwheat 118–120  
camelthorn 121–122  
cheatgrass 26–27  
coyote willow 323–324  
dandelion 202–203  
death camas 204–205  
fescue, Arizona 40–41  
fescue, six weeks 42–43  
filaree 211–212  
foxtail barley 44–45  
geranium 215–216  
halogeton 223–224  
horsetail 352–353  
horsetail, smooth 354–355  
jointed goatgrass 48–49  
junegrass, prairie 58–59  
lambsquarters 245–246  
lichen 356–357  
lupine 259–260  
Mormon tea 132–133  
mustard 265–266  
needle-and-threadgrass 69–70  
needlegrass, desert 71–72  
New Mexico feathergrass 75–76  
prickly pear 315–316  
quaking aspen 334–335  
rabbitbrush 138–139  
redtop 77–78  
ricegrass, Indian 79–80  
roundleaf buffaloberry 142–143  
rushes 104–105  
scarlet globemallow 219–220  
sedges 108–109  
sego lily 283–284  
snakeweed 158–159  
snowberry 160–161  
subalpine fir 344–345  
three-awns (annual) 85–86  
water hemlock 305–306  
wheatgrass, crested 91–92  
wheatgrass, slender 93–94  
wild rose 164–165  
wildrye, Canadian 97–98  
wildrye, Russian 99–100

yucca 317–318

#### Warm Season

alder 319–320  
antelope bitterbrush 112–113  
antelopehorn 173–174  
arrowgrass 101  
beepiant, Rocky Mountain 175–176  
birdbeak 177–178  
biscuitroot 114–115  
blackbrush 116–117  
blanket flower 181–182  
bluebells 185–186  
blue-eyed grass 183–184  
bluestem, little false 12  
brome, red 20–21  
brome, ripgut 22–23  
bulrush 102–103  
bundle flower 187–188  
California burclover 189–190  
chokecherry 123–124  
cholla 313–314  
cliff fendlerbush 125–126  
cliffrose 127  
cocklebur 191–192  
common purslane 193–194  
cordgrass, alkali 28–29  
cow parsnip 196–197  
curlycup gumweed 198–199  
daisy, rock 200–201  
desert sand verbena 206  
devilweed 207–208  
dropseed, giant 30–31  
dropseed, pine 32–33  
dropseed, sand 34–35  
dropseed, spike 36–37  
evening primrose 209–210  
field bindweed 170  
field sandbur 83–84  
four o'clock 213–214  
four-winged saltbush 128–129  
Fremont cottonwood 325–326  
Gambel oak 327  
gilia 217–218  
grama, black 50–51  
grama, blue 52–53  
grama, sideoats 54–55  
greasewood 130–131  
groundsel, threadleaf 221–222  
horsemint 225–226  
horsetail, smooth 354–355  
Indian paintbrush 227–228

iris 229–230

Jacob's ladder 231–232  
James' galleta 46–47  
jimmyweed 233–234  
jimsonweed 235–236  
johnsongrass 56–57  
juniper 328–329  
knapweed 237–238  
knapweed, diffuse 239–240  
knapweed, Russian 241–242  
kochia 243–244  
larkspur 247–248  
leafy spurge 249–250  
lichen 356–357  
loco, Ellis 251–252  
loco, rattleweed 253–254  
loco, woolly 255–256  
milkweed 261–262  
mountain mahogany 330–331  
muhly, mountain 60–61  
muhly, ring 62–63  
muhly, sandhill 64–65  
muhly, spike 66–67  
mullein 263–264  
mutton grass 68  
Navajo tea 267–268  
Navajo tobacco 269–270  
Navajo wild carrot 271–272  
nightshade 273–274  
nutsedge, yellow 106–107  
Oregon grape 134–135  
pepperweed 277–278  
pingue 136–137  
poison ivy 279–280  
ponderosa pine 332–333  
puncturevine 171–172  
rabbitbrush, green 140–141  
red plum 336–337  
red stem peavine 257–258  
roundleaf buffaloberry 142–143  
Russian olive 338–339  
sacaton, alkali 38–39  
sagebrush, big 144–145  
sagebrush, bigelow 146–147  
sagebrush, black 148–149  
sagebrush, fringed 150–151  
sagebrush, sand 152–153  
saltcedar 340–341  
saltgrass 81–82  
sedges 108–109  
senecio 285–286  
shadscale 154–155

## Index

*continued*

### Growing Season

Warm Season *continued*

Siberian elm 342–343  
skunkbush 156–157  
snakeweed 158–159  
sneezeweed 287–288  
spreading wallflower 289–290  
St. Johnswort 291–292  
subalpine fir 344–345  
sulphur cinquefoil 293–294  
sunflower 295–296  
thistle, Canada 299–300  
thistle, musk 297–298  
thistle, Scotch 301–302  
three-awns (annual) 85–86  
three-awns (perennial) 87–88  
tree of heaven 346–347  
tufted horse grass 89–90  
tumbleweed 303–304  
Utah serviceberry 348–349  
water birch 350–351  
wax currant 162–163  
western yarrow 307  
wheatgrass, western 95–96  
whitetop 308–309  
winterfat 166–167  
wolfberry 168–169  
wooly Indian wheat 310–311  
wormsweed 312



## Index

*continued*

### Habitat

#### Brushlands

algerita 110–111  
antelope bitterbrush 112–113  
beepplant, Rocky Mountain 175–176  
birdbeak 177–178  
biscuitroot 114–115  
blackbrush 116–117  
bottlebrush squirreltail 13  
brome, field 14–15  
brome, red 20–21  
buckwheat 118–120  
California burclover 189–190  
camelthorn 121–122  
chokecherry 123–124  
cliff fendlerbush 125–126  
cliffrose 127  
cocklebur 191–192  
cow parsnip 196–197  
curlycup gumweed 198–199  
daisy, rock 200–201  
dandelion 202–203  
death camas 204–205  
dropseed, giant 30–31  
field bindweed 170  
field sandbur 83–84  
filaree 211–212  
four-winged saltbush 128–129  
gilia 217–218  
greasewood 130–131  
groundsel, threadleaf 221–222  
halogeton 223–224  
Indian paintbrush 227–228  
jimmyweed 233–234  
knapweed 237–238  
knapweed, diffuse 239–240  
knapweed, Russian 241–242  
leafy spurge 249–250  
loco, Ellis 251–252  
loco, rattleweed 253–254  
loco, wooly 255–256  
lupine 259–260  
Mormon tea 132–133  
muhly, ring 62–63  
Navajo tea 267–268  
Navajo wild carrot 271–272  
needlegrass, desert 71–72  
New Mexico feathergrass 75–76  
nightshade 273–274  
Oregon grape 134–135  
penstemon 275–276  
pingue 136–137

prickly pear 315–316  
puncturevine 171–172  
rabbitbrush 138–139  
rabbitbrush, green 140–141  
red stem peavine 257–258  
rubberweed, bitter 281–282  
sagebrush, big 144–145  
sagebrush, Bigelow 146–147  
sagebrush, black 148–149  
sagebrush, sand 152–153  
saltgrass 81–82  
sego lily 283–284  
senecio 285–286  
shadscale 154–155  
skunkbush 156–157  
snakeweed 158–159  
snowberry 160–161  
thistle, Canada 299–300  
thistle, musk 297–298  
three-awns (perennial) 87–88  
tumbleweed 303–304  
wax currant 162–163  
western yarrow 307  
wheatgrass, crested 91–92  
wild rose 164–165  
wildrye, Russian 99–100  
winterfat 166–167  
wolfberry 168–169  
wooly Indian wheat 310–311  
yucca 317–318

#### Grasslands

antelope bitterbrush 112–113  
antelopehorn 173–174  
arrowgrass 101  
birdbeak 177–178  
bitterweed 179–180  
blanket flower 181–182  
bluebells 185–186  
blue-eyed grass 183–184  
bluegrass, Kentucky 10–11  
bluestem, little false 12  
bottlebrush squirreltail 13  
brome, field 14–15  
brome, Japanese 16–17  
brome, mountain 18–19  
brome, red 20–21  
brome, rigput 22–23  
brome, smooth 24–25  
bulrush 102–103  
bundle flower 187–188  
California burclover 189–190  
cheatgrass 26–27

cholla 313–314  
cordgrass, alkali 28–29  
cow parsnip 196–197  
curlycup gumweed 198–199  
daisy, rock 200–201  
dandelion 202–203  
desert sand verbena 206  
devilweed 207–208  
dropseed, giant 30–31  
dropseed, pine 32–33  
dropseed, sand 34–35  
dropseed, spike 36–37  
fescue, Arizona 40–41  
fescue, six weeks 42–43  
field sandbur 83–84  
filaree 211–212  
foxtail barley 44–45  
geranium 215–216  
gilia 217–218  
grama, black 50–51  
grama, blue 52–53  
grama, sideoats 54–55  
greasewood 130–131  
groundsel, threadleaf 221–222  
horsetail 352–353  
Indian paintbrush 227–228  
Jacob's ladder 231–232  
James' galleta 46–47  
jimsonweed 235–236  
johnsongrass 56–57  
jointed goatgrass 48–49  
junegrass, prairie 58–59  
knapweed, diffuse 239–240  
knapweed, Russian 241–242  
kochia 243–244  
larkspur 247–248  
leafy spurge 249–250  
loco, rattleweed 253–254  
loco, wooly 255–256  
milkweed 261–262  
muhly, mountain 60–61  
muhly, ring 62–63  
muhly, sandhill 64–65  
muhly, spike 66–67  
mustard 265–266  
mutton grass 68  
Navajo tobacco 269–270  
Navajo wild carrot 271–272  
needle-and-threadgrass 69–70  
needlegrass, desert 71–72  
New Mexico feathergrass 75–76  
nightshade 273–274

## Index

*continued*

### Habitat

#### Grasslands *continued*

penstemon 275–276  
pepperweed 277–278  
poison ivy 279–280  
prickly pear 315–316  
red stem peavine 257–258  
redtop 77–78  
ricegrass, Indian 79–80  
rubberweed, bitter 281–282  
rushes 104–105  
sacaton, alkali 38–39  
sagebrush, bigelow 146–147  
sagebrush, fringed 150–151  
saltgrass 81–82  
scarlet globemallow 219–220  
sedges 108–109  
senecio 285–286  
shadscale 154–155  
sneezeweed 287–288  
spreading wallflower 289–290  
St. Johnswort 291–292  
sunflower 295–296  
thistle, Canada 299–300  
three-awns (annual) 85–86  
three-awns (perennial) 87–88  
tufted horse grass 89–90  
tumbleweed 303–304  
wheatgrass, crested 91–92  
wheatgrass, western 95–96  
whitetop 308–309  
wild rose 164–165  
wildrye, Canadian 97–98  
wooly Indian wheat 310–311

#### Piñon-Juniper Woodland

birdbeak 177–178  
bitterweed 179–180  
blackbrush 116–117  
cliff fendlerbush 125–126  
cliffrose 127  
Colorado piñon 321–322  
daisy, rock 200–201  
dropseed, giant 30–31  
dropseed, pine 32–33  
evening primrose 209–210  
fescue, six weeks 42–43  
foxtail barley 44–45  
geranium 215–216  
gilia 217–218  
James' galleta 46–47  
jointed goatgrass 48–49  
juniper 328–329  
knapweed, diffuse 239–240

loco, Ellis 251–252  
loco, rattleweed 253–254  
mountain mahogany 330–331  
muhly, spike 66–67  
Navajo tea 267–268  
needlegrass, Scribner 73  
prickly pear 315–316  
red stem peavine 257–258  
roundleaf buffaloberry 142–143  
saltgrass 81–82  
thistle, Canada 299–300  
thistle, Scotch 301–302  
wild rose 164–165  
wildrye, Russian 99–100  
Ponderosa Pine/Mixed Conifer  
bitterweed 179–180  
bluestem, little false 12  
daisy, rock 200–201  
devilweed 207–208  
dropseed, pine 32–33  
fescue, six weeks 42–43  
foxtail barley 44–45  
Gambel oak 327  
geranium 215–216  
Jacob's ladder 231–232  
James' galleta 46–47  
knapweed, diffuse 239–240  
loco, Ellis 251–252  
muhly, mountain 60–61  
mustard 265–266  
needlegrass, Scribner 73  
ponderosa pine 332–333  
prickly pear 315–316  
quaking aspen 334–335  
roundleaf buffaloberry 142–143  
sedges 108–109  
sego lily 283–284  
snowberry 160–161  
subalpine fir 344–345  
thistle, Canada 299–300  
wax currant 162–163  
wild rose 164–165  
wildrye, Canadian 97–98  
River Valleys  
alder 319–320  
bitterweed 179–180  
brome, field 14–15  
bulrush 102–103  
copperweed 195  
coyote willow 323–324  
dandelion 202–203

dropseed, giant 30–31  
foxtail barley 44–45  
Fremont cottonwood 325–326  
geranium 215–216  
gilia 217–218  
horsemint 225–226  
horsetail 352–353  
horsetail, smooth 354–355  
iris 229–230  
kochia 243–244  
larkspur 247–248  
leafy spurge 249–250  
lupine 259–260  
Navajo wild carrot 271–272  
nutsedge, yellow 106–107  
pepperweed 277–278  
poison ivy 279–280  
quaking aspen 334–335  
red plum 336–337  
rubberweed, bitter 281–282  
rushes 104–105  
Russian olive 338–339  
sagebrush, black 148–149  
saltcedar 340–341  
saltgrass 81–82  
sedges 108–109  
Siberian elm 342–343  
snowberry 160–161  
sunflower 295–296  
tufted horse grass 89–90  
Utah serviceberry 348–349  
water birch 350–351  
water hemlock 305–306  
wheatgrass, slender 93–94  
whitetop 308–309  
wild rose 164–165  
Wastelands  
beeplant, Rocky Mountain 175–176  
bottlebrush squirreltail 13  
brome, Japanese 16–17  
brome, ripgut 22–23  
cholla 313–314  
cliff fendlerbush 125–126  
cliffrose 127  
copperweed 195  
death camas 204–205  
desert sand verbena 206  
dropseed, spike 36–37  
filaree 211–212  
foxtail barley 44–45

## Index

*continued*

### Habitat

#### Wastelands *continued*

grama, black 50–51  
groundsel, threadleaf 221–222  
Indian paintbrush 227–228  
James' galleta 46–47  
jimsonweed 235–236  
kochia 243–244  
lambsquarters 245–246  
leafy spurge 249–250

loco, rattleweed 253–254  
milkweed 261–262  
Mormon tea 132–133  
mullein 263–264  
Navajo tobacco 269–270  
needlegrass, desert 71–72  
nightshade 273–274  
prickly pear 315–316

sagebrush, black 148–149  
sneezeweed 287–288  
tumbleweed 303–304  
wheatgrass, crested 91–92  
winterfat 166–167  
wolfberry 168–169  
wormsweed 312  
yucca 317–318

### Plant Type

#### Brush

algerita 110–111  
antelope bitterbrush 112–113  
biscuitroot 114–115  
blackbrush 116–117  
buckwheat 118–120  
camelthorn 121–122  
chokecherry 123–124  
cliff fendlerbush 125–126  
cliffrose 127  
four-winged saltbush 128–129  
greasewood 130–131  
Mormon tea 132–133  
Oregon grape 134–135  
pingue 136–137  
rabbitbrush 138–139  
rabbitbrush, green 140–141  
roundleaf buffaloberry 142–143  
sagebrush, big 144–145  
sagebrush, Bigelow 146–147  
sagebrush, black 148–149  
sagebrush, fringed 150–151  
sagebrush, sand 152–153  
shadscale 154–155  
skunkbush 156–157  
snakeweed 158–159  
snowberry 160–161  
thistle, Scotch 301–302  
wax currant 162–163  
wild rose 164–165  
winterfat 166–167  
wolfberry 168–169

#### Cacti

cholla 313–314  
prickly pear 315–316  
yucca 317–318

#### Forbs

antelopehorn 173–174  
beepplant, Rocky Mountain 175–176  
birdbeak 177–178  
bitterweed 179–180  
blanket flower 181–182  
bluebells 185–186  
blue-eyed grass 183–184  
bundle flower 187–188  
California burclover 189–190  
cocklebur 191–192  
common purslane 193–194  
copperweed 195  
cow parsnip 196–197  
curlycup gumweed 198–199  
daisy, rock 200–201  
dandelion 202–203  
death camas 204–205  
desert sand verbena 206  
devilweed 207–208  
evening primrose 209–210  
filaree 211–212  
four o'clock 213–214  
geranium 215–216  
gilia 217–218  
groundsel, threadleaf 221–222  
halogeton 223–224  
horsemint 225–226  
Indian paintbrush 227–228  
iris 229–230  
Jacob's ladder 231–232  
jimmyweed 233–234  
jimsonweed 235–236  
knapweed 237–238  
knapweed, diffuse 239–240

knapweed, Russian 241–242  
kochia 243–244  
lambsquarters 245–246  
larkspur 247–248  
leafy spurge 249–250  
loco, Ellis 251–252  
loco, rattleweed 253–254  
loco, woolly 255–256  
lupine 259–260  
milkweed 261–262  
mullein 263–264  
mustard 265–266  
Navajo tea 267–268  
Navajo wild carrot 271–272  
nightshade 273–274  
penstemon 275–276  
pepperweed 277–278  
poison ivy 279–280  
red stem peavine 257–258  
rubberweed 281–282  
scarlet globemallow 219–220  
sego lily 283–284  
senecio 285–286  
sneezeweed 287–288  
spreading wallflower 289–290  
St. Johnswort 291–292  
sulphur cinquefoil 293–294  
sunflower 295–296  
thistle, Canada 299–300  
thistle, musk 297–298  
thistle, Scotch 301–302  
tumbleweed 303–304  
water hemlock 305–306  
western yarrow 307  
whitetop 308–309  
wooly Indian wheat 310–311



## Index

continued

### Plant Type

Forbs *continued*

wormsweed 312

Grass

bluegrass, Kentucky 10–11

bluestem, little false 12

bottlebrush squirreltail 13

brome, field 14–15

brome, Japanese 16–17

brome, mountain 18–19

brome, red 20–21

brome, ripgut 22–23

brome, smooth 24–25

cheatgrass 26–27

cordgrass, alkali 28–29

dropseed, giant 30–31

dropseed, pine 32–33

dropseed, sand 34–35

dropseed, spike 36–37

fescue, Arizona 40–41

fescue, six weeks 42–43

field sandbur 83–84

foxtail barley 44–45

grama, black 50–51

grama, blue 52–53

grama, sideoats 54–55

Grasslike

arrowgrass 101

bulrush 102–103

nutsedge, yellow 106–107

rushes 104–105

sedges 108–109

James' galleta 46–47

johnsongrass 56–57

jointed goatgrass 48–49

junegrass, prairie 58–59

knapweed, Russian 241–242

muhly, mountain 60–61

muhly, ring 62–63

muhly, sandhill 64–65

muhly, spike 66–67

mutton grass 68

Navajo tobacco 269–270

needle-and-threadgrass 69–70

needlegrass, desert 71–72

needlegrass, Scribner 73

New Mexico feathergrass 75–76

redtop 77–78

ricegrass, Indian 79–80

sacaton, alkali 38–39

saltgrass 81–82

three-awns (annual) 85–86

three-awns (perennial) 87–88

tufted horse grass 89–90

wheatgrass, crested 91–92

wheatgrass, slender 93–94

wheatgrass, western 95–96

wildrye, Canadian 97–98

wildrye, Russian 99–100

None

horsetail 352–353

horsetail, smooth 354–355

lichen 356–357

Trees

alder 319–320

Colorado piñon 321–322

coyote willow 323–324

Fremont cottonwood 325–326

Gambel oak 327

juniper 328–329

mountain mahogany 330–331

ponderosa pine 332–333

quaking aspen 334–335

red plum 336–337

Russian olive 338–339

saltcedar 340–341

Siberian elm 342–343

subalpine fir 344–345

tree of heaven 346–347

Utah serviceberry 348–349

water birch 350–351

Vines

field bindweed 170

puncturevine 171–172

### Scientific Name

*Abies lasiocarpa* subsp. *Arizona*  
344–345

*Abronia* spp. 206

*Achillea lanulosa* 307

*Achnatherum hymenoides* 79–80

*Achnatherum scribneri* 73–74

*Acroptilon repens* 241–242

*Aegilops cylindrica* 48–49

*Agropyron smithii* 95–96

*Agrostis gigantea* 77–78

*Agropyron cristatum* 91–92

*Ailanthus altissima* 346–347

*Alhagi camelorum* 121–122

*Alhagi maurorum* 121–122

*Allionia incarnata* 213–214

*Alnus* spp. 319–320

*Amelanchier utahensis* 348–349

*Aristida* spp. 85–86, 85–88

*Artemisia bigelovii* 146–147

*Artemisia filifolia* 152–153

*Artemisia frigida* 150–151

*Artemisia nova* 148–149

*Artemisia tridentata* 144–145

*Asclepias asperula* 173–174

*Asclepias* spp. 261–262

*Aster spinosus* 207–208

*Astragalus allochrous* 253–254

*Astragalus amphioxys* 251, 253,  
255, 258

*Astragalus beathii* 251, 253, 255,  
258

*Astragalus ceramicus* E. Sheldon var  
*ceramicus* 251, 253, 255, 258

*Astragalus cronquistii* 251, 253,  
255, 258

*Astragalus cutleri* 251, 253, 255,  
258

*Astragalus emoryanus* 257–258

*Astragalus heilii* 251, 253, 255, 258

*Astragalus humillimus* 251, 253,  
255, 258

*Astragalus humistratus* 251, 253,  
255, 258

## Index

continued

### Scientific Name *continued*

- Astragalus lentiginosus* Douglas ex Hooker var *australis* Barnely 251, 253, 255, 258  
*Astragalus lonchocarpus* 251, 253, 255, 258  
*Astragalus missouriensis* 251, 253, 255, 258  
*Astragalus mollisium* 251, 253, 255, 258  
*Astragalus mollissimus* 255–256  
*Astragalus naturitensis* 251, 253, 255, 258  
*Astragalus nuttallianus* 251, 253, 255, 257, 258  
*Astragalus praelongus* Sheldon var *ellisiae* 251, 253, 255, 258  
*Astragalus praelongus* var. *ellisiae* 251–252  
*Atriplex canescens* 128–129  
*Atriplex confertifolia* 154–155  
*Atriplex garrettii* var. *navajoensis* 128  
*Bassia scoparia* 243–244  
*Berberis aquifolium* 134–135  
*Berberis trifoliolata* 110–111  
*Betula occidentalis* 350–351  
*Blepharoneuron tricholepis* 32–33  
*Bouteloua curtipendula* 54–55  
*Bouteloua eriopoda* 50–51  
*Bouteloua gracilis* 52–53  
*Brassica* spp. 265–266  
*Bromus arvensis* 14–15  
*Bromus carinatus* 18–19  
*Bromus diandrus* 22–23  
*Bromus inermis* 24–25  
*Bromus japonicus* 16–17  
*Bromus rubens* 20–21  
*Bromus tectorum* 26–27  
*Calochortus nuttallii* 283–284  
*Campanula rotundifolia* 185–186  
*Cardaria draba* 308–309  
*Carduus nutans* 297–298  
*Carex specuicola* 108  
*Carex* spp. 108–109  
*Castilleja* spp. 227–228  
*Cenchrus incertus* 14–15, 83–84  
*Cenchrus spinifex* 83–84  
*Centaurea diffusa* 239–240  
*Centaurea picris* 241–242  
*Centaurea* spp. 237–238  
*Ceratoides lanata* 166  
*Cercocarpus montanus* 330–331  
*Chenopodium album* var. *missouriense* 245–246  
*Chenopodium missouriense* Aellen 245–246  
*Chloracantha spinosa* 207–208  
*Chrysothamnus viscidiflorus* 140–141  
*Cicuta douglasii* 305–306  
*Cirsium arvense* 299–300  
*Cleome serrulata* 175–176  
*Coleogyne ramosissima* 116–117  
*Convolvulus arvensis* 170  
*Cordylanthus wrightii* 177–178  
*Cowania neomexicana* 127  
*Cylindropuntia imbricata* 313  
*Cylindropuntia* spp. 313–314  
*Cylindropuntia versicolor* 313  
*Cylindropuntia whipplei* 313  
*Cyperus esculentus* 106–107  
*Datura* spp. 235–236  
*Delphinium* spp. 247–248  
*Deschampsia cespitosa* 89–90  
*Desmanthus cooleyi* 187–188  
*Distichlis spicata* 81–82  
*Dysphania ambrosioides* 312  
*Elaeagnus angustifolia* 338–339  
*Elymus canadensis* 97–98  
*Elymus elymoides* 13  
*Elymus trachycaulus* 93–94  
*Ephedra* spp. 132–133  
*Equisetum arvense* 352–353  
*Equisetum laevigatum* 354–355  
*Ericameria arizonica* 138  
*Ericameria nauseosa* 138–139  
*Eriogonum alatum* 119  
*Eriogonum cernuum* 119  
*Eriogonum clavellatum* 119  
*Eriogonum corymbosum* 119  
*Eriogonum deflexum* 119  
*Eriogonum divaricatum* 119  
*Eriogonum effusum* 119  
*Eriogonum gordonii* 119  
*Eriogonum hieraciifolium* 119  
*Eriogonum hookeri* 119  
*Eriogonum inflatum* 119  
*Eriogonum jamesii* 119  
*Eriogonum lachnogynum* 119  
*Eriogonum lachnogynum* var. *sarahiae* 119  
*Eriogonum leptocladon* 119  
*Eriogonum leptophyllum* 119  
*Eriogonum lonchophyllum* 119  
*Eriogonum microthecum* 118–120  
*Eriogonum ovalifolium* 119  
*Eriogonum pharnaceoides* 119  
*Eriogonum racemosum* 119  
*Eriogonum scabrellum* 119  
*Eriogonum shockleyi* 119  
*Eriogonum* spp. 118–120  
*Eriogonum subreniforme* 119  
*Eriogonum umbellatum* 119  
*Eriogonum viridulum* 119  
*Eriogonum wetherillii* 119  
*Eriogonum wrightii* 119  
*Erodium cicutarium* 211–212  
*Erysimum repandum* 289–290  
*Euphorbia esula* 249–250  
*Eurotia ceratoides* 166–167  
*Eurotia lanata* 166–167  
*Fendlera rupicola* 125–126  
*Festuca arizonica* 40–41  
*Gaillardia* spp. 181–182  
*Geranium* spp. 215–216  
*Gilia* spp. 217–218  
*Grindelia squarrosa* 198–199  
*Gutierrezia sarothrae* 158–159  
*Halogeton glomeratus* 223–224  
*Haplopappus heterophyllus* 233–234  
*Haplopappus pluriflorus* 233–234  
*Helenium autumnale* 287–288  
*Helianthus annuus* 295–296  
*Helianthus anomalus* 295  
*Helianthus maximiliani* 295  
*Helianthus nuttallii* 295  
*Helianthus petiolaris* 295  
*Heracleum maximum* 196–197  
*Hesperostipa comata* 69–70  
*Hesperostipa comata* subsp. *comata* 69–70  
*Hesperostipa neomexicana* 75–76  
*Hilaria jamesii* 46–47  
*Hordeum jubatum* 44–45  
*Hymenoxys odorata* 180, 281–282  
*Hymenoxys richardsonii* 136–137  
*Hymenoxys* spp. 179–180  
*Hypericum perforatum* 291–292  
*Iris missouriensis* 229–230  
*Isocoma pluriflora* 233–234  
*Isocoma wrightii* 233–234  
*Juncus arcticus* 104  
*Juncus bufonius* 104  
*Juncus ensifolius* 104

## Index

continued

### Scientific Name *continued*

- Juncus interior* 104  
*Juncus longistylis* 104  
*Juncus* spp. 104–105  
*Juncus torreyi* 104  
*Juniperus monosperma* 328–329  
*Kochia scoparia* 243–244  
*Koeleria macrantha* 58–59  
*Krascheninnikovia lanata* 166–167  
*Lepidium alyssoides* 277–278  
*Lepidium draba* 308–309  
*Lepidium virginicum* 277–278  
*Lichen* 356–357  
*Lomatium* spp. 114–115  
*Lupinus* spp. 259–260  
*Lycium* spp. 168–169  
*Mahonia aquifolium* 134–135  
*Medicago polymorpha* 189–190  
*Monarda* spp. 225–226  
*Muhlenbergia montana* 60–61  
*Muhlenbergia pungens* 64–65  
*Muhlenbergia torreyi* 62–63  
*Muhlenbergia wrightii* 66–67  
*Oenothera* spp. 209–210  
*Onopordum acanthium* 301–302  
*Opuntia chlorotica* 315  
*Opuntia engelmannii* 315  
*Opuntia macrocentra* 315  
*Opuntia macrorhiza* var. *pottsii* 315–316  
*Opuntia phaeacantha* 315  
*Opuntia polyacantha* 315–316  
*Opuntia* spp. 315–316  
*Oryzopsis hymenoides* 79–80  
*Oxytenia acerosa* 195  
*Oxytropis lambertii* 251, 253, 255, 258  
*Pappostipa speciosa* 71–72  
*Pascopyrum smithii* 95–96  
*Penstemon navajoa* 275  
*Penstemon* spp. 275–276  
*Penstemon strictus* 275–276  
*Perityle emoryi* 200–201  
*Pinus edulis* 321–322  
*Pinus ponderosa* 332–333  
*Plantago patagonica* 310–311  
*Pleuraphis jamesii* 46–47  
*Poa fendleriana* 68  
*Poa pratensis* 10–11  
*Polemonium foliosissimum* 231–232  
*Populus fremontii* 325–326  
*Populus tremuloides* 334–335  
*Portulaca oleracea* 193–194  
*Potentilla recta* 293–294  
*Prunus americana* 336–337  
*Prunus virginiana* 123–124  
*Psathyrostachys juncea* 99–100  
*Purshia mexicana* 127  
*Purshia tridentata* 112–113  
*Quercus gambelii* 327  
*Rhaponticum repens* 241–242  
*Rhus trilobata* 156–157  
*Ribes cereum* 162–163  
*Rosa woodsii* 164–165  
*Rumex crispus* 269–270  
*Rumex hymenosepalus* 271–272  
*Salix bebbiana* 162  
*Salix exigua* 323–324  
*Salsola kali* 303–304  
*Salsola tragus* 303–304  
*Sarcobatus vermiculatus* 130–131  
*Schizachyrium scoparium* var. *sco-*  
*parium* 12  
*Schoenoplectus acutus* 102–103  
*Senecio flaccidus* var. *flaccidus* 221–222  
*Senecio longilobus* 221–222  
*Senecio* spp. 285–286  
*Senecio vulgaris* 285  
*Senecio warnockii* 286  
*Shepherdia rotundifolia* 142–143  
*Sisyrinchium* spp. 183–184  
*Solanum* spp. 273–274  
*Sorghum halapense* 57  
*Sorghum x alnum* 56–57  
*Spartina gracilis* 28–29  
*Sphaeralcea coccinea* 219–220  
*Sphaeralcea* spp. 219–220  
*Sporobolus airoides* 38–39  
*Sporobolus contractus* 36–37  
*Sporobolus cryptandrus* 34–35  
*Sporobolus giganteus* 30–31  
*Stipa comata* 69–70  
*Stipa cometa* 69–70  
*Stipa hymenoides* 79–80  
*Stipa neomexicana* 75–76  
*Stipa speciosa* 71–72  
*Symphoricarpos albus* 160–161  
*Symphoricarpos* spp. 160–161  
*Tamarix ramosissima* 341  
*Tamarix* spp. 340–341  
*Taraxacum officinale* 202–203  
*Thelesperma megapotamicum* 267–268  
*Toxicodendron rydbergii* 279–280  
*Tribulus terrestris* 171–172  
*Triglochin maritima* 101  
*Ulmus pumila* 342–343  
*Verbascum thapsus* 263–264  
*Vulpia octoflora* 42–43  
*Xanthium saccharatum* 191–192  
*Yucca angustissima* 317–318  
*Yucca baccata* 317–318  
*Yucca baileyi* 318  
*Yucca* spp. 317–318  
*Zigadenus paniculatus* 204–205  
*Zigadenus vaginatus* 204

### Special Considerations

- Harmful or invasive  
algerita 110–111  
antelopehorn 173–174  
arrowgrass 101  
bitterweed 179–180  
bluegrass, Kentucky 10–11  
brome, field 14–15  
brome, Japanese 16–17  
brome, red 20–21  
brome, ripgut 22–23  
brome, smooth 24–25  
bulrush 102–103  
California burclover 189–190  
camelthorn 121–122  
cheatgrass 26–27



## Index

*continued*

### Special Considerations

Harmful or invasive *continued*

chokecherry 123–124  
cholla 313–314  
cocklebur 191–192  
common purslane 193–194  
copperweed 195  
cow parsnip 196–197  
death camas 204–205  
field sandbur 83–84  
filaree 211–212  
foxtail barley 44–45  
Gambel oak 327  
greasewood 130–131  
groundsel, threadleaf 221–222  
halogeton 223–224  
horsetail 352–353  
jimmyweed 233–234  
jimsonweed 235–236  
johnsongrass 56–57  
jointed goatgrass 48–49  
knapweed 237–238  
knapweed, diffuse 239–240  
knapweed, Russian 241–242  
kochia 243–244  
larkspur 247–248  
leafy spurge 249–250  
loco, Ellis 251–252  
loco, rattleweed 253–254  
loco, woolly 255–256  
lupine 259–260  
milkweed 261–262  
Mormon tea 132–133  
mustard 265–266  
Navajo tobacco 269–270  
needle-and-threadgrass 69–70  
needlegrass, desert 71–72  
New Mexico feathergrass 75–76  
nightshade 273–274  
nutsedge, yellow 106–107  
pepperweed 277–278  
pingue 136–137  
poison ivy 279–280  
ponderosa pine 332–333  
prickly pear 315–316  
puncturevine 171–172  
rabbitbrush, green 140–141  
red stem peavine 257–258  
redtop 77–78  
rubberweed, bitter 281–282  
Russian olive 338–339  
sagebrush, sand 152–153  
saltcedar 340–341

saltgrass 81–82  
senecio 285–286  
Siberian elm 342–343  
snakeweed 158–159  
sneezeweed 287–288  
spreading wallflower 289–290  
St. Johnswort 291–292  
sulphur cinquefoil 293–294  
thistle, Canada 299–300  
thistle, musk 297–298  
thistle, Scotch 301–302  
three-awns (annual) 85–86  
three-awns (perennial) 87–88  
tree of heaven 346–347  
tumbleweed 303–304  
water hemlock 305–306  
wheatgrass, crested 91–92  
whiteweed 308–309

#### Important

grama, black 50–51  
muhly, ring 62–63  
Navajo tea 267–268  
Navajo wild carrot 271–272

#### Particularly helpful on rangelands

algerita 110–111  
antelope bitterbrush 112–113  
birdbeak 177–178  
biscuitroot 114–115  
bluegrass, Kentucky 10–11  
bottlebrush squirreltail 13  
brome, mountain 18–19  
brome, smooth 24–25  
bulrush 102–103  
cliffrose 127  
Colorado piñon 321–322  
common purslane 193–194  
cordgrass, alkali 28–29  
cow parsnip 196–197  
coyote willow 323–324  
dandelion 202–203  
dropseed, pine 32–33  
dropseed, spike 36–37  
filaree 211–212  
four-winged saltbush 128–129  
Gambel oak 327  
gilia 217–218  
grama, black 50–51  
grama, blue 52–53  
grama, sideoats 54–55  
horsemint 225–226

junegrass, prairie 58–59  
lichen 356–357  
milkweed 261–262  
mountain mahogany 330–331  
muhly, mountain 60–61  
muhly, sandhill 64–65  
muhly, spike 66–67  
mutton grass 68–69  
Navajo wild carrot 271–272  
needle-and-threadgrass 69–70  
New Mexico feathergrass 75–76  
penstemon 275–276  
ponderosa pine 332–333  
prickly pear 315–316  
quaking aspen 334–335  
rabbitbrush 138–139  
red plum 336–337  
redtop 77–78  
ricegrass, Indian 79–80  
roundleaf buffaloberry 142–143  
rushes 104–105  
sacaton, alkali 38–39  
sagebrush, big 144–145  
sagebrush, Bigelow 146–147  
sagebrush, sand 152–153  
saltcedar 340–341  
saltgrass 81–82  
scarlet globemallow 219–220  
shadscale 154–155  
sneezeweed 287–288  
snowberry 160–161  
St. Johnswort 291–292  
sunflower 295–296  
water birch 350–351  
wax currant 162–163  
western yarrow 307  
wheatgrass, crested 91–92  
wheatgrass, slender 93–94  
wheatgrass, western 95–96  
wild rose 164–165  
winterfat 166–167  
wormsweed 312  
yucca 317–318